

Table 1. Characteristics of participants, overall and stratified by double infection status

	All, N=81	MI, N=74	DI, N=7	p-value
Age (year), median (1st – 3rd quartiles)	32.9 [27.6;41.2]	33.1 [27.8;41.2]	32.0 [26.8;39.2]	0.775
Age (year), N (%)				1.000
20 - 29	32 (39.5%)	29 (39.2%)	3 (42.9%)	
30 - 39	26 (32.1%)	24 (32.4%)	2 (28.6%)	
≥ 40	23 (28.4%)	21 (28.4%)	2 (28.6%)	
Country, N (%)				0.188
Spain	56 (69.1%)	53 (71.6%)	3 (42.9%)	
Other	25 (30.9%)	21 (28.4%)	4 (57.1%)	
Time since HIV infection (months), median (1st – 3rd quartiles)	12.9 [2.23;38.9]	11.1 [2.23;39.0]	25.7 [11.2;36.7]	0.347
Previous STD, N (%)				0.660
No	23 (28.4%)	22 (29.7%)	1 (14.3%)	
Yes	58 (71.6%)	52 (70.3%)	6 (85.7%)	
STD since HIV Diagnosis, N (%)				1.000
No	16 (19.8%)	15 (20.3%)	1 (14.3%)	
Yes	65 (80.2%)	59 (79.7%)	6 (85.7%)	
N of STD, N (%)				0.283
None	14 (17.5%)	13 (17.8%)	1 (14.3%)	
1 - 2	33 (40.0%)	32 (42.5%)	1 (14.3%)	
≥ 3	34 (42.5%)	29 (39.7%)	5 (71.4%)	
N of SP lifetime, N (%)				0.098
0 - 500	49 (60.5%)	47 (63.5%)	2 (28.6%)	
> 501	32 (39.5%)	27 (36.5%)	5 (71.4%)	
N of SP last year, N (%)				0.017
≤25	47 (58.0%)	46 (62.2%)	1 (14.3%)	
>25	34 (42.0%)	28 (37.8%)	6 (85.7%)	
Viral load (copies/mL), N (%)				0.246
0 - 999	25 (30.9%)	21 (28.4%)	4 (57.1%)	
1000 - 4999	30 (37.0%)	29 (39.2%)	1 (14.3%)	
≥5000	26 (32.1%)	24 (32.4%)	2 (28.6%)	
CD4+ T cells/μL, N (%)				1.000
≤500	18 (22.5%)	17 (23.3%)	1 (14.3%)	

M.I: Mono-infected individuals. D.I: Double-infected individuals. STD: Sexually transmitted disease  
 SP: Sexual partner