

This is the peer reviewed version of the following article:

Low adherence to the western and high adherence to the mediterranean dietary patterns could prevent colorectal cancer

Adela Castelló, Pilar Amiano, Nerea Fernández de Larrea, Vicente Martín, Maria Henar Alonso, Gemma Castaño-Vinyals, Beatriz Pérez-Gómez, Rocío Olmedo-Requena, Marcela Guevara, Guillermo Fernandez-Tardon, Trinidad Dierssen-Sotos, Cristobal Llorens-Ivorra, Jose María Huerta, Rocío Capelo, Tania Fernández-Villa, Anna Díez-Villanueva, Carmen Urtiaga, Jesús Castilla, Jose Juan Jiménez-Moleón, Víctor Moreno, Verónica Dávila-Batista, Manolis Kogevinas, Nuria Aragonés, Marina Pollán, On behalf of MCC-Spain researchers

[Eur J Nutr.](#) 2019 Jun;58(4):1495-1505.

which has been published in final form at

<https://doi.org/10.1007/s00394-018-1674-5>

1 **AUTHORS LIST:** Adela Castelló^{1,2,3}, Pilar Amiano^{2,4,5}, Nerea Fernández de Larrea^{1,2}, Vicente
2 Martín⁶, Maria Henar Alonso^{7,8}, Gemma Castaño-Vinyals^{2,9,10,11}, Beatriz Pérez-Gómez^{1,2},
3 Rocío Olmedo-Requena^{2,12,13}, Marcela Guevara^{2,14}, Guillermo Fernandez-Tardon^{2,15}, Trinidad
4 Dierssen-Sotos¹⁶, Cristobal Llorens-Ivorra¹⁷, Jose M. Huerta^{2,18}, Rocío Capelo¹⁹, Tania
5 Fernández-Villa⁶, Anna Díez-Villanueva⁷, Carmen Urtiaga⁵, Jesús Castilla^{2,14}, Jose J. Jiménez-
6 Moleón^{2,12,13}, Víctor Moreno^{7,8}, Verónica Dávila-Batista⁶, Manolis Kogevinas^{2,9,10,11}, Nuria
7 Aragonés^{1,2}, Marina Pollán^{1,2} on behalf of MCC-Spain researchers

8 **TITLE: Low adherence to the Western and high adherence to the Mediterranean dietary**
9 **patterns could prevent colorectal cancer.**

10 **AFFILIATIONS:**

11 ¹ Cancer Epidemiology Unit, National Center for Epidemiology, Carlos III Institute of Health.
12 Avenida Monforte de Lemos 5, 28029, Madrid, Spain.

13 ² Consortium for Biomedical Research in Epidemiology & Public Health (CIBERESP), Carlos
14 III Institute of Health. Avenida Monforte de Lemos 5, 28029, Madrid, Spain.

15 ³ Faculty of Medicine, University of Alcalá, Campus Universitario - C/ 19, Av. de Madrid,
16 Km 33,600, 28871 Alcalá de Henares, Madrid

17 ⁴ Public Health Department of Gipuzkoa, Government of the Basque Country, Avenida
18 Navarra, 4, 20013, San Sebastián, Spain.

19 ⁵ Biodonostia Research Institute. Paseo Dr Beguiristain s/n, 20014, San Sebastián, Spain.

20 ⁶ The Research Group in Gene-Environment and Health Interactions, Vegazana Campus,
21 University of León. Campus Vegazana, s/n, 24071, León, Spain

22 ⁷ Unit of Biomarkers and Susceptibility, Cancer Prevention and Control Program, Catalan
23 Institute of Oncology (ICO) and IDIBELL. Gran Via km 2.7, 08907, L'Hospitalet de Llobregat,
24 Spain.

25 ⁸ Department of Clinical Sciences, Faculty of Medicine, University of Barcelona, Campus de
26 Bellvitge, Pavelló de Govern, Feixa Llarga s/n 08907, L'Hospitalet de Llobregat, Spain.

27 ⁹ ISGlobal, Centre for Research in Environmental Epidemiology (CREAL). Carrer del Doctor
28 Aiguader 88, 08003, Barcelona, Spain.

29 ¹⁰ Universitat Pompeu Fabra (UPF). Carrer del Doctor Aiguader 88, 08003, Barcelona, Spain

30 ¹¹ IMIM (Hospital del Mar Medical Research Institute). Carrer del Doctor Aiguader 88,
31 08003, Barcelona, Spain

32 ¹² Department of Preventive Medicine and Public Health, University of Granada. Av, de la
33 Investigación, 11, 18016, Granada, Spain.

34 ¹³ Instituto de Investigación Biosanitaria ibs.GRANADA, Complejo Hospitales Universitarios
35 de Granada/Universidad de Granada. Edificio Licinio de la Fuente, Calle Dr. Azpitarte, 4,
36 18012, Granada, Spain.

37 ¹⁴ Public Health Institute of Navarra. Calle Leyre 15, 31003, Pamplona, Spain

38 ¹⁵ IUOPA, University of Oviedo. Facultad de Medicina, Planta 7, Campus de El Cristo B,
39 33006, Oviedo, Spain.

40 ¹⁶ Universidad de Cantabria – IDIVAL. Avenida Cardenal Herrera Oria s/n, 39011,
41 Santander, Spain.

42 ¹⁷ Centro de Salud Pública de Dénia. Consellería de Sanidad Universal y Salud Pública. Plaza
43 Jaime I, 5, 03700, Denia, Spain.

44 ¹⁸ Department of Epidemiology, Murcia Regional Health Council, IMIB-Arrixaca. C/ Luis
45 Fontes Pagán nº 9 - 1ª planta. C.P.- 30003, Murcia, Spain.

46 ¹⁹ Centro de Investigación en Salud y Medio Ambiente (CYSMA). Universidad de Huelva.
47 Campus Universitario de El Carmen, 21071, Huelva, Spain.

48 **CORRESPONDING AUTHOR:**

49 Dr. Adela Castelló

50 Cancer Epidemiology Unit, National Center for Epidemiology.

51 Instituto de Salud Carlos III.

52 Avenida Monforte de Lemos, 5, 28029, Madrid, Spain.

53 ORCID: 0000-0002-1308-9927

54 Phone: +34 91 822 2667.

55 Fax: +34 91 387 7815.

56 e-mail: acastello@isciii.es

57

58 **ABSTRACT**

59 **Purpose:** To assess if the associations found between three previously identified dietary
60 patterns with breast, prostate and gastric cancer are also observed for colorectal cancer (CRC).

61 **Methods:** MCC-Spain is a multicase-control study that collected information of 1629 incident
62 cases of CRC and 3509 population-based controls from 11 Spanish provinces. Western, Prudent
63 and Mediterranean data-driven dietary patterns - derived in another Spanish case-control study-
64 were reconstructed in MCC-Spain. Their association with CRC was assessed using mixed
65 multivariable logistic regression models considering a possible interaction with sex. Risk by
66 tumor site (proximal colon, distal colon, and rectum) was evaluated using multinomial
67 regression models.

68 **Results:** While no effect of the Prudent pattern on CRC risk was observed, a high adherence to
69 the Western dietary pattern was associated with increased CRC risk for both males (OR_{fourth}
70 $(Q4)_{\text{vs.first}}(Q1)_{\text{quartile}}(95\%CI):1.45(1.11;1.91)$) and females ($OR_{Q4vsQ1}(95\%CI):1.50(1.07;2.09)$)
71 but seem to be confined to distal colon ($OR_{\text{fourth}}(Q4)_{\text{vs.first}}(Q1)_{\text{quartile}}(95\%CI):2.02(1.44;2.84)$) and
72 rectal ($OR_{Q4vsQ1}(95\%CI):1.46(1.05;2.01)$) tumors. The protective effect of the Mediterranean
73 dietary pattern against CRC was observed for both sexes (Males:
74 $OR_{Q4vsQ1}(95\%CI):0.71(0.55;0.92)$; Females: $OR_{Q4vsQ1}(95\%CI):0.56(0.40;0.77)$) and for all
75 cancer sites: proximal colon ($OR_{Q4vsQ1}(95\%CI):0.70(0.51;0.97)$), distal colon
76 ($OR_{Q4vsQ1}(95\%CI):0.65(0.48;0.89)$), and rectum ($OR_{Q4vsQ1}(95\%CI):0.60(0.45;0.81)$).

77 **Conclusion:** Our results are consistent with most of the associations previously found between
78 these patterns and breast, prostate and gastric cancer risk and indicate that consuming whole
79 fruits, vegetables, legumes, olive oil, nuts and fish and avoiding red and processed meat, refined
80 grains, sweets, caloric drinks, juices, convenience food and sauces might reduce CRC risk.

81 **KEYWORDS:** “Colonic Neoplasms”; “Rectal Neoplasms”; “prevention and control”;
82 “Principal Component Analysis”; “Dietary Patterns”; "Diet"; "Diet, Western"; "Diet,
83 Mediterranean".

84 **ACKNOWLEDGEMENTS:** This work was supported by Carlos III Institute of Health
85 grants (PI12/00488, PI12/00265, PI12/00715, PI12/01270, PI11/01403, PI11/01889,
86 PI11/00226, PI11/01810, PI11/02213, PI09/00773, PI09/01286, PI09/01903, PI09/02078,
87 PI09/01662, PI08/1770, PI08/0533, PI08/1359), Spanish Ministry of Economy and
88 Competitiveness (IJCI-2014-20900), Consejería de Salud de la Junta de Andalucía (PI-0306-
89 2011; PI-0571-2009); Catalan Government DURSGrant (2014SGR647); Instituto de Salud
90 Carlos III, co-funded by FEDER funds –a way to build Europe– PI14-00613; Fundación
91 Marqués de Valdecilla (API 10/09); Acción Transversal del Cancer, approved by the Spanish
92 Ministry Council on October 11, 2007; Red Temática de Investigación del Cáncer del ISCIII
93 (RD12/0036/0036); Junta de Castilla y León (LE22A10-2); Consejería de Salud de la Junta
94 de Andalucía (2009-S0143); Conselleria de Sanitat de la Generalitat Valenciana
95 (AP_061/10); Recercaixa (2010ACUP 00310); Regional Government of the Basque Country;
96 Consejería de Sanidad de la Región de Murcia; European Commission grants (FOOD-CT-
97 2006-036224-HIWATE); Spanish Association Against Cancer Scientific Foundation;
98 Fundación Caja de Ahorros de Asturias; University of Oviedo.

99 None of the sponsors intervened in any of the stages of the research.

100 **INTRODUCTION**

101 The incidence of colorectal cancer (CRC) has increased in Europe in the last decades
102 [1], being the second most diagnosed cancer in 2012 [2]. According to the scientific evidence,
103 40-50% of CRC cases are attributable to modifiable risk factors such as diet, physical activity
104 and body weight [3,4], providing major opportunities for prevention. The current evidence,
105 points to a possible protective effect of foods containing dietary fiber and calcium against CRC
106 [5,6] and a detrimental effect of red and processed meat [6,7] and alcohol consumption [6,8,9].

107 Despite the importance of these findings for individual foods, some authors suggest that
108 the evaluation of the effects of full dietary patterns might be more appropriate, since it allows
109 the exploration of the effect of food and nutrient interactions in disease [10-12]. Many indexes
110 have been developed in the last years that evaluate dietary quality against predefined criteria
111 [13,14] and a recent metaanalysis found an inverse association between a high score for these
112 indexes and cancer mortality and/or incidence [15]. However, these indexes are based on results
113 in the area of cardiovascular disease and they refer to a theoretical diet that do not necessarily
114 reflect the eating habits of a particular population. Moreover, moderate alcohol intake is
115 positively considered in most of these indexes although alcohol consumption as low as one
116 drink per day increases the risk of several tumors, including colorectal cancer [6]. In fact, some
117 authors suggested that the lack of concordance of the results found for diet quality indexes and
118 cancer might be due to their positive scoring for alcohol consumption [16]. As an alternative
119 approach, dietary patterns that accurately represent the diet in a specific population can be
120 identified with statistical methods like principal component analysis. These patterns also
121 present the advantage of being extracted independently of disease associations, which allows
122 exploration of the role of actual dietary habits in different health conditions. The scarce existing
123 results for data-driven dietary patterns and CRC, indicate a possible protective effect of the so
124 called Mediterranean/Healthy/Prudent dietary pattern [17-22] on CRC and a harmful effect of
125 a pattern labelled as Western/Unhealthy diet [17-23], but the evidence is still insufficient [6].

126 A recent Spanish study on female breast cancer (BC) –EpiGEICAM- identified three
127 data-driven dietary patterns [24] labelled as Western (associated with increased risk of BC),
128 Prudent (not associated with BC) and Mediterranean (protective against BC). EpiGEICAM
129 presents the novelty of being able to identify, over the same population, two different patterns
130 (Prudent and Mediterranean) commonly interchanged in the literature [9,18,20,21,23,25].
131 However, these patterns do not always represent the same dietary habits and the differences

132 might be determinant in their association with disease risk, as it was the case for BC in the
133 EpiGEICAM context [25]. Therefore, the replication of these patterns in different populations
134 and the exploration of their association with tumors other than BC are of great scientific interest.
135 In fact, the reproducibility of the patterns found has already been assessed in a different sample
136 of Spanish women [26] and similar associations have been observed for other tumours and
137 individuals. The detrimental effect of a high adherence to the Western dietary pattern has been
138 corroborated for breast cancer [27] and also observed for gastric [28] cancer. These studies also
139 show different results for the Prudent (null effect) and Mediterranean (protective) patterns in
140 the case of breast [27], prostate [29] and gastric cancer [28].

141 The objective of the present study is to assess if the associations found between these
142 dietary patterns -Western, Prudent and Mediterranean- with breast [24,27], prostate [29] and
143 gastric cancer [28] risk in our country are also observed for CRC risk, and to evaluate possible
144 differences by sex and cancer site.

145 **PATIENTS AND METHODS**

146 The multicase-control MCC-Spain study [30] recruited between 2008 and 2013
147 histologically confirmed incident cases of five tumors: breast, prostate, colorectal, stomach and
148 chronic lymphocytic leukemia. Cases were recruited in 23 hospitals from 12 provinces and a
149 single set of population controls, frequency matched by age and sex with the overall distribution
150 of cases in each province, were randomly selected from the lists of residents assigned to primary
151 health-care centers belonging to the same catchment area of each collaborating hospital. MCC-
152 Spain recruited 2140 histologically confirmed CRC cases and 3950 population-based controls
153 from 11 of the 12 contributing provinces. These numbers, represented the 64% of the CRC
154 cases and the 53% of controls invited to participate (supplementary material, figure S1).
155 Potential participants had to be able to answer the questionnaire, had to live in the study area for

156 at least 6 months before the diagnosis (cases) or at recruitment (controls) and had to be 20-85
157 years old. Cases were identified as soon as possible after the diagnosis, and histologically
158 confirmed incident cases of colon (ICD10 codes C18: malignant neoplasm of colon and
159 D01.0: Carcinoma in situ of colon) or rectum (ICD10 codes C19: Malignant neoplasm of
160 rectosigmoid junction ; C20: Malignant neoplasm of rectum; D01.1: Carcinoma in situ of
161 rectosigmoid junction and D01.2: Carcinoma in situ of rectum) cancer with no prior history of
162 the disease and diagnosed within the recruitment period were included. They were classified
163 according to the localization of tumor in proximal colon (including caecum, ascending &
164 transverse colon and hepatic and splenic flexures), distal colon (including descending and
165 sigmoid colon) and rectal cancer. When more than one tumor in different locations were
166 diagnosed at the same time, the site in which the tumor was more invasive was assigned.

167 Information on socio-demographic factors, lifestyle and personal/family medical history
168 was collected with a questionnaire administered by trained personnel in a face-to-face
169 interview. Subsequent telephone contact was made to complete missing values on key
170 variables. Height and weight at different ages were self-reported and diet was assessed with a
171 154-items semi-quantitative food frequency questionnaire (FFQ), which was based on an
172 instrument validated in Spain [31]. Dietary information referred to the previous year before
173 diagnosis (cases) or before interview (controls).

174 In the present study, three dietary patterns identified in a previous Spanish case-control
175 study (EpiGEICAM) on female breast cancer (BC) [24] are examined: A Western dietary
176 pattern positively associated with BC risk that is characterized by high intakes of high-fat dairy
177 products, processed meat, refined grains, sweets, caloric drinks, convenience food and sauces
178 and by low intakes of low-fat dairy products and whole grains; A Prudent pattern not related to
179 BC that represented high intakes of low-fat dairy products, vegetables, fruits, whole grains and
180 juices; and a Mediterranean pattern that seemed to be protective and denoted a high intake of

181 fish, vegetables, legumes, boiled potatoes, fruits, olives and vegetable oil – mainly olive oil
182 (72%), and olives (22%) in our context-, and a low intake of juices. These three dietary patterns
183 were identified in the EpiGEICAM sample in 2014 by grouping the dietary intake information
184 collected with a 117 FFQ into 26 inter-correlated food groups and applying principal
185 components analysis (PCA) without rotation of the variance-covariance matrix over these 26
186 food groups [32]. This method defines a set of weights (pattern loadings) associated with each
187 food group that represents the correlation between food consumption and the
188 component/pattern scores. Pattern loadings can be used to reproduce such patterns in other
189 samples as explained in detail elsewhere [25,26]. Briefly, we grouped 146 of the 154 items of
190 the MCC-Spain FFQ (excluding non-caloric and alcoholic beverages) into 26 food groups
191 defined in the EpiGEICAM study (see **Table 1** for detailed information on the composition of
192 food groups and their weight in the patterns). Afterwards, the scores of adherence to the
193 Western, Prudent and Mediterranean dietary patterns of the MCC-Spain participants were
194 calculated as a linear combination of the pattern loadings for each food group and dietary
195 pattern extracted from the EpiGEICAM study [24] (**Table 1**) and the food group consumption
196 reported by the MCC-Spain participants.

197 After describing the data, crude and adjusted associations between adherence to each
198 dietary pattern and CRC risk were evaluated using logistic regression models with random
199 province-specific intercepts. As fixed-effects terms, caloric and alcohol intake, self-reported
200 body mass index (BMI) and physical activity (metabolic equivalents (METs)) during the 10
201 years before diagnosis (cases) / interview (controls), age, smoking status, education, family
202 history of CRC and sex were considered as potential confounders. Scores of adherence were
203 analyzed both, as categorical (grouping the scores of adherence into quartiles of their
204 distribution among controls) and continuous variables (1-standard deviation increase taking
205 into account the dispersion among controls). Heterogeneity of the effects by sex was tested

206 including in the models an interaction term between the score of adherence and sex.
207 Multinomial logistic regression models were used to evaluate the association of the adherence
208 to the Western, Prudent and Mediterranean dietary patterns with proximal colon, distal colon
209 and rectal cancer separately. These models were adjusted by the same set of variables described
210 before but including province of residence as a fixed effect term.

211 Finally, assuming a causal relationship between the adherence to each of the patterns
212 and CRC risk for all analyses, the population attributable fraction (PAF%) was calculated using
213 Hanley's J.A. formula [33] to estimate the proportion of total cancer in the population that
214 hypothetically would not have occurred if all participants were in the optimal quartile of
215 adherence to the dietary patterns (first quartile for Western and fourth quartile for Prudent and
216 Mediterranean dietary patterns). Confidence intervals for PAF were computed using bootstrap
217 with 500 iterations.

218 Analyses were performed using STATA/MP (version 14.1, 2015, StataCorp LP) and
219 statistical significance was set at 2-sided $p < 0.05$.

220 **RESULTS**

221 Among the initially recruited participants, 3509 (89%) controls and 1889 (88%) cases
222 reported data on diet. Cases that provided dietary information later than 6 months after
223 diagnosis were excluded ($n=260$). Therefore, 1629 cases and 3509 controls aged 22 to 85 years
224 were included in the study (supplementary figure S1). The multivariable analyses were carried
225 out over 1530 cases and 3240 controls because data on either BMI ($<5\%$), physical activity
226 ($<1\%$), smoking status ($<1\%$), total energy ($<2\%$) or alcohol intake ($<2\%$) was missing for 99
227 cases and 269 controls.

228 Compared to controls, CRC cases were more adherent to the Western ($p < 0.001$) and
229 Mediterranean ($p = 0.015$) dietary patterns and reported higher energy ($p < 0.001$) and alcohol

230 (p=0.001) intake. CRC cases were also older (p<0.001) and reported higher BMI (p<0.001) and
231 lower levels of physical activity (p<0.001). The proportion of former smokers (p<0.001), males
232 (p<0.001), participants with no formal education (p<0.001) or with family history of CRC
233 (p<0.001) was also higher among cases (**Table 2**).

234 Results from **Table 3** revealed a positive association between a high adherence to the
235 Western dietary pattern and global CRC ($OR_{\text{fourth}(Q4)\text{vs.first}(Q1)\text{quartile}}$ (95% CI):1.50(1.20;1.87))
236 risk that was similar among males and females (p-interaction=0.615). Once the difference in
237 calorie intake was taken into account, a high adherence to the Mediterranean pattern appeared
238 to be protective ($OR_{Q4\text{vs}Q1}$ (95% CI):0.65(0.53;0.80)), with this effect slightly stronger among
239 females ($OR_{Q4\text{vs}Q1}$ (95% CI):0.56(0.40;0.77)) than among males ($OR_{Q4\text{vs}Q1}$
240 (95% CI):0.71(0.55;0.92)), though the p-value for the heterogeneity of the linear effects was not
241 significant (p-interaction=0.733). Assuming a causal relationship between diet and CRC risk,
242 the estimations indicate that 1/4 and 1/5 of CRC cases could have been prevented if all the
243 participants had been in the lowest category of adherence to the Western and in the highest
244 category of adherence to the Mediterranean dietary patterns respectively.

245 Stratified results by tumor subtype (**Table 4**) also indicate a detrimental effect of a high
246 adherence to the Western dietary pattern over CRC risk that seems to be confined to distal colon
247 ($OR_{Q4\text{vs}Q1}$ (95% CI):2.02(1.44;2.84)) and rectal tumors ($OR_{Q4\text{vs}Q1}$ (95% CI):1.46(1.05;2.01); p-
248 heterogeneity=0.087), while the protective effect of the Mediterranean dietary pattern was
249 similar for all tumor sites (Proximal colon: $OR_{Q4\text{vs}Q1}$ (95% CI):0.70(0.51;0.97); Distal Colon:
250 $OR_{Q4\text{vs}Q1}$ (95% CI):0.65(0.48;0.89); Rectum: $OR_{Q4\text{vs}Q1}$ (95% CI):0.60 (0.45;0.81); p-
251 heterogeneity=0.746). In agreement with these results, it was estimated that more than 1/3 of
252 distal colon and 1/4 of rectum tumors could have been prevented if all the study participants
253 were in the lowest quartile of adherence to the Western dietary pattern and 1/5 for distal colon

254 and 1/4 for rectum tumors could have been prevented with the highest adherences to
255 Mediterranean dietary pattern.

256 A high adherence to the prudent pattern did not show an association with CRC risk.

257 **DISCUSSION**

258 The detrimental effect of a high adherence to the Western dietary pattern for breast
259 [24,27] and gastric [28] cancer and the differential effect of a high adherence to the Prudent
260 (null) and to the Mediterranean (protective) dietary patterns over breast [24,27], prostate [29]
261 and gastric cancer [28] identified in the previous studies was also found for CRC in the present
262 work. Concretely, we found that a high adherence to the Western dietary pattern might increase
263 CRC risk in both males and females and that such risk might be confined to distal colon and
264 rectal cancer. Also, a high adherence to the Mediterranean dietary pattern showed a general
265 protective effect against CRC that was very similar among males and females and for all cancer
266 sites. On the contrary, the adherence to the Prudent dietary pattern was not associated to CRC.

267 Some recent reviews and metaanalysis [9,19,22], also report a positive association
268 between a high adherence to the Western dietary pattern and CRC risk and a protective effect
269 of a diet rich in fruits, vegetables, legumes and/or fish. The studies published after these
270 reviews, also report a positive association of a high adherence to the Western dietary pattern
271 with global CRC risk [18,20,21,23] and a possible protective effect of a Healthy diet against
272 this tumor [18,20,21]. In agreement with our results, some authors conclude that the effect of
273 the Western and Healthy diet might be stronger for distal colon and rectal cancer [21,22] or
274 indicate stronger effects of the Western diet in distal colon tumors [9]. Only a few of these
275 studies provide information of a possible interaction between diet and sex [18,20,21] and none
276 of them report significant differences.. Similarly, the current evidence for index based dietary
277 patterns point to a detrimental effect of pro-inflammatory diets (similar to our Western pattern)

278 for CRC risk [34] and a protective effect of diets that share common characteristics with our
279 Mediterranean pattern against this type of tumor [34,35]. One of the most important findings of
280 the present study is the difference in the associations found for Prudent and Mediterranean
281 dietary patterns. To understand these differences, we explored the associations of CRC risk
282 with individual food groups (supplementary Table S1). We believe that the protective effect of
283 the Mediterranean pattern against the null effect of the Prudent might be greatly explained by
284 the protective effect of oily fish, nuts and olives and olive oil, only present in the Mediterranean
285 pattern, but also by the detrimental effect of juices intake, only included in the Prudent pattern,
286 that might counteract the positive effect of a high consumption of fruits, vegetables and whole
287 grains characteristic of this pattern.

288 Some biological mechanisms support the associations found. On the one hand, the
289 “Western”-like diet high in fat, refined grains, red and processed meats and sweets has been
290 associated with higher levels of inflammatory markers [36] and with inflammation-related
291 chronic diseases [37]. Moreover, the high content of iron in meat products present in this pattern
292 generates free radicals that attack DNA and damage the tissue [38]. Additionally, processing
293 meat at high temperatures produces carcinogens such as N-nitroso and polycyclic aromatic
294 hydrocarbons [39]. On the other hand, the antioxidants from fruits, vegetables and legumes
295 present in the Mediterranean pattern may reduce risk by quenching free radicals and reducing
296 oxidative damage to DNA [40]. Furthermore, fiber dilutes faecal content, decreases transit time
297 and increases stool weight [41] contributing to a healthier gastrointestinal tract. Different
298 carcinogenic pathways in proximal and distal tumors have been suggested, based on their
299 molecular differences [42]. In this sense, the higher effect of the Western dietary pattern
300 (characterized by a low dietary fiber intake) in distal colon and rectal tumors, might reflect a
301 higher susceptibility to dietary carcinogens due to a less mature phenotype and lower immune
302 activity of dendritic cells involved in immunologic surveillance at this location [43]. Olive oil

303 intake has also been suggested to inhibit colon cancer development by inducing apoptosis and
304 down-regulating the expression of cyclooxygenase2 and Bcl-2 proteins that have a crucial role
305 in colorectal carcinogenesis [44]. Finally, the gut microbiome seems to play an important role
306 in colorectal carcinogenesis [45], and dietary habits strongly influence it [46]. Turnbaugh et al.
307 [46] recently demonstrated in an animal model that changing from low-fat, plant based diets to
308 high-fat, high-sugar diets can shift the structure of the microbiota, modify the representation of
309 metabolic pathways in the microbiome, and alter microbiome gene expression .

310 Our results should be interpreted in the context of the study's limitations. Recall bias is
311 always a concern in case-control studies. Anticipating the existence of this bias, some questions
312 about general dietary habits were included in the questionnaire and used to adjust the responses
313 to the FFQ [47]. Additionally, only cases that responded to the questionnaire within the 6
314 months following their diagnosis were included. On the other hand, the participation rates (64%
315 among CRC cases and 53% among controls) might give rise to some concerns about a possible
316 selection bias. In this sense, participating controls might have healthier lifestyles than the
317 general population, resulting in an overestimation of the effects. However, no effect was found
318 for the prudent pattern that includes consumption of products widely known as "Healthy".
319 Therefore, we believe that this bias, if it exists, would be non-differential. Finally, the biological
320 plausibility of the associations found, their strength, their consistency across sex and tumor site,
321 their consistency with the results from other studies on CRC [9,17-23] and the reproducibility
322 of the results across different studies and tumors [24,27-29], deem it unlikely that our findings
323 are a result of recall or selection bias.

324 One of the main strengths of the current research is the recruitment of histologically
325 confirmed incident cases of CRC and population controls. Furthermore, the geographical
326 variability of the recruited participants, coming from 11 provinces from the North, South,
327 Center, West and East of the country, ensured the representation of the different diets coexisting

328 within Spain. Also, the sample size allowed the evaluation of potential interactions of diet and
329 sex and the exploration of the associations by tumor localization. We also carried out a
330 sensitivity analysis to explore all the associations excluding 42 in situ cases and obtained very
331 similar results that led to the same exact conclusions (supplementary material tables S2 and
332 S3). Additionally, as mentioned before, we explored the associations of CRC risk with
333 individual food groups to ensure the associations found for patterns are not only due to the
334 presence in the pattern of one or two foods associated with this tumor (supplementary material
335 table S1). High consumers of high fat dairy products, meats, refined grains and sweets (products
336 characteristic of the Western Pattern) showed higher risk of CRC, while high consumers of oily
337 fish, vegetables, fruits, nuts and olive oil (foods present in the Mediterranean pattern) seemed
338 to be protected against this tumor. Therefore, most of the components of the two patterns
339 associated with CRC were also individually associated with this tumor, making it unlikely that
340 the associations found for the whole dietary patterns are due only to the association of CRC
341 with some individual foods. Finally, the reproducibility [26] and applicability [25] of the
342 Western, Prudent and Mediterranean dietary patterns applied here was previously tested,
343 demonstrating that the scores of adherence to these patterns can be calculated following the
344 exact same rules over different populations, resulting in different levels of adherence but still
345 being valid, which is supported by the similitude of the results found for breast [24,27], prostate
346 [29] and gastric cancer [28] and the present results found for CRC.

347 Our results provide evidence about very specific associations between diet and CRC
348 that could be useful to clinical practitioners and public health professionals to offer nutritional
349 recommendations based on avoiding the Western dietary pattern and promoting the
350 Mediterranean diet. Even though other risk factors are involved in the genesis of these type of
351 tumors, diet is a key risk factor for colorectal cancer. In this sense, if a country like Spain, with
352 a high compliance with the Mediterranean diet and a moderate adherence to the Western diet,

353 can benefit from abandoning the latter in favor of the former, the benefit might be greater in
354 countries with unhealthier diets.

355

356 **CONCLUSION**

357 A high consumption of fruits, vegetables and whole grains combined with a low dietary
358 fat intake might not be enough to prevent CRC. A fair percentage of colorectal cancer cases
359 could be reduced in the general population by providing dietary recommendations based in a
360 decrease of the consumption of high-fat dairy products, red and processed meat, refined grains,
361 sweets, caloric drinks, juices, convenience food and sauces in favor of an increase in the intake
362 of whole fruits, vegetables, legumes, olive oil, nuts and fish, especially for distal colon and
363 rectal tumors.

364

365

366

367 **ETHICAL STANDARDS:**

368 The MCC-Spain study protocol was approved by the Ethics Committee of each the
369 participating institutions and has been performed in accordance with the ethical standards as
370 laid down in the 1964 Declaration of Helsinki and its later amendments. All participants were
371 informed about the study objectives and signed an informed consent.

372 **CONFLICT OF INTEREST:**

373 The authors declare that they have no conflict of interest.

374 **REFERENCES:**

- 375 1. Arnold M, Karim-Kos HE, Coebergh JW, Byrnes G, Antilla A, Ferlay J, Renehan AG, Forman D,
376 Soerjomataram I (2015) Recent trends in incidence of five common cancers in 26 European countries since
377 1988: Analysis of the European Cancer Observatory. *European journal of cancer* 51 (9):1164-1187.
378 doi:10.1016/j.ejca.2013.09.002
- 379 2. Stewart B, Wild C, Editors (2014) (2014) *World Cancer Report 2014*. Lyon, France: International
380 Agency for Research on Cancer. .
- 381 3. WCRF (2015) Cancer preventability estimates for diet, nutrition, body fatness, and physical activity.
382 [http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-estimates-diet-](http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-estimates-diet-nutrition)
383 [nutrition](http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-estimates-diet-nutrition). Accessed 26/10/2016 2016
- 384 4. Whiteman DC, Webb PM, Green AC, Neale RE, Fritschi L, Bain CJ, Parkin DM, Wilson LF, Olsen
385 CM, Nagle CM, Pandeya N, Jordan SJ, Antonsson A, Kendall BJ, Hughes MC, Ibiebele TI, Miura K, Peters S,
386 Carey RN (2015) Cancers in Australia in 2010 attributable to modifiable factors: summary and conclusions.
387 *Australian and New Zealand journal of public health* 39 (5):477-484. doi:10.1111/1753-6405.12471
- 388 5. Song M, Garrett WS, Chan AT (2015) Nutrients, foods, and colorectal cancer prevention.
389 *Gastroenterology* 148 (6):1244-1260.e1216. doi:10.1053/j.gastro.2014.12.035
- 390 6. WCRF/AICR (2017) *World Cancer Research Fund / American Institute for Cancer Research.*
391 *Continuous Update Project Report. Diet, nutrition, physical activity and colorectal cancer. 2017*
392 http://www.wcrf.org/sites/default/files/CUP%20Colorectal%20Report_2017_Digital.pdf. Accessed 10th October
393 2017
- 394 7. Aune D, Chan DS, Vieira AR, Navarro Rosenblatt DA, Vieira R, Greenwood DC, Kampman E, Norat T
395 (2013) Red and processed meat intake and risk of colorectal adenomas: a systematic review and meta-analysis of
396 epidemiological studies. *Cancer causes & control : CCC* 24 (4):611-627. doi:10.1007/s10552-012-0139-z
- 397 8. Cai S, Li Y, Ding Y, Chen K, Jin M (2014) Alcohol drinking and the risk of colorectal cancer death: a
398 meta-analysis. *European journal of cancer prevention : the official journal of the European Cancer Prevention*
399 *Organisation (ECP)* 23 (6):532-539. doi:10.1097/cej.0000000000000076
- 400 9. Magalhaes B, Peleteiro B, Lunet N (2012) Dietary patterns and colorectal cancer: systematic review and
401 meta-analysis. *European journal of cancer prevention : the official journal of the European Cancer Prevention*
402 *Organisation (ECP)* 21 (1):15-23. doi:10.1097/CEJ.0b013e3283472241
- 403 10. Barkoukis H (2007) Importance of understanding food consumption patterns. *J Am Diet Assoc* 107
404 (2):234-236
- 405 11. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*
406 13 (1):3-9
- 407 12. Jacques PF, Tucker KL (2001) Are dietary patterns useful for understanding the role of diet in chronic
408 disease? *The American journal of clinical nutrition* 73 (1):1-2

- 409 13. Fung TT, McCullough ML, Newby PK, Manson JE, Meigs JB, Rifai N, Willett WC, Hu FB (2005)
410 Diet-quality scores and plasma concentrations of markers of inflammation and endothelial dysfunction. *The*
411 *American journal of clinical nutrition* 82 (1):163-173
- 412 14. Liese AD, Krebs-Smith SM, Subar AF, George SM, Harmon BE, Neuhauser ML, Boushey CJ, Schap
413 TE, Reedy J (2015) The Dietary Patterns Methods Project: synthesis of findings across cohorts and relevance to
414 dietary guidance. *The Journal of nutrition* 145 (3):393-402. doi:10.3945/jn.114.205336
- 415 15. Schwingshackl L, Bogensberger B, Hoffmann G (2018) Diet Quality as Assessed by the Healthy Eating
416 Index, Alternate Healthy Eating Index, Dietary Approaches to Stop Hypertension Score, and Health Outcomes:
417 An Updated Systematic Review and Meta-Analysis of Cohort Studies. *Journal of the Academy of Nutrition and*
418 *Dietetics* 118 (1):74-100.e111. doi:10.1016/j.jand.2017.08.024
- 419 16. Giacosa A, Barale R, Bavaresco L, Gatenby P, Gerbi V, Janssens J, Johnston B, Kas K, La Vecchia C,
420 Mainguet P, Morazzoni P, Negri E, Pelucchi C, Pezzotti M, Rondanelli M (2013) Cancer prevention in Europe:
421 the Mediterranean diet as a protective choice. *European journal of cancer prevention : the official journal of the*
422 *European Cancer Prevention Organisation (ECP)* 22 (1):90-95. doi:10.1097/CEJ.0b013e328354d2d7
- 423 17. Safari A, Shariff ZM, Kandiah M, Rashidkhani B, Fereidooni F (2013) Dietary patterns and risk of
424 colorectal cancer in Tehran Province: a case-control study. *BMC public health* 13:222. doi:10.1186/1471-2458-
425 13-222
- 426 18. Park Y, Lee J, Oh JH, Shin A, Kim J (2016) Dietary patterns and colorectal cancer risk in a Korean
427 population: A case-control study. *Medicine* 95 (25):e3759. doi:10.1097/md.0000000000003759
- 428 19. Feng YL, Shu L, Zheng PF, Zhang XY, Si CJ, Yu XL, Gao W, Zhang L (2017) Dietary patterns and
429 colorectal cancer risk: a meta-analysis. *European journal of cancer prevention : the official journal of the*
430 *European Cancer Prevention Organisation (ECP)* 26 (3):201-211. doi:10.1097/cej.0000000000000245
- 431 20. Haslam A, Wagner Robb S, Hebert JR, Huang H, Ebell MH (2017) Association between dietary pattern
432 scores and the prevalence of colorectal adenoma considering population subgroups. *Nutrition & dietetics: the*
433 *journal of the Dietitians Association of Australia*. doi:10.1111/1747-0080.12400
- 434 21. Mehta RS, Song M, Nishihara R, Drew DA, Wu K, Qian ZR, Fung TT, Hamada T, Masugi Y, da Silva
435 A, Shi Y, Li W, Gu M, Willett WC, Fuchs CS, Giovannucci EL, Ogino S, Chan AT (2017) Dietary Patterns and
436 Risk of Colorectal Cancer: Analysis by Tumor Location and Molecular Subtypes. *Gastroenterology* 152
437 (8):1944-1953.e1941. doi:10.1053/j.gastro.2017.02.015
- 438 22. Randi G, Edefonti V, Ferraroni M, La Vecchia C, Decarli A (2010) Dietary patterns and the risk of
439 colorectal cancer and adenomas. *Nutrition reviews* 68 (7):389-408. doi:10.1111/j.1753-4887.2010.00299.x
- 440 23. Tayyem RF, Bawadi HA, Shehadah I, Agraib LM, AbuMweis SS, Al-Jaberi T, Al-Nusairr M, Bani-
441 Hani KE, Heath DD (2017) Dietary patterns and colorectal cancer. *Clinical nutrition (Edinburgh, Scotland)* 36
442 (3):848-852. doi:10.1016/j.clnu.2016.04.029
- 443 24. Castello A, Pollan M, Buijsse B, Ruiz A, Casas AM, Baena-Canada JM, Lope V, Antolin S, Ramos M,
444 Munoz M, Lluch A, de Juan-Ferre A, Jara C, Jimeno MA, Rosado P, Diaz E, Guillem V, Carrasco E, Perez-
445 Gomez B, Vioque J, Boeing H, Martin M (2014) Spanish Mediterranean diet and other dietary patterns and
446 breast cancer risk: case-control EpiGEICAM study. *British journal of cancer* 111 (7):9.
447 doi:10.1038/bjc.2014.434
- 448 25. Castello A, Buijsse B, Martin M, Ruiz A, Casas AM, Baena-Canada JM, Pastor-Barriuso R, Antolin S,
449 Ramos M, Munoz M, Lluch A, de Juan-Ferre A, Jara C, Lope V, Jimeno MA, Arriola-Arellano E, Diaz E,
450 Guillem V, Carrasco E, Perez-Gomez B, Vioque J, Pollan M, researchers G (2016) Evaluating the Applicability
451 of Data-Driven Dietary Patterns to Independent Samples with a Focus on Measurement Tools for Pattern
452 Similarity. *Journal of the Academy of Nutrition and Dietetics* 116 (12):1914-1924 e1916.
453 doi:10.1016/j.jand.2016.05.008
- 454 26. Castello A, Lope V, Vioque J, Santamarina C, Pedraz-Pingarron C, Abad S, Ederra M, Salas-Trejo D,
455 Vidal C, Sanchez-Contador C, Aragones N, Perez-Gomez B, Pollan M (2016) Reproducibility of data-driven
456 dietary patterns in two groups of adult Spanish women from different studies. *The British journal of nutrition*
457 116 (4):734-742. doi:10.1017/S000711451600252X
- 458 27. Castello A, Boldo E, Perez-Gomez B, Lope V, Altzibar JM, Martin V, Castano-Vinyals G, Guevara M,
459 Dierssen-Sotos T, Tardon A, Moreno V, Puig-Vives M, Llorens-Ivorra C, Alguacil J, Gomez-Acebo I, Castilla J,
460 Gracia-Lavedan E, Davila-Batista V, Kogevinas M, Aragones N, Amiano P, Pollan M (2017) Adherence to the
461 Western, Prudent and Mediterranean dietary patterns and breast cancer risk: MCC-Spain study. *Maturitas* 103:8-
462 15. doi:10.1016/j.maturitas.2017.06.020
- 463 28. Castelló A, Fernández de Larrea N, Martin M, Dávila-Batista V, Boldo E, Guevara M, Moreno V,
464 Castaño-Vinyals G, Gómez-Acebo I, Fernandez-Tardon G, Peiro R, Olmedo-Requena R, Capelo R, Navarro C,
465 Pacho-Valbuena S, Pérez-Gómez B, Kogevinas M, Pollán M, Aragónés N (2017) High Adherence to the
466 Western, Prudent and Mediterranean Dietary Patterns and Risk of Gastric Adenocarcinoma. (In Press). *Gastric*
467 *Cancer*

- 468 29. Castello A, Boldo E, Amiano P, Castano-Vinyals G, Aragones N, Gomez-Acebo I, Peiro R, Jimenez-
469 Moleon JJ, Alguacil J, Tardon A, Cecchini L, Lope V, Dierssen-Sotos T, Mengual L, Kogevinas M, Pollan M,
470 Perez-Gomez B, researchers M-S (2017) Mediterranean dietary pattern is associated to low risk of aggressive
471 prostate cancer: MCC-Spain study. *The Journal of urology*. doi:10.1016/j.juro.2017.08.087
- 472 30. Castano-Vinyals G, Aragones N, Perez-Gomez B, Martin V, Llorca J, Moreno V, Altzibar JM, Ardanaz
473 E, de Sanjose S, Jimenez-Moleon JJ, Tardon A, Alguacil J, Peiro R, Marcos-Gragera R, Navarro C, Pollan M,
474 Kogevinas M, Group MC-SS (2015) Population-based multicase-control study in common tumors in Spain
475 (MCC-Spain): rationale and study design. *Gaceta sanitaria / SESPAS* 29 (4):308-315.
476 doi:10.1016/j.gaceta.2014.12.003
- 477 31. Garcia-Closas R, Garcia-Closas M, Kogevinas M, Malats N, Silverman D, Serra C, Tardon A, Carrato
478 A, Castano-Vinyals G, Dosemeci M, Moore L, Rothman N, Sinha R (2007) Food, nutrient and heterocyclic
479 amine intake and the risk of bladder cancer. *European journal of cancer* 43 (11):1731-1740.
480 doi:10.1016/j.ejca.2007.05.007
- 481 32. Burt C (1948) Factor Analysis and canonical correlations. *Br J Math Stat Psychol* 1 (2):95-106
- 482 33. Hanley JA (2001) A heuristic approach to the formulas for population attributable fraction. *Journal of*
483 *epidemiology and community health* 55 (7):508-514
- 484 34. Steck SE, Guinter M, Zheng J, Thomson CA (2015) Index-based dietary patterns and colorectal cancer
485 risk: a systematic review. *Advances in nutrition (Bethesda, Md)* 6 (6):763-773. doi:10.3945/an.115.009746
- 486 35. Schwingshackl L, Schwedhelm C, Galbete C, Hoffmann G (2017) Adherence to Mediterranean Diet
487 and Risk of Cancer: An Updated Systematic Review and Meta-Analysis. *Nutrients* 9 (10).
488 doi:10.3390/nu9101063
- 489 36. Barbaresko J, Koch M, Schulze MB, Nothlings U (2013) Dietary pattern analysis and biomarkers of
490 low-grade inflammation: a systematic literature review. *Nutrition reviews* 71 (8):511-527.
491 doi:10.1111/nure.12035
- 492 37. Thorburn AN, Macia L, Mackay CR (2014) Diet, metabolites, and "western-lifestyle" inflammatory
493 diseases. *Immunity* 40 (6):833-842. doi:10.1016/j.immuni.2014.05.014
- 494 38. Ashmore JH, Rogers CJ, Kelleher SL, Lesko SM, Hartman TJ (2016) Dietary Iron and Colorectal
495 Cancer Risk: A Review of Human Population Studies. *Critical reviews in food science and nutrition* 56
496 (6):1012-1020. doi:10.1080/10408398.2012.749208
- 497 39. Santarelli RL, Pierre F, Corpet DE (2008) Processed meat and colorectal cancer: a review of
498 epidemiologic and experimental evidence. *Nutrition and cancer* 60 (2):131-144.
499 doi:10.1080/01635580701684872
- 500 40. Fang YZ, Yang S, Wu G (2002) Free radicals, antioxidants, and nutrition. *Nutrition (Burbank, Los*
501 *Angeles County, Calif)* 18 (10):872-879
- 502 41. Giovannucci E, Willett WC (1994) Dietary factors and risk of colon cancer. *Annals of medicine* 26
503 (6):443-452
- 504 42. Missiaglia E, Jacobs B, D'Ario G, Di Narzo AF, Sonesson C, Budinska E, Popovici V, Vecchione L,
505 Gerster S, Yan P, Roth AD, Klingbiel D, Bosman FT, Delorenzi M, Tejpar S (2014) Distal and proximal colon
506 cancers differ in terms of molecular, pathological, and clinical features. *Annals of oncology : official journal of*
507 *the European Society for Medical Oncology* 25 (10):1995-2001. doi:10.1093/annonc/mdu275
- 508 43. Bernardo D, Durant L, Mann ER, Bassity E, Montalvillo E, Man R, Vora R, Reddi D, Bayiroglu F,
509 Fernandez-Salazar L, English NR, Peake ST, Landy J, Lee GH, Malietzis G, Siaw YH, Murugananthan AU,
510 Hendy P, Sanchez-Recio E, Phillips RK, Garrote JA, Scott P, Parkhill J, Paulsen M, Hart AL, Al-Hassi HO,
511 Arranz E, Walker AW, Carding SR, Knight SC (2016) Chemokine (C-C Motif) Receptor 2 Mediates Dendritic
512 Cell Recruitment to the Human Colon but Is Not Responsible for Differences Observed in Dendritic Cell
513 Subsets, Phenotype, and Function Between the Proximal and Distal Colon. *Cellular and molecular*
514 *gastroenterology and hepatology* 2 (1):22-39 e25. doi:10.1016/j.jcmgh.2015.08.006
- 515 44. Pelucchi C, Bosetti C, Negri E, Lipworth L, La Vecchia C (2011) Olive oil and cancer risk: an update of
516 epidemiological findings through 2010. *Current pharmaceutical design* 17 (8):805-812
- 517 45. Schwabe RF, Jobin C (2013) The microbiome and cancer. *Nature reviews Cancer* 13 (11):800-812.
518 doi:10.1038/nrc3610
- 519 46. Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, Gordon JI (2009) The effect of diet on the
520 human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *Science translational medicine*
521 1 (6):6ra14. doi:10.1126/scitranslmed.3000322
- 522 47. Calvert C, Cade J, Barrett JH, Woodhouse A (1997) Using cross-check questions to address the
523 problem of mis-reporting of specific food groups on Food Frequency Questionnaires. UKWCS Steering Group.
524 United Kingdom Women's Cohort Study Steering Group. *European journal of clinical nutrition* 51 (10):708-712
525

527 **Table 1:** Composition of food groups based on the food frequency questionnaire of the MCC-
 528 Spain study and component loadings for each pattern identified in the EpiGEICAM study²⁵.

FOOD GROUP	FOOD ^a	West ^b	Prud ^b	Med ^b
HIGH FAT DAIRY	Whole-fat milk, double cream, condensed milk, whole-fat yogurt, semi-cured, cured, or creamy cheese, blue cheese ^c , custard, milk shake ^c , ice-cream,	0.60	-0.11	0.20
LOW FAT DAIRY	Semi-skimmed and skimmed milk, soy milk ^c , skimmed yogurt, curd, cottage or fresh white cheese.	-0.49	0.60	-0.01
EGGS	Eggs.	0.19	0.08	0.16
WHITE MEAT	Chicken, rabbit and duck.	0.08	0.17	0.18
RED MEAT	Pork, beef, lamb, liver (beef, pork or chicken), entrails, hamburgers (pork or beef) and meatballs (pork or beef) ^c .	0.27	0.09	0.22
PROCESSED MEAT	Sausages, serrano ham ^c and other cold meat, bacon, pâté, foie-gras.	0.36	0.10	0.26
WHITE FISH	Fresh or frozen white fish (hake, sea bass, sea bream), $\frac{1}{2}$ salted fish ^c and $\frac{1}{2}$ smoked fish ^c .	0.01	0.24	0.34
OILY FISH	Fresh or frozen blue fish (tuna, swordfish, sardines, anchovies, salmon), canned fish, $\frac{1}{2}$ salted fish ^c and $\frac{1}{2}$ smoked fish ^c .	0.05	0.24	0.44
SEAFOOD/SHELL FISH	Clams, mussels, oysters, squid, cuttlefish, octopus, prawn, crab, shrimp and similar products.	0.17	0.27	0.35
LEAFY VEGETABLES	Spinach, chard, lettuce and other leafy vegetables.	-0.11	0.34	0.40
FRUITING VEGETABLES	Tomato, eggplant, zucchini, cucumber, pepper, artichoke and avocado ^c .	0.00	0.36	0.45
ROOT VEGETABLES	Carrot, pumpkin and radish ^c .	0.05	0.35	0.44
OTHER VEGETABLES	Cooked cabbage, cauliflower or broccoli, onion, green beans, asparagus, mushrooms ^c , corn, garlic, gazpacho ^c , vegetable soup ^c and other vegetables ^c .	-0.04	0.40	0.42
LEGUMES ^d	Peas ^c , lentils ^c , chickpeas ^c , beans ^c and broad beans ^c .	0.21	0.15	0.34
POTATOES	Roasted or boiled potatoes and sweet potatoes ^c .	0.17	0.25	0.40
FRUITS	Orange, grapefruit ^c , mandarin, banana, apple, pear, grapes, kiwi, strawberries ^c , cherries ^c , peach, figs ^c , melon or watermelon, prunes, mango ^c and papaya ^c and other fresh or dried fruits ^c .	-0.07	0.31	0.31
NUTS	Almonds, peanuts, pine nuts, hazelnut	0.18	0.22	0.29

FOOD GROUP	FOOD^a	West^b	Prud^b	Med^b
REFINED GRAINS	White-flour bread, rice, pasta	0.37	0.15	0.23
WHOLE GRAINS	Whole-grain bread and breakfast cereals	-0.43	0.47	-0.06
OLIVES AND VEGETABLE OIL	Olives, added olive oil to salads, bread and dishes, other vegetable oils (sunflower, corn, and soybean).	0.12	0.19	0.34
OTHER EDIBLE FATS	Margarine, butter and lard^c .	0.22	0.02	0.11
SWEETS	Chocolate and other sweets, cocoa powder, plain cookies, chocolate cookies, pastries (croissant, donut, cake, pie or similar)	0.35	0.18	0.05
SUGARY	Jam, honey, sugar and fruit in sugar syrup^c .	0.24	0.05	0.00
JUICES	Tomato juice^c , freshly squeezed orange juice, juice (other than freshly squeezed)	0.25	0.67	-0.39
CALORIC DRINKS	Sugar-sweetened soft drinks and nut milk^c .	0.74	0.21	-0.25
CONVENIENCE FOOD AND SAUCES	Croquette, fish sticks, dumplings^c , kebab^c , fried potatoes, crisps, pizza, instant soup^c , mayonnaise, tomato sauce, hot sauce^c , ketchup and other sauces^c .	0.47	0.12	0.24

529 ^a Log-transformed centered intake in grams.

530 ^b West: Western; Prud: Prudent; Med: Mediterranean

531 ^c in bold items that are included in the FFQ from MCC-Spain study that were not included in
532 EpiGEICAM.

533 ^d FFQ questionnaire from EpiGEICAM only included a single general question on legumes
534 intake whereas MCC-Spain included more detailed information on the type of legumes.

535

536 **Table 2.** Description of scores of adherence to Western, Prudent and Mediterranean dietary
 537 patterns and other baseline characteristics for colorectal cancer cases and controls.

	Controls n=3509	Cases n=1629	p
Western mean (sd)^a	-0.38 (3.52)	0.14 (3.52)	<0.001
Prudent mean (sd)^a	-0.09 (3.29)	-0.19 (3.32)	0.353
Mediterranean mean (sd)^a	-0.02 (2.89)	0.19 (2.75)	0.015
Energy (kcal/day) mean (sd)	1903.81 (570.75)	2008.24 (638.31)	<0.001
Alcohol (g/day) median (IQR)	7.22 (0.00;23.21)	9.38 (0.00;34.72)	0.001
BMI^b (kg/m²) mean (sd)	26.61 (4.41)	27.59 (4.46)	<0.001
Physical activity (METs^b/week) n (%^c)			<0.001
0	1341 (38%)	855 (52%)	
0.1-8	506 (14%)	183 (11%)	
8-15.9	422 (12%)	135 (8%)	
>=16	1202 (34%)	456 (28%)	
Unknown	38 (1%)	0 (0%)	
Age (years) mean (sd)	63.20 (11.69)	67.09 (10.63)	<0.001
Smoking n (%^c)			<0.001
Never Smoker	1549 (44%)	680 (42%)	
Former Smoker	1224 (35%)	660 (41%)	
Current Smoker	724 (21%)	279 (17%)	
Unknown	12 (0%)	10 (1%)	
Education n (%^c)			<0.001
No formal Education	619 (18%)	522 (32%)	
Primary School	1143 (33%)	648 (40%)	
Secondary School	1010 (29%)	311 (19%)	
University or more	737 (21%)	148 (9%)	
Previous history of CRC^b n (%^c)			<0.001
No	3101 (88%)	1295 (79%)	
2nd Degree	107 (3%)	62 (4%)	
One of 1st degree	281 (8%)	231 (14%)	
More than one of 1st degree	20 (1%)	41 (3%)	
Sex			<0.001
Male	1813 (52%)	1043 (64%)	
Female	1696 (48%)	586 (36%)	

538 ^a The pairwise Pearson correlations for the level of adherence to the three identified dietary
 539 patterns were 0.329 for the Western and Prudent, 0.231 for the Western and Mediterranean and
 540 0.485 for the Prudent and Mediterranean.

541 ^b BMI: Body mass index; CRC: Colorectal cancer; METS: Metabolic equivalent.

542 ^c Percentages might not add up 100 because of rounding.

543
544

Table 3. Association between colorectal cancer incidence and the scores of adherence to Western, Prudent and Mediterranean dietary patterns and attributable fractions for all participants and by sex.

		ALL n=4770				MALE n=2688		FEMALE n=2082		
		Co/Ca ^a	OR ^b (95%CI)	Co/Ca ^a	aOR ^c (95%CI)	Co/Ca ^a	aOR ^c (95%CI)	Co/Ca ^a	aOR ^c (95%CI)	p-int
WESTERN										
Quartiles	Q1	877/322	1	772/292	1	335/160	1	437/132	1	
	Q2	878/409	1.27 (1.06;1.51)	824/390	1.29 (1.06;1.57)	405/227	1.16 (0.89;1.52)	419/163	1.46 (1.10;1.94)	
	Q3	877/423	1.36 (1.14;1.62)	831/401	1.43 (1.17;1.75)	449/260	1.33 (1.02;1.73)	382/141	1.56 (1.16;2.10)	
	Q4	877/475	1.47 (1.23;1.75)	813/447	1.50 (1.20;1.87)	511/341	1.45 (1.11;1.91)	302/106	1.50 (1.07;2.09)	
p-trend			<0.001		<0.001		0.004		0.009	
1SD-increase			1.16 (1.09;1.24)		1.19 (1.10;1.30)		1.21 (1.09;1.34)		1.17 (1.04;1.31)	0.615
PAF^d%					24% (12%;36%)		21% (5%;37%)		18% (3%;33%)	
PRUDENT										
Quartiles	Q1	878/440	1	783/398	1	485/292	1	298/106	1	
	Q2	876/384	0.83 (0.70;0.98)	811/362	0.87 (0.72;1.05)	430/235	0.84 (0.66;1.05)	381/127	0.95 (0.69;1.31)	
	Q3	877/403	0.89 (0.75;1.05)	827/389	1.00 (0.83;1.21)	412/241	0.94 (0.74;1.19)	415/148	1.13 (0.83;1.54)	
	Q4	878/402	0.88 (0.74;1.04)	819/381	0.94 (0.76;1.15)	373/220	0.88 (0.68;1.13)	446/161	1.05 (0.77;1.44)	
p-trend			0.242		0.875		0.475		0.515	
1SD^a-increase			0.96 (0.90;1.02)		0.97 (0.90;1.05)		0.95 (0.86;1.04)		1.02 (0.90;1.15)	0.330
PAF^d%					2% (-12%;15%)		4% (-12%;21%)		3% (-12%;19%)	
MEDITERRANEAN										
Quartiles	Q1	878/394	1	796/359	1	398/206	1	398/153	1	
	Q2	877/412	0.98 (0.83;1.17)	821/386	0.91 (0.75;1.10)	390/236	0.99 (0.77;1.27)	431/150	0.83 (0.63;1.10)	
	Q3	876/371	0.80 (0.67;0.96)	815/357	0.72 (0.59;0.87)	425/219	0.71 (0.55;0.92)	390/138	0.74 (0.55;0.99)	
	Q4	878/452	0.90 (0.76;1.07)	808/428	0.65 (0.53;0.80)	487/327	0.71 (0.55;0.92)	321/101	0.56 (0.40;0.77)	
p-trend			0.073		0.000		0.001		0.000	
1SD^a-increase			0.98 (0.92;1.05)		0.87 (0.80;0.94)		0.88 (0.79;0.96)		0.85 (0.76;0.96)	0.733
PAF^d%					20% (8%;33%)		15% (2%;29%)		18% (3%;33%)	

545 ^a Co: Controls; Ca: Cases; SD: Standard deviation.

546 ^b Crude odds ratio of colorectal cancer associated with the adherence to the Western, Prudent and Mediterranean dietary patterns

547 ^c Odds ratio of colorectal cancer associated with the adherence to the Western, Prudent and Mediterranean dietary patterns adjusted by sex, age,
548 education, BMI, family history of colorectal cancer, physical activity, smoking status, caloric intake and alcohol intake as fixed effects and
549 province of residence as a random effect.

550 ^c Same as ^b including an interaction term with sex.

551 ^d PAF= Population attributable fraction. Proportion of colorectal cancer cases that could be prevented if all participants were in the most
552 beneficial category of adherence to each pattern (Q1 for Western and Q4 for Prudent and Mediterranean)

$$PAF = \frac{PF_{Q1} \cdot (OR_{Q1} - 1) + PF_{Q2} \cdot (OR_{Q2} - 1) + PF_{Q3} \cdot (OR_{Q3} - 1) + PF_{Q4} \cdot (OR_{Q4} - 1)}{1 + [PF_{Q1} \cdot (OR_{Q1} - 1) + PF_{Q2} \cdot (OR_{Q2} - 1) + PF_{Q3} \cdot (OR_{Q3} - 1) + PF_{Q4} \cdot (OR_{Q4} - 1)]} \cdot 100$$

553 PF=Proportion of population in the specific exposure category

OR= Odds ratio for the especific exposure category

SUPPLEMENTARY MATERIAL

Figure S1. Flow of colorectal cancer cases and controls through the MCC-Spain study stages. CRC, Colorectal cancer.




Table S1: Odds Ratio of colorectal cancer associated to quartiles of consumption of 26 food groups not adjusting and adjusting by the consumption of the rest of the foods.

	2nd Quartile	3rd Quartile	4th Quartile		2nd Quartile	3rd Quartile	4th Quartile	
	OR ^a (95%CI)	OR ^a (95%CI)	OR ^a (95%CI)	p for trend ^a	OR ^b (95%CI)	OR ^b (95%CI)	OR ^b (95%CI)	p for trend ^b
High fat dairy ^c	1.13(0.93;1.37)	1.21(1.00;1.48)	1.39(1.15;1.70)	0.001	1.08(0.88;1.32)	1.11(0.89;1.37)	1.19(0.94;1.50)	0.169
Low fatdairy ^c	0.81(0.68;0.98)	0.86(0.72;1.04)	0.70(0.58;0.85)	0.001	0.88(0.72;1.08)	0.99(0.8;1.23)	0.86(0.68;1.08)	0.394
Eggs ^d	.	1.13(0.95;1.34)	1.46(1.18;1.82)	0.003	.	1.00(0.84;1.20)	1.22(0.96;1.53)	0.261
White meat ^c	1.02(0.83;1.25)	1.12(0.93;1.36)	1.30(1.08;1.57)	0.003	0.98(0.79;1.21)	1.06(0.87;1.30)	1.30(1.06;1.59)	0.005
Red meat ^c	1.15(0.94;1.40)	1.53(1.26;1.87)	1.65(1.35;2.03)	0.000	1.11(0.90;1.37)	1.42(1.15;1.75)	1.39(1.12;1.74)	0.001
Processed meat ^c	1.14(0.94;1.40)	1.10(0.90;1.35)	1.58(1.28;1.94)	0.000	1.12(0.91;1.39)	1.02(0.82;1.27)	1.49(1.19;1.86)	0.001
White fish ^c	1.07(0.88;1.29)	1.10(0.91;1.32)	1.14(0.94;1.38)	0.173	1.07(0.88;1.32)	1.15(0.94;1.41)	1.32(1.07;1.63)	0.009
Oily fish ^c	0.96(0.80;1.15)	0.81(0.67;0.97)	0.78(0.65;0.95)	0.003	1.01(0.83;1.22)	0.84(0.69;1.03)	0.79(0.64;0.98)	0.014
Seafood ^c	1.01(0.84;1.22)	0.90(0.74;1.09)	0.86(0.71;1.05)	0.076	0.98(0.81;1.19)	0.87(0.71;1.06)	0.83(0.67;1.03)	0.049
Leafy vegetables ^c	0.90(0.75;1.08)	0.73(0.60;0.88)	0.56(0.46;0.68)	0.000	1.01(0.82;1.23)	0.90(0.73;1.12)	0.75(0.59;0.96)	0.017
Fruiting Vegetables ^c	0.77(0.64;0.92)	0.62(0.51;0.75)	0.59(0.48;0.71)	0.000	0.80(0.66;0.98)	0.73(0.59;0.90)	0.86(0.68;1.09)	0.155
Root Vegetables ^c	0.86(0.72;1.03)	0.75(0.62;0.90)	0.61(0.50;0.74)	0.000	0.90(0.74;1.08)	0.83(0.68;1.01)	0.76(0.61;0.95)	0.014
Other Vegetables ^c	0.92(0.77;1.10)	0.70(0.58;0.84)	0.64(0.53;0.78)	0.000	1.00(0.82;1.23)	0.82(0.66;1.02)	0.87(0.68;1.10)	0.081
Legumes ^d	.	0.87(0.73;1.03)	0.94(0.79;1.11)	0.375	.	0.91(0.76;1.08)	0.99(0.82;1.19)	0.833
Potatoes ^c	1.27(1.03;1.56)	1.61(1.32;1.97)	1.64(1.34;2.01)	0.000	1.29(1.04;1.60)	1.61(1.30;1.98)	1.65(1.33;2.04)	0.000
Fruits ^c	0.93(0.77;1.13)	0.85(0.70;1.03)	0.64(0.52;0.78)	0.000	1.03(0.85;1.26)	0.99(0.8;1.22)	0.80(0.64;1.00)	0.042
Nuts ^d	.	0.94(0.81;1.09)	0.73(0.60;0.89)	0.008	.	1.00(0.85;1.18)	0.90(0.73;1.10)	0.528
Olives and Vegetable Oil ^c	0.87(0.69;1.08)	0.86(0.73;1.02)	0.79(0.65;0.96)	0.019	0.90(0.72;1.14)	0.92(0.77;1.10)	0.90(0.73;1.12)	0.259
Other Edible Fats ^c	.	0.96(0.79;1.18)	1.16(0.98;1.36)	0.157	.	0.90(0.72;1.11)	1.02(0.86;1.22)	0.981
Refined Grains ^c	1.34(1.09;1.63)	1.37(1.12;1.68)	1.42(1.13;1.78)	0.004	1.20(0.96;1.49)	1.17(0.93;1.47)	1.24(0.96;1.6)	0.245
Whole grains ^d	.	0.79(0.66;0.94)	0.69(0.58;0.81)	0.000	.	0.82(0.68;0.99)	0.85(0.70;1.03)	0.018
Sweets ^c	1.16(0.95;1.40)	1.32(1.09;1.59)	1.29(1.05;1.58)	0.007	1.14(0.93;1.40)	1.27(1.03;1.55)	1.23(0.98;1.54)	0.092
Sugary ^c	1.25(1.03;1.53)	1.22(1.02;1.47)	1.44(1.19;1.75)	0.000	1.17(0.95;1.44)	1.10(0.90;1.34)	1.30(1.06;1.60)	0.027
Juices ^d	.	1.26(1.07;1.48)	1.39(1.18;1.64)	0.000	.	1.33(1.12;1.58)	1.58(1.32;1.88)	0.000
Caloric Drinks ^d	.	0.83(0.70;0.98)	0.98(0.83;1.15)	0.520	.	0.78(0.65;0.93)	0.84(0.70;1.00)	0.031
Convenience Food ^c	0.98(0.81;1.18)	1.11(0.92;1.34)	1.10(0.90;1.34)	0.212	0.87(0.72;1.07)	0.93(0.76;1.15)	0.84(0.67;1.05)	0.170

^a Adjusted by sex, age, education, BMI, family history of colorectal cancer, physical activity, smoking status, caloric intake and alcohol intake as fixed effects and province of residence as a random effect.

^b Adjusted by sex, age, education, BMI, family history of colorectal cancer, physical activity, smoking status, caloric intake, alcohol intake and food group intake as fixed effects and province of residence as a random effect.

^c Reference intake is first quartile.

^d Reference intake is first + second quartile due to the more uniform distribution of data.

1 **Table S2:** Association between colorectal cancer incidence and the scores of adherence to Western,
 2 Prudent and Mediterranean dietary patterns excluding in situ cases.

	ALL n=4729		MALE n=2662		FEMALE n=2067		
	Co/Ca	OR(95%CI)	Co/Ca	OR(95%CI)	Co/Ca	OR(95%CI)	p-het
WESTERN							
Q1	772/285	1	335/157	1	437/128	1	
Q2	824/375	1.26 (1.03;1.54)	405/221	1.14 (0.87;1.49)	419/154	1.42 (1.06;1.89)	
Q3	831/394	1.43 (1.17;1.76)	449/255	1.31 (1.00;1.71)	382/139	1.59 (1.18;2.15)	
Q4	813/435	1.48 (1.18;1.85)	511/329	1.40 (1.06;1.84)	302/106	1.54 (1.10;2.15)	
p-trend		<0.001		0.010		0.005	
1SD-increase		1.19 (1.09;1.29)		1.19 (1.08;1.32)		1.18 (1.05;1.33)	0.865
PRUDENT							
Q1	783/390	1	485/286	1	298/104	1	
Q2	812/351	0.85 (0.70;1.02)	431/229	0.82 (0.64;1.03)	381/122	0.92 (0.67;1.27)	
Q3	826/378	0.98 (0.81;1.19)	411/234	0.92 (0.72;1.17)	415/144	1.10 (0.81;1.51)	
Q4	819/370	0.90 (0.73;1.12)	373/213	0.84 (0.65;1.09)	446/157	1.02 (0.74;1.41)	
p-trend		0.659		0.322		0.606	
1SD-increase		0.96 (0.89;1.05)		0.94 (0.86;1.04)		1.01 (0.89;1.15)	0.347
MEDITERRANEAN							
Q1	796/346	1	398/198	1	398/148	1	
Q2	821/375	0.91 (0.75;1.11)	390/227	0.98 (0.76;1.27)	431/148	0.84 (0.63;1.12)	
Q3	816/348	0.71 (0.58;0.87)	425/215	0.71 (0.55;0.93)	391/133	0.72 (0.54;0.97)	
Q4	807/420	0.65 (0.52;0.80)	487/322	0.71 (0.55;0.92)	320/98	0.55 (0.40;0.76)	
p-trend		<0.001		0.002		<0.001	
1SD-increase		0.87 (0.80;0.94)		0.87 (0.79;0.96)		0.85 (0.75;0.96)	0.725

3
 4 **Table S3.** Adjusted odds ratios for the association between proximal colon, distal colon and rectal
 5 cancer incidence and scores of adherence to Western, Prudent and Mediterranean diet excluding in
 6 situ cases.

			Proximal n=447		Distal n=487		Rectal n=546		
	Co	Ca	OR(95%CI)	Ca	OR(95%CI)	Ca	OR(95%CI)	p-het	
WESTERN									
Q1	772	105	1	82	1	96	1		
Q2	824	110	1.01 (0.75;1.37)	133	1.61 (1.19;2.19)	131	1.26 (0.94;1.70)		
Q3	831	109	1.09 (0.80;1.49)	125	1.66 (1.21;2.28)	157	1.62 (1.20;2.18)		
Q4	813	123	1.16 (0.83;1.63)	147	1.99 (1.41;2.80)	162	1.44 (1.04;2.00)		
p-trend			0.325		<0.001		0.013		
1SD-increase			1.07 (0.94;1.22)		1.28 (1.13;1.45)		1.22 (1.08;1.38)	0.085	
PRUDENT									
Q1	783	111	1	129	1	149	1		
Q2	812	110	0.91 (0.68;1.22)	118	0.86 (0.65;1.15)	121	0.77 (0.58;1.01)		
Q3	826	115	1.02 (0.75;1.37)	114	0.91 (0.68;1.22)	146	0.99 (0.75;1.30)		
Q4	819	111	0.91 (0.66;1.27)	126	1.01 (0.74;1.38)	130	0.80 (0.59;1.08)		
p-trend			0.777		0.899		0.409		
1SD-increase			0.99 (0.87;1.12)		0.98 (0.87;1.11)		0.92 (0.83;1.03)	0.613	
MEDITERRANEAN									
Q1	796	97	1	115	1	132	1		
Q2	821	111	0.93 (0.69;1.26)	128	0.96 (0.72;1.27)	130	0.84 (0.64;1.11)		
Q3	816	106	0.75 (0.55;1.03)	113	0.73 (0.54;0.98)	129	0.67 (0.51;0.90)		
Q4	807	133	0.71 (0.51;0.99)	131	0.66 (0.48;0.91)	155	0.58 (0.43;0.79)		
p-trend			0.019		0.003		<0.001		
1SD-increase			0.89 (0.79;1.01)		0.88 (0.78;0.99)		0.83 (0.74;0.93)	0.596	

7
 8
 9
 10

Table 4. Adjusted odds ratios for the association between proximal colon, distal colon and rectal cancer incidence and scores of adherence to Western, Prudent and Mediterranean dietary patterns.

			Proximal Colon		Distal Colon		Rectum		
			n=457		n=503		n=560		
	Co	Ca	aOR ^b (95%CI)	Ca	aOR ^b (95%CI)	Ca	aOR ^b (95%CI)	p-het	
WESTERN									
Quartiles									
Q1	772	108	1	84	1	98	1		
Q2	824	111	1.00 (0.75;1.35)	141	1.70 (1.26;2.29)	137	1.30 (0.97;1.74)		
Q3	831	110	1.07 (0.79;1.46)	128	1.67 (1.22;2.29)	159	1.60 (1.19;2.15)		
Q4	813	128	1.19 (0.85;1.66)	150	2.02 (1.44;2.84)	166	1.46 (1.05;2.01)		
p-trend			0.275		<0.001		0.013		
1SD-increase			1.07 (0.95;1.22)		1.28 (1.13;1.45)		1.23 (1.09;1.38)	0.087	
PAF^c%			7% (-12%;25%)		40% (21%;60%)		27% (11%;44%)		
PRUDENT									
Quartiles									
Q1	783	114	1	132	1	151	1		
Q2	811	113	0.92 (0.69;1.24)	123	0.91 (0.69;1.19)	124	0.79 (0.60;1.03)		
Q3	827	117	1.01 (0.75;1.37)	118	0.94 (0.71;1.26)	151	1.02 (0.78;1.33)		
Q4	819	113	0.92 (0.67;1.28)	130	1.06 (0.78;1.44)	134	0.83 (0.62;1.12)		
p-trend			0.798		0.680		0.545		
1SD-increase			0.98 (0.87;1.11)		1.00 (0.88;1.12)		0.94 (0.84;1.05)	0.686	
PAF^c%			4% (-15%;24%)		-8% (-28%;12%)		9% (-9%;28%)		
MEDITERRANEAN									
Quartiles									
Q1	796	100	1	124	1	133	1		
Q2	821	113	0.92 (0.68;1.24)	131	0.92 (0.70;1.22)	136	0.87 (0.66;1.15)		
Q3	815	109	0.75 (0.55;1.03)	115	0.71 (0.53;0.95)	133	0.70 (0.53;0.93)		
Q4	808	135	0.70 (0.51;0.97)	133	0.65 (0.48;0.89)	158	0.60 (0.45;0.81)		
p-trend			0.017		0.002		<0.001		
1SD-increase			0.89 (0.78;1.00)		0.88 (0.78;0.99)		0.84 (0.75;0.94)	0.746	
PAF^c%			16% (-3%;34%)		20% (3%;38%)		24% (9%;38%)		

^a Co: Controls; Ca: Cases; SD: Standard Deviation.

^b Odds ratio of colon and rectal cancer associated to the adherence to the Western, Prudent and Mediterranean diet patterns adjusted by sex, age, education, BMI, family history of colorectal cancer, physical activity, smoking status, caloric intake and alcohol intake and province of residence as fixed effects.

^c PAF= Population attributable fraction. Proportion of colorectal cancer cases that could be prevented if all participants were in the most beneficial category of adherence to each pattern (Q1 for Western and Q4 for Prudent and Mediterranean)

$$PAF = \frac{PF_{Q1} \cdot (OR_{Q1} - 1) + PF_{Q2} \cdot (OR_{Q2} - 1) + PF_{Q3} \cdot (OR_{Q3} - 1) + PF_{Q4} \cdot (OR_{Q4} - 1)}{1 + [PF_{Q1} \cdot (OR_{Q1} - 1) + PF_{Q2} \cdot (OR_{Q2} - 1) + PF_{Q3} \cdot (OR_{Q3} - 1) + PF_{Q4} \cdot (OR_{Q4} - 1)]} \cdot 100$$

PF=Proportion of population in the specific exposure category

OR= Odds ratio for the especific exposure category