

ORIGINAL ARTICLE

## Urban air pollution and hospital admissions for asthma and acute respiratory disease in Murcia city (Spain)<sup>☆</sup>



Juan Antonio Ortega-García<sup>a,b,\*</sup>, Indra Martínez-Hernández<sup>b</sup>, Elena Boldo<sup>c</sup>, Alberto Cárceles-Álvarez<sup>b</sup>, Carmen Solano-Navarro<sup>d</sup>, Rebeca Ramis<sup>c</sup>, Estefanía Aguilar-Ros<sup>b</sup>, Manuel Sánchez-Solis<sup>e</sup>, Fernando López-Hernández<sup>f</sup>

<sup>a</sup> Comité de Salud Medioambiental, Asociación Española de Pediatría (AEP), Spain

<sup>b</sup> Unidad de Salud Medioambiental, Servicio de Pediatría, Hospital Clínico Universitario Virgen de la Arrixaca, Salud y Ambiente Lab, IMIB-Arrixaca, Universidad de Murcia, Murcia, Spain

<sup>c</sup> Unidad de Epidemiología Ambiental y del Cáncer, Centro Nacional de Epidemiología, Instituto de Salud Carlos III, Madrid, Spain. CIBER en Epidemiología y Salud Pública-CIBERESP, Madrid, Spain

<sup>d</sup> Sección de Urgencias, Servicio de Pediatría, Hospital Clínico Universitario Virgen de la Arrixaca, Murcia, Spain

<sup>e</sup> Sección de Neumología Pediátrica, Servicio de Pediatría, Hospital Clínico Universitario Virgen de la Arrixaca, Murcia, Spain

<sup>f</sup> Departamento de Métodos Cuantitativos, Universidad Politécnica de Cartagena, Cartagena, Murcia, Spain

Received 18 November 2019; accepted 16 January 2020

Available online 14 July 2020

### KEYWORDS

Air pollution;  
Environmental health;  
Hospital admission;  
Asthma;  
Respiratory disease;  
Risk factors

### Abstract

**Introduction:** Urban air pollution (UAP) is a major threat to child and adolescent health. Children are more vulnerable to its effects, being associated with higher morbidity and mortality due to acute and chronic diseases, especially respiratory ones. A study is performed on the relationship between UAP and the rate of hospital admissions due to acute respiratory diseases.

**Patients and methods:** An ecological study was conducted on young people under 19 years-old in the city of Murcia, who had visited hospital emergency departments due to respiratory diseases (ICD-9) during 2015. A logistic regression was performed on the risk of hospital admission that included consultations in relation to the average daily levels of environmental pollutants (NO<sub>2</sub>, O<sub>3</sub>, PM<sub>10</sub>, SO<sub>2</sub>) obtained from the Air Quality Surveillance and Control network of the Region of Murcia. Other control variables, such as gender, age, average daily ambient temperature, and season of the year.

**Results:** A total of 12,354 (56% boys and 44% girls) children consulted in the emergency department for respiratory disease. Of those, 3.5% were admitted, with a mean age of 2.54 (95% CI; 2.16–2.91) years. The Odds Ratio (OR) of hospital admission for respiratory diseases: NO<sub>2</sub> 1.02

<sup>☆</sup> Please cite this article as: Ortega-García JA, Martínez-Hernández I, Boldo E, Cárceles-Álvarez A, Solano-Navarro C, Ramis R, et al. Contaminación atmosférica urbana e ingresos hospitalarios por asma y enfermedades respiratorias agudas en la ciudad de Murcia (España). An Pediatr (Barc). 2020;93:95–102.

\* Corresponding author.

E-mail address: [ortega@pehsu.org](mailto:ortega@pehsu.org) (J.A. Ortega-García).

**PALABRAS CLAVE**

Contaminación del aire;  
Salud medioambiental;  
Ingresos hospitalarios;  
Asma;  
Enfermedad respiratoria;  
Factores de riesgo

(95% CI; 1.01–1.04;  $P < .01$ ),  $O_3$  1.01 (95% CI; 1.00–1.03;  $P < .01$ ) male 1.4 (95% CI 1.11–1.79;  $P < .01$ ) and winter 2.10 (95% CI 1.40–3.21;  $P < .01$ ). Admissions for asthma: PM10 1.02 (95% CI; 1.01–1.04;  $P < .05$ ),  $O_3$  1.04 (95% CI; 1.01–1.06;  $P < .01$ ). Admissions for bronchiolitis: Age 0.69 (95% CI; 0.48–0.99;  $P < .05$ );  $NO_2$  1.03 (95% CI; 1.01–1.05;  $P < .01$ ).

**Conclusions:** UAP increases hospital admissions in children due to acute respiratory diseases, especially asthma and bronchiolitis. Implementing preventive measures, expanding time series and collaborative studies with open data, would help improve public health and air quality in the cities.

© 2020 Published by Elsevier España, S.L.U. on behalf of Asociación Española de Pediatría. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Contaminación atmosférica urbana e ingresos hospitalarios por asma y enfermedades respiratorias agudas en la ciudad de Murcia (España)

**Resumen**

**Introducción:** La contaminación atmosférica urbana (CAU) es una amenaza principal para la salud infanto-juvenil. Los niños son más vulnerables a sus efectos asociando mayor morbimortalidad de enfermedades agudas y crónicas, especialmente respiratorias. Pretendemos estudiar los efectos de la CAU en la tasa de ingresos hospitalarios por causa aguda respiratoria.

**Pacientes y métodos:** Estudio ecológico durante 2015 de las visitas con patología respiratoria (CIE-9) de urgencias hospitalarias en menores de 19 años en el municipio de la ciudad de Murcia. Regresión logística para el riesgo de ingreso hospitalario entre las consultas en relación con los niveles promedios diarios de contaminantes ambientales ( $NO_2$ ,  $O_3$ , PM10,  $SO_2$ ) obtenidas de la red de Vigilancia y Control de la Calidad del Aire de la Región de Murcia. Otras variables de control: sexo, edad, temperatura ambiental media diaria, y estación del año.

**Resultados:** Un total de 12,354 niños consultaron en urgencias por patología respiratoria, 56% niños y 44% niñas. Ingresaron el 3,5%, con edad media de 2,54 (IC95% 2,16–2,91) años. Para el riesgo de ingreso hospitalario por enfermedades respiratorias:  $NO_2$  1,02 (IC95% 1,01–1,04;  $P < ,01$ ),  $O_3$  1,01 (IC95% 1,00–1,03;  $P < ,01$ ) sexo masculino 1,4 (IC95% 1,11–1,79;  $P < ,01$ ) e invierno 2,10 (IC95% 1,40–3,21;  $P < ,01$ ). Ingresos por asma: PM10 1,02 (IC95% 1,01–1,04;  $P < ,05$ ),  $O_3$  1,04 (IC95% 1,01–1,06;  $P < ,01$ ) Ingresos por bronquiolitis: edad 0,69 (IC95% 0,48–0,99;  $P < ,05$ );  $NO_2$  1,03 (IC95% 1,01–1,05;  $P < ,01$ ).

**Conclusiones:** La CAU aumenta los ingresos hospitalarios en pediatría por patología aguda respiratoria, especialmente por crisis de asma y bronquiolitis. Poner en marcha medidas preventivas, ampliar las series temporales y estudios colaborativos con datos en abierto ayudarían a mejorar la salud pública y la calidad del aire en las ciudades.

© 2020 Publicado por Elsevier España, S.L.U. en nombre de Asociación Española de Pediatría. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Introduction**

Urban air pollution (UAP) is one of the main environmental health threats for the paediatric age group.<sup>1</sup> Although the entire population is affected by poor air quality, children and youth are more vulnerable to its effects due to the physiological immaturity of their respiratory and immune systems, their higher metabolism, higher respiratory rate and greater requirements of air per kilogram of body weight compared to adults.<sup>2</sup> A growing body of scientific evidence demonstrates the contribution of UAP to the increased morbidity and mortality of children worldwide.<sup>3,4</sup> Exposure to UAP during childhood has been associated with an increase in respiratory complaints and associated hospital admissions, inflammation of the airways, impaired respiratory function, asthma attacks and acute lower airway infections.<sup>5–15</sup>

The World Health Organization (WHO) regularly publishes guidelines on air quality that constitute the main reference to establish standards worldwide.<sup>16</sup> It also offers information on the air quality of more than 3000 cities in 103 countries (<http://breathelife2030.org/>). Table 1 presents the main types of air pollutants highlighted in studies of respiratory morbidity<sup>3</sup> and the maximum levels recommended by the WHO.<sup>16</sup> The primary components of UAP involved in morbidity are particulate matter, including fine particles 2.5  $\mu\text{m}$  in diameter and smaller (PM2.5) or 10  $\mu\text{m}$  in diameter and smaller (PM10), nitrogen dioxide ( $NO_2$ ), sulphur dioxide ( $SO_2$ ), and ozone ( $O_3$ ). Tropospheric  $O_3$  is found in the lowest layers of the atmosphere and is considered a secondary pollutant, as it is not directly emitted to the atmosphere but rather forms from certain precursors (volatile nonmethane organic compounds, carbon

**Table 1** Air pollutants that affect the respiratory tract, main sources of these pollutants and mean value limits recommended by the WHO.

	Sources of emissions	WHO limit
Particulate matter	Combustion/burning	Annual mean 10 $\mu\text{g}/\text{m}^3$ 24-h mean 25 $\mu\text{g}/\text{m}^3$
Particulate matter <10 $\mu\text{m}$ in diameter (PM10)	Construction, dust resuspended in roads, traffic and wind	Annual mean 20 $\mu\text{g}/\text{m}^3$  24-h mean 50 $\mu\text{g}/\text{m}^3$
Nitrogen dioxide (NO <sub>2</sub> )	Emissions from diesel fuels and associated combustion associated with motor vehicle traffic	Annual mean 40 $\mu\text{g}/\text{m}^3$  1-h mean 200 $\mu\text{g}/\text{m}^3$
Sulphur dioxide (SO <sub>2</sub> )	Combustion of fossil fuels in industrial plants	24-h mean 20 $\mu\text{g}/\text{m}^3$ 10-min mean 500 $\mu\text{g}/\text{m}^3$
Ozone (O <sub>3</sub> )	Photochemical reactions in presence of light and oxides or volatile organic compounds	8-h mean 100 $\mu\text{g}/\text{m}^3$

monoxide [CO], nitrogen oxides and, to a lesser degree, methane [CH<sub>4</sub>]) produced by combustion processes (traffic and industry). Through the action of sunlight, these chemical substances react to form O<sub>3</sub>. Since sunlight is one of the main factors at play in these reactions, the concentrations of these pollutants peak in spring and summer.

The aim of our study was to analyse the association between the level of UAP and hospital admissions in individuals aged less than 18 years due to acute respiratory illness in the city of Murcia, Spain.

## Sample and methods

### Study sample

The city of Murcia has a population of 441 003 inhabitants (2016), including 100 539 aged less than 19 years.<sup>17</sup> The Paediatric Emergency Unit (PED) of the Hospital Clínico Universitario Virgen de la Arrixaca is the only hospital-based paediatric emergency care unit in the city of Murcia in the Murcia public health system.

The population under study consisted of all children aged less than 17 years residing in the city and urban area of Murcia that visited the emergency department of the Hospital Clínico Universitario Virgen de la Arrixaca between January 1, 2015 and December 31, 2015 given any diagnosis of upper or lower respiratory tract infection based on the recorded International Classification of Diseases, Ninth Revision (ICD-9) code. The ICD-9 and hospital admission code were registered by the Department of Clinical Documentation following discharge.

Study variables: hospital admission (yes/no), sex, age in years, grouped into categories (<1 year, 1–3 years, 4–9 years, >9 years), ICD-9 diagnostic code and season of the year.

The study protocol was reviewed and approved by the Clinical Research Ethics Committee of the Hospital Clínico Universitario Virgen de la Arrixaca.

### Geographic location and urban air pollution data

Murcia is a town in southeast Spain on the banks of the Segura river in the coastal depression of the Sierra de Carascoy, 40 km from the Mediterranean, with mild winds and a semi-arid dry climate. Its winters are mild and its summers very hot, with an annual mean temperature of 18.6 °C. The area does not have any refineries, thermal power stations or incineration plants based on the *Registro Estatal de Emisiones y Fuentes Contaminantes* (National Registry of Emissions and Pollution Sources, <http://www.prtr-es.es>), has experienced considerable urban development and a rapid population increase in the past 25 years. The main sources of pollution are motor vehicle traffic and the burning of biomass or farming residues associated with traditional agriculture in the huerta of Murcia (PM10 and NO<sub>2</sub>). The periodic intrusion of Saharan dust sometimes causes variations in PM10 levels. The intensity of sunlight and frequent heat in Murcia, combined with UAP and the orography of the area have contributed to an increasing frequency of episodes of excessive levels of the secondary pollutant O<sub>3</sub>.

We obtained data on environmental pollutants from January 1, 2015 to December 31, 2015 from the website of the Air Quality Surveillance and Control Network of the General Directorate of the Environment of the Region of Murcia.<sup>18</sup> It has a measuring station in the square of San Basilio in the city of Murcia. The UAP data obtained for the months under study included the following pollutants: NO<sub>2</sub>, PM10, O<sub>3</sub>, SO<sub>2</sub>. The levels of PM2.5 are not measured. The mean daily temperature was based on the measurements made by the station of the Air Quality Surveillance and Control Network (Murcia-Alcantarilla). We compared these values with the air quality standards of the WHO.

Variables under study: NO<sub>2</sub>, PM10, O<sub>3</sub>, SO<sub>2</sub> (daily mean), daily mean temperature, month and season of the year of visit.

### Statistical analysis

We performed a descriptive analysis with frequency and contingency tables. We fitted a multivariate logistic regression model to assess the risk of admission with a 95% level

**Table 2** Sociodemographic variables and univariate analysis.

	Discharge home		Admitted to hospital		ANOVA / CW/ $\chi^2$ *
	n (%)	Mean (95% CI)	n (%)	Mean (95% CI)	
Sex					.05
Male	6661 (96.2)		264 (3.8)		
Female	5257 (96.8)		172 (3.2)		
Age (years)		3.11 (3.06–3.16)		2.54 (2.16–2.91)	< .01
<1	2961 (24.0)		257 (2.1)		
1–3	5726 (46.3)		91 (0.7)		
4–9	2739 (22.2)		50 (0.4)		
>=10	492 (4.0)		38 (0.3)		
Mean T (°C)		17.35 (17.23–17.47)		15.26 (14.68–15.84)	< .01
Season of year					< .01
Winter	4850 (40.7)		255 (58.5)		
Spring	2509 (21.1)		76 (17.4)		
Summer	2375 (19.9)		49 (11.2)		
Autumn	2184 (18.3)		56 (12.8)		
UAP (daily mean $\mu\text{g}/\text{m}^3$ )					
NO <sub>2</sub>		46.92 (46.60–47.23)		53.88 (52.06–55.69)	< .01
PM10		34.81(34.47–35.16)		37.13 (35.01–39.25)	.02
O <sub>3</sub>		44.35 (43.95–44.75)		38.27 (36.16–40.38)	< .01
SO <sub>2</sub>		7.84 (7.80–7.88)		7.61 (7.39–7.82)	.02

CI, confidence interval; UAP, urban air pollution; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone; PM10, particulate matter with diameter  $\leq 10 \mu\text{m}$ ; SO<sub>2</sub>, sulphur dioxide; T, temperature.

Significant variables ( $P < .05$ ) presented in boldface.

\* =Chi-Square.


of confidence. We performed bidirectional tests to prevent confounding due to seasonal variations.<sup>19</sup>

## Results

The PED received 12 354 visits of residents of the municipality of Murcia due to acute upper or lower respiratory tract disease in 2015. Cold/upper airway infection, asthma/bronchospasm, pharyngitis or sinusitis/abscesses, otitis, laryngitis, bronchiolitis and pneumonia/bronchitis amounted to 38.7%. 18.9%. 14.9%. 10.0%. 7.9%. 6.9% and 2.6, respectively. A total of 436 patients (corresponding to 3.5% of visits for respiratory disease) were admitted to hospital; Fig. 1 shows the distribution of hospital admissions by respiratory disease. Table 2 presents the profiles of the patients that sought care and data on air pollution.

The annual mean level of NO<sub>2</sub> in the municipality of Murcia, of 46.92  $\mu\text{g}/\text{m}^3$  (95% confidence interval, 46.60–47.23), exceeds the limit recommended by the WHO (40  $\mu\text{g}/\text{m}^3$ ), with levels exceeding this recommended limit 52% of days a year.<sup>20</sup> The annual mean of PM10 is 34.81  $\mu\text{g}/\text{m}^3$  (34.47–35.16), and the levels of PM10 exceed the annual mean recommended by the WHO (50  $\mu\text{g}/\text{m}^3$ ) 10% of the days. Fig. 2 presents the annual distribution of daily mean values for the 4 pollutants.

Table 3 presents the results of the multivariate logistic regression analysis. The risk of hospital admission due to respiratory complaints was greater in male patients, increased with the mean daily level of NO<sub>2</sub> and O<sub>3</sub> and decreased with age.




**Figure 1** Distribution of hospital admissions due to respiratory complaints in 2015 in the city of Murcia.

## Discussion

Our findings reveal that a higher short-term level of UAP (O<sub>3</sub> and NO<sub>2</sub>), male sex and winter were associated with an increased risk of hospital admission due to respiratory illness. This association was strongest for asthma (with O<sub>3</sub> and PM10) and bronchiolitis (with NO<sub>2</sub>).

The increased hospital admission rate in male patients has already been described in the previous literature,<sup>14,15</sup>



**Figure 2** Distribution of the daily mean of air pollutants in the city of Murcia throughout the year.

although it diverges from the increased vulnerability to  $O_3$  described in female patients.<sup>21</sup> Some of the possible explanations proposed for this are models of exposure (boys tend to play outdoors for longer periods and engage in more aerobic activity) and physiological (differences in hormones, lung size, inflammation of airways).

From a physiological standpoint, it makes sense that air pollutants could increase the severity of asthma and bronchiolitis on account of their known effect on lung function and airway inflammation.<sup>22</sup>

There is evidence that short-term exposure to particulate matter,  $O_3$ ,<sup>18</sup>  $SO_2$  and  $NO_2$  is associated to an increase in asthma exacerbations.<sup>10</sup> The effect of  $PM_{10}$  on asthma could be explained by the oxidative stress induced by the presence of  $PM_{10}$  in the epithelium, with an increase in prostaglandin E that leads to defective macrophage phagocytosis, inflammation, oedema and cytotoxicity. The evidence on hospital admissions due to asthma in relation to  $O_3$  exposure requires further research,<sup>14,23,24</sup> but a review of 87 studies found that 71 reported a significant increase in the risk of hospital admission due to asthma.<sup>15</sup>

Bronchiolitis is an infection of the small airways that mainly affects children aged less than 2 years. It causes inflammation of the airways and necrosis of the bronchial epithelium.<sup>25</sup> In 80% of cases it is associated with infection by respiratory syncytial virus, and it is one of the main reasons for hospital admission in children aged less than 2 years in both high-income<sup>26</sup> and low-income<sup>27</sup> countries. Although there are isolated cases throughout the year, there are usually one to two outbreaks of bronchiolitis in the cold

months of the year, between November and March. The risk factors known to increase the risk of hospital admission are preterm birth, low birth weight, heart disease or exposure to tobacco smoke. We found an association between the mean daily level of  $NO_2$  and the risk of hospital admission due to bronchiolitis. The current evidence suggests an association between short (<24 h) and recent exposure to  $PM_{2.5}$ ;  $PM_{10}$ ;  $NO_2$  and  $SO_2$ , among other pollutants, and an increased incidence of lower respiratory tract infections<sup>11</sup> and an increased risk of hospital admission due to bronchiolitis.<sup>28</sup> The greater effect observed in other studies with chronic exposure to particulate matter (several days or weekly mean) on hospital admission due to bronchiolitis may reflect different pathogenic mechanisms.<sup>29</sup> Chronic exposure to  $PM_{2.5}$  and  $PM_{10}$  may have a chronic proinflammatory effect,<sup>30</sup> whereas exposure to  $NO_2$  and  $SO_2$  may be associated with more acute airway damage.<sup>31</sup>

Ozone levels usually peak during hot seasons, which are far apart from the periods when outbreaks of infection by respiratory syncytial virus tend to occur. This would make it difficult to identify an association with bronchiolitis. In winter, other contaminants, such as  $NO_2$ , are probably more relevant contributors to respiratory disease. As has been done by other authors,<sup>32</sup> we analysed synergistic effects and interactions in a model taking into account multiple pollutants and age groups, and we did not find any differences in the results.

In our study, the annual mean level of  $NO_2$  in the city of Murcia exceeded the recommended limit for the health of its residents. Murcia is the city that ranks 8th

**Table 3** Multivariate logistic regression model. Risk of hospital admission due to acute respiratory illness.

## Risk of hospital admission / all admissions due to respiratory illness

	B	SE	Wald	Sig	Exp (B)	Lower bound	Upper bound
Age (years)	-0.035	0.021	2.661	0.103	0.966	0.926	1.007
Sex (male)	0.346	0.121	8.260	0.004	1.414	1.116	1.791
T (°C)	-0.003	0.019	0.033	0.856	0.997	0.960	1.035
NO <sub>2</sub> (µg/m <sup>3</sup> )	-0.024	0.006	16.786	0.000	1.024	1.013	1.036
O <sub>3</sub> (µg/m <sup>3</sup> )	0.016	0.006	6.493	0.001	1.017	1.004	1.030
PM10 (µg/m <sup>3</sup> )	-0.005	0.004	1.313	0.252	0.995	0.987	1.003
SO <sub>2</sub> (µg/m <sup>3</sup> )	-0.021	0.028	0.602	0.438	0.979	0.927	1.033
Season of year			15.421	0.001			
Winter	0.744	0.215	11.993	0.001	2.105	1.403	3.211
Spring	0.130	0.246	0.279	0.597	1.139	0.879	1.846
Summer	-0.158	0.296	0.284	0.594	0.854	0.478	1.525
Constant	-5.285	0.583	82.133	0.000	0.005		

## Risk of hospital admission / asthma

	B	SE	Wald	Sig	Exp (B)	Lower bound	Upper bound
Age (years)	0.040	0.036	1.221	0.269	1.041	0.969	1.118
Sex (male)	0.481	0.268	3.217	0.073	1.618	0.956	2.738
T (°C)	-0.028	0.040	0.495	0.482	0.972	0.898	1.052
NO <sub>2</sub> (µg/m <sup>3</sup> )	0.014	0.012	1.356	0.244	1.014	0.990	1.039
O <sub>3</sub> (µg/m <sup>3</sup> )	0.036	0.013	7.835	0.005	1.037	1.011	1.063
PM10 (µg/m <sup>3</sup> )	0.019	0.008	5.194	0.023	1.019	1.003	1.036
SO <sub>2</sub> (µg/m <sup>3</sup> )	0.005	0.062	0.008	0.931	1.005	0.891	1.134
Season of year			2.691	0.442			
Winter	-0.017	0.448	0.001	0.970	0.983	0.409	2.365
Spring	-0.710	0.502	1.998	0.158	0.492	0.184	1.316
Summer	-0.424	0.544	0.609	0.435	0.654	0.225	1.899
Constant	-5.970	1.195	24.958	0.000	0.003		

## Risk of hospital admission / bronchiolitis

	B	SE	Wald	Sig	Exp (B)	Lower bound	Upper bound
Age (years)	-0.365	0.184	3.927	0.048	0.694	0.484	0.996
Sex (male)	0.377	0.206	3.354	0.067	1.457	0.974	2.180
T (°C)	-0.006	0.035	0.034	0.854	0.994	0.928	1.064
NO <sub>2</sub> (µg/m <sup>3</sup> )	0.031	0.010	10.031	0.002	1.032	1.012	1.052
O <sub>3</sub> (µg/m <sup>3</sup> )	0.019	0.011	2.953	0.086	1.019	0.997	1.041
PM10 (µg/m <sup>3</sup> )	-0.008	0.007	1.497	0.221	0.992	0.979	1.005
SO <sub>2</sub> (µg/m <sup>3</sup> )	0.001	0.046	0.001	0.979	1.001	0.915	1.096
Season of year			4.254	0.235			
Winter	0.342	0.375	0.832	0.362	1.408	0.675	2.938
Spring	0.547	0.417	1.721	0.190	1.729	0.763	3.916
Summer	-0.920	0.884	1.084	0.298	0.398	0.070	2.252
Constant	-3.388	1.114	9.252	0.002	0.034		

Input variable(s): NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone; PM10, particulate matter ≤10 µm; SO<sub>2</sub>, sulphur dioxide; daily mean, season of year, daily mean temperature in the city of Murcia, age in years, sex.

in Spain in terms of the highest annual mean level of NO<sub>2</sub>. The increasing urbanization and concentration of the population in megacities,<sup>33</sup> in absence of corresponding air quality control measures, has a direct impact on child health. At present, 98% of cities in low- and medium-income countries and 56% of cities in high-income countries have levels of pollution exceeding the limits recommended by the WHO.<sup>34</sup>

One significant limitation of our study involves the scarcity of sources of pollution data. The municipality of Murcia has 400 000 inhabitants, making it the seventh most populous municipality in Spain, yet it only has one pollutant measurement station, which does not collect information for PM<sub>2.5</sub>, among other pollutants. The season and the temperature play a role in the variations in levels of UAP, so we controlled our analyses for these factors.<sup>35–39</sup> Other

population- and individual-based variables, such as smoking, dietary factors and lifestyle, may be a source of bias. Smoking is particularly important, given that approximately 50% of children in Spain live with a smoker at home.<sup>40</sup> The low level of documentation and the lack of an established coding scheme to document exposure to second-hand smoke in minors that visited the PED precluded the analysis of its impact on our findings. We believe it essential to train paediatricians in the use of brief opportunistic intervention during PED visits. On the other hand, additional collaborative cohort and longitudinal studies with a standardised methodological approach are needed to better analyse the associations identified in our study.

Although the proportion of patients requiring hospital admission due asthma or bronchiolitis that could be attributed to UAP may seem small, UAP probably has a substantial impact in morbidity and the global burden of disease at the population level. The good news is that exposure to UAP is totally preventable. Successful interventions are already being developed and implemented in numerous cities and regions across the globe. The reduction of UAP has been associated with public health benefits, with a decrease in mortality associated to cardiovascular and respiratory diseases in adults and a decrease in the rate of hospital admission due to respiratory complaints in children (of up to 87% in cities that started out with high levels of pollutants), a decrease in the incidence of preterm birth and a reduction in lung inflammation.<sup>33,41–45</sup> Thus, improving air quality surveillance systems and implementing measures for control of UAP (public transport, urban design, banning brush burning, promoting cycling, limiting the use of high-emission motor vehicles, improving monitoring of UAP and its effects in vulnerable groups, among others) contribute to reducing the rate of hospital admission in the paediatric age group.

Such a pathway requires courageous leadership, the creation of new structures that connect health and environment, substantial resources and widespread changes in society.<sup>40</sup> The growing social awareness of the association between health and environment is a key driver of change. We take this opportunity to ask Spanish paediatricians to help increase the sample size, providing free access to our data (<http://pehsu.org/wp-content/uploads/UAP.IngresosRespi.zip>) to other paediatrics research groups in Spain. The combined analysis of clinical and environmental surveillance data is a powerful tool toward achieving adequate monitoring and for creating models that can contribute to improving health and air quality in cities.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## Acknowledgments

We thank Dr Picazo Román and his entire clinical documentation team at the Hospital Clínico Universitario Virgen de la Arrixaca for their collaboration in the project. We thank the groups Madres por el Clima Murcia (Murcia Mothers for Climate) and Stop Quemadas Murcia (Stop Burning Murcia) for

their work in raising social awareness of the need to improve air quality in the city of Murcia.

## References

- Ortega-García JA, Sánchez-Solís M, Ferris-Tortajada J. Air pollution and children's health. *An Pediatr (Barc)*. 2018;89:77–9.
- Schwartz J. Air pollution and children's health. *Pediatrics*. 2004;113 4 Suppl::1037–43.
- Unicef. Clear the air for children: the impact of air pollution on children. New York: UNICEF; 2016.
- Global Burden of Disease 2016 Risk Factors Collaborators. Global Burden of Disease 2016 Risk Factors Collaborators Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390:1345–422.
- Goldizen FC, Sly PD, Knibbs LD. Respiratory effects of air pollution on children. *Pediatr Pulmonol*. 2016;51:94–108.
- Gehring U, Gruzieva O, Agius RM, Beelen R, Custovic A, Cyrys J, et al. Air pollution exposure and lung function in children: the ESCAPE project. *Environ Health Perspect*. 2013;121:1357–64.
- Luong LMT, Phung D, Dang TN, Sly PD, Morawska L, Thai PK. Seasonal association between ambient ozone and hospital admission for respiratory diseases in Hanoi, Vietnam. *PLoS One*. 2018;13:e0203751.
- Bono R, Romanazzi V, Bellisario V, Tassinari R, Trucco G, Urbino A, et al. Air pollution, aeroallergens and admissions to pediatric emergency room for respiratory reasons in Turin, northwestern Italy. *BMC Public Health*. 2016;16:722.
- Nicolai T. Environmental air pollution and lung disease in children. *Monaldi Arch Chest Dis*. 1999;54:475–8.
- Orellano P, Quaranta N, Reynoso J, Balbi B, Vasquez J. Effect of outdoor air pollution on asthma exacerbations in children and adults: systematic review and multilevel meta-analysis. *PLoS One*. 2017;12:e0174050.
- Mehta S, Shin H, Burnett R, North T, Cohen AJ. Ambient particulate air pollution and acute lower respiratory infections: a systematic review and implications for estimating the global burden of disease. *Air Qual Atmos Health*. 2013;6:69–83.
- Götschi T, Heinrich J, Sunyer J, Künzli N. Long-term effects of ambient air pollution on lung function: a review. *Epidemiology*. 2008;19:690–701.
- Schultz ES, Gruzieva O, Bellander T, Bottai M, Hallberg J, Kull I, et al. Traffic-related air pollution and lung function in children at 8 years of age: a birth cohort study. *Am J Respir Crit Care Med*. 2012;186:1286–91.
- Kuo CY, Chan CK, Wu CY, Phan DV, Chan CKL. The short-term effects of ambient air pollutants on childhood asthma hospitalization in Taiwan: a national study. *Int J Environ Res Public Health*. 2019;16:203.
- Zheng XY, Ding H, Jiang LN, Chen SW, Zheng JP, Qiu M, et al. Association between air pollutants and asthma emergency room visits and hospital admissions in time series studies: a systematic review and meta-analysis. *PLoS One*. 2015;10:e0138146.
- World Health Organization (WHO) (2006) World Health Organization (WHO) Air quality guidelines. [http://www.euro.who.int/\\_data/assets/pdf\\_file/0005/78638/E90038.pdf](http://www.euro.who.int/_data/assets/pdf_file/0005/78638/E90038.pdf), 2006.
- Instituto Nacional de Estadística. Estadística del Padrón Continuo a 1 de enero de 2016. Datos por municipios. Murcia [Accessed 18 Oct 2019]. Available from: <https://www.ine.es/jaxi/Datos.htm?path=/t20/e245/p05/a2016/l0/&file=00030001.px>.
- Red de Vigilancia y Control de la Calidad del Aire de la Dirección General de Medio Ambiente de la Comunidad Autónoma

- de la Región de Murcia [Accessed 15 Sep 2019]. Available from: <https://sinqlair.carm.es/calidadaire/>.
19. Bateson TF, Schwartz J. Control for seasonal variation and time trend in case-crossover studies of acute effects of environmental exposures. *Epidemiology*. 1999;10:539–44.
  20. Air pollution country profile. Spain. European Environment Agency [Accessed 18 One 2018]. Available from: <https://www.eea.europa.eu/themes/air/country-fact-sheets/spain>.
  21. Mikerov AN, Phelps DS, Gan X, Umstead TM, Haque R, Wang G, et al. Effect of ozone exposure and infection on bronchoalveolar lavage: sex differences in response patterns. *Toxicol Lett*. 2014;230:333–44.
  22. Barraza-Villarreal A, Sunyer J, Hernandez-Cadena L, Escamilla-Núñez MC, Sienra-Monge JJ, Ramírez-Aguilar M, et al. Air pollution, airway inflammation, and lung function in a cohort study of Mexico City schoolchildren. *Environ Health Perspect*. 2008;116:832–8.
  23. Gharibi H, Entwistle MR, Ha S, Gonzalez M, Brown P, Schweizer D, et al. Ozone pollution and asthma emergency department visits in the Central Valley, California, USA, during June to September of 2015: a time-stratified case-crossover analysis. *J Asthma*. 2018;9:1–12.
  24. Nhung NTT, Schindler C, Dien TM, Probst-Hensch N, Perez L, Künzli N. Acute effects of ambient air pollution on lower respiratory infections in hanoi children: an eight-year time series study. *Environ Int*. 2018;110:139–48.
  25. Cano-Garcinuño A, Praena-Crespo M, Mora-Gandarillas I, Carvajal-Urueña I, Callén-Blecua MT, García-Merino Á. Grupo de Expertos para el Diagnóstico de la Bronquiolitis. Criterios heterogeneidad en el diagnóstico de aguda bronquiolitis en España. *An Pediatr (Barc)*. 2019;90:109–17.
  26. Green CA, Yeates D, Goldacre A, Sande C, Parslow RC, McShane P, et al. Admission to hospital for bronchiolitis in England: trends over five decades, geographical variation and association with perinatal characteristics and subsequent asthma. *Arch Dis Child*. 2016;101:140–6.
  27. Robertson SE, Roca A, Alonso P, Simoes EA, Kartasasmita CB, Olaleye DO, et al. Respiratory syncytial virus infection: denominator-based studies in Indonesia, Mozambique, Nigeria and South Africa. *Bull World Health Organ*. 2004;82:914–22.
  28. King C, Kirkham J, Hawcutt D, Sinha I. The effect of outdoor air pollution on the risk of hospitalisation for bronchiolitis in infants: a systematic review. *PeerJ*. 2018;6:e5352.
  29. Yitshak-Sade M, Yudovitch D, Novack V, Tal A, Kloog I, Goldberg A. Air pollution and hospitalization for bronchiolitis among young children. *Ann Am Thorac Soc*. 2017;14:1796–802.
  30. Calderón-Garcidueñas L, Mora-Tiscareño A, Fordham LA, Valencia-Salazar G, Chung CJ, Rodríguez-Alcaraz A, et al. Respiratory damage in children exposed to urban pollution. *Pediatr Pulmonol*. 2003;36:148–61.
  31. Chen TM, Gokhale J, Shofer S, Kuschner WG. Outdoor air pollution: nitrogen dioxide, sulfur dioxide, and carbon monoxide health effects. *Am J Med Sci*. 2007;333:249–56.
  32. Karr C, Lumley T, Schreuder A, Davis R, Larson T, Ritz B, et al. Effects of subchronic and chronic exposure to ambient air pollutants on infant bronchiolitis. *Am J Epidemiol*. 2007;165:553–60.
  33. Landrigan PJ, Fuller R, Acosta NJR, Adeyi O, Arnold R, Basu NN, et al. The Lancet Commission on Pollution and health. *Lancet*. 2018;391:462–512.
  34. World Health Organization (WHO) (2016) World Health Organization (WHO) WHO's urban ambient air pollution database—update 2016. 2016. [Accessed 21 Nov 2017]. Available from: [http://www.who.int/phe/health\\_topics/outdoorair/databases/cities/en/](http://www.who.int/phe/health_topics/outdoorair/databases/cities/en/).
  35. Karr CJ, Demers PA, Koehoorn MW, Lencar CC, Tamburic L, Brauer M. Influence of ambient air pollutant sources on clinical encounters for infant bronchiolitis. *Am J Respir Crit Care Med*. 2009;180:995–1001.
  36. Girgis MS, Strickland MJ, Hu X, Liu Y, Chang HH, Belanoff C, et al. Chronic PM(2.5) exposure and risk of infant bronchiolitis and otitis media clinical encounters. *Int J Hyg Environ Health*. 2017;220:1055–63.
  37. Karr CJ, Demers PA, Koehoorn MW, Lencar CC, Tamburic L, Brauer M. Infant exposure to fine particulate matter and traffic and risk of hospitalization for RSV bronchiolitis in a region with lower ambient air pollution. *Environ Res*. 2009;109:321–7.
  38. Ségala C, Poizeau D, Mesbah M, Willems S, Maidenberg M. Winter air pollution and infant bronchiolitis in Paris. *Environ Res*. 2008;106:96–100.
  39. Martín Martín R, Sánchez Bayle M. Impact of air pollution in paediatric consultations in Primary Health Care: ecological study. *An Pediatr (Barc)*. 2018;89:80–5.
  40. Ortega-García JA, Tellerías L, Ferrís-Tortajada J, Boldo E, Campillo-López F, van den Hazel P, et al. Threats, challenges and opportunities for paediatric environmental health in Europe, Latin America and the Caribbean. *An Pediatr (Barc)*. 2019;90:124.e1–11.
  41. Rich DQ, Liu K, Zhang J, Thurston SW, Stevens TP, Pan Y, et al. Differences in birth weight associated with the 2008 Beijing Olympics Air Pollution Reduction: results from a natural experiment. *Environ Health Perspect*. 2015;123:880–7.
  42. Clancy L, Goodman P, Sinclair H, Dockery DW. Effect of air-pollution control on death rates in Dublin, Ireland: an intervention study. *Lancet*. 2002;360:1210–4.
  43. Izquierdo R, García dos Santos S, Borge R, de la Paz D, Sari-giannis D, Gotti A, et al. Health impact assessment by the implementation of Madrid City air-quality plan in 2020. *Environ Res*. 2020;183:1090212.
  44. Zhang Z, Wang J, Liu F, Yuan L, Yuan J, Chen L, et al. Impacts of event-specific air quality improvements on total hospital admissions and reduced systemic inflammation in COPD patients. *PLoS One*. 2019;14:e0208687.
  45. Pope CA 3<sup>rd</sup>. Respiratory disease associated with community air pollution and a steel mill, Utah Valley. *Am J Public Health*. 1989;79:623–8.