

Table S4. Single and multiple enteric protist infections/colonisations detected in faecal samples from non-human primates in the present study.

Protist species combination	Faecal samples (<i>n</i>)	Relative frequency (%)
<i>Blastocystis</i> sp. only	56	30.1
<i>G. duodenalis</i> only	58	31.2
<i>E. dispar</i> only	17	9.1
<i>B. coli</i> only	2	1.1
<i>T. abrasarti</i> only	6	3.2
<i>E. bienewisi</i> only	3	1.6
<i>Cryptosporidium</i> spp. only	1	0.5
<i>Blastocystis</i> sp. + <i>G. duodenalis</i>	15	8.1
<i>Blastocystis</i> sp. + <i>E. dispar</i>	13	7.0
<i>Blastocystis</i> sp. + <i>B. coli</i>	3	1.6
<i>E. dispar</i> + <i>G. duodenalis</i>	3	1.6
<i>B. coli</i> + <i>Cryptosporidium</i> spp.	1	0.5
<i>B. coli</i> + <i>G. duodenalis</i>	1	0.5
<i>Blastocystis</i> sp. + <i>Cryptosporidium</i> spp.	1	0.5
<i>Blastocystis</i> sp. + <i>E. bienewisi</i>	1	0.5
<i>Cryptosporidium</i> spp. + <i>G. duodenalis</i>	1	0.5
<i>G. duodenalis</i> + <i>T. abrasarti</i>	1	0.5
<i>Blastocystis</i> sp. + <i>E. dispar</i> + <i>G. duodenalis</i>	3	1.6
Total	186	100.0