

Supplementary Table 1. Sequence of primers used for qRT-PCR studies.

Gene		Sequence (5' - 3')
<i>Circadian locomotor output cycles kaput (Clock)</i>	Fw	CCTTGCGTCTGTGGGTGTT
	Rv	TGTCATCTTCTTCCACCAATCCA
<i>Aryl hydrocarbon receptor nuclear translocator-like (Arntl)</i>	Fw	CGCCTCTACCTGTTCAAAGAAAAA
	Rv	TCACCCGTATTTCCCCGTTCC
<i>Period circadian clock 1 (Per1)</i>	Fw	AAACGGCAAGCGGATGG
	Rv	GCCATACAGTGGAGGACGAAA
<i>Period circadian clock 2 (Per2)</i>	Fw	CAGAGGAGAAGACTCCGCAC
	Rv	TTGCTGTCGCTGGATGATGT
<i>Period circadian clock 3 (Per3)</i>	Fw	ACGAAGCTCCTCGAATCCCT
	Rv	GAATCTGACGGGCGAGTGTT
<i>Cryptochrome 1 (Cry1)</i>	Fw	CGGTAGAGGAAGTCGGGGTG
	Rv	AGACACTGAAGCAAAAATCGCC
<i>Cryptochrome 2 (Cry2)</i>	Fw	TGAAGAACTGACCAAGCGGG
	Rv	GTTGGTGATTGGCTTCTCTGC
<i>Hypoxanthine-guanine phosphoribosyltransferase (Hprt)</i>	Fw	GTTGGGCTTACCTCACTGCT
	Rv	TAATCACGACGCTGGGACTG