

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Urinary Phthalate Metabolites and Slow Walking Speed in the Korean Elderly Environmental Panel II Study

Jeonggyo Yoon, Esther García-Esquinas, Junghoon Kim, Jung Hyun Kwak, Hongsoo Kim, Sungroul Kim, Kyoung-Nam Kim, Yun-Chul Hong, and Yoon-Hyeong Choi

Table of Contents

Table S1. Timeline of KEEP II data collection.

Table S2. Distribution of urinary phthalate concentrations ($\mu\text{g/L}$) at enrollment and follow-up in KEEP II study participants.

Table S3. Distribution of creatinine-corrected urinary phthalate concentrations ($\mu\text{g/g cre}$) at enrollment and follow-up in KEEP II study participants.

Table S4. Geometric means (95% CIs) of urinary phthalate metabolite concentrations according to participant characteristics at enrollment in the KEEP II study (numeric values of Figure 2).

Table S5. Spearman's correlation coefficients between enrollment and follow-up visits for each phthalate metabolite in KEEP II study participants.

Table S6. Odds ratios (95% CI) for slowness (< 1.0 m/s) according to urinary phthalate metabolite concentrations (non-adjusted and covariate-adjusted) from cross-sectional analysis at enrollment ($n = 1,190$) and longitudinal analysis ($n = 2,218$).

Table S7. Change (95% CI) in walking speed (m/s) according to urinary phthalate metabolite concentrations (non-adjusted and covariate-adjusted) from cross-sectional analysis at enrollment ($n = 1,190$) and longitudinal analysis ($n = 2,218$).

Table S8. Odds ratios (95% CI) for slowness (< 0.8 m/s) according to urinary phthalate metabolite concentrations ($\mu\text{g/g cre}$) from cross-sectional analysis at enrollment ($n = 1,190$) and longitudinal analysis ($n = 2,218$).

Table S9. Odds ratios (95% CI) for sex-and height-adjusted slowness by urinary phthalate metabolite concentrations ($\mu\text{g/g cre}$) from cross-sectional analysis at enrollment ($n = 1190$) and longitudinal analysis ($n = 2218$).

Table S10. Odds ratios (95% CI) for slowness (< 1.0 meter/second) by urinary phthalate concentrations ($\mu\text{g/g cre}$) from longitudinal analysis after further adjustments for socioeconomic status ($n = 1,041$) and depression ($n = 1,189$).

Table S11. Odds ratios (95% CI) for slowness (< 1.0 meter/second) according to urinary phthalate concentrations ($\mu\text{g/g cre}$) in men and women, from longitudinal analysis ($n = 2,218$).

Table S12. Overall effects of phthalate metabolite mixtures on changes in walking speed (m/s, meter/second) estimated by Bayesian kernel machine regression (numeric values of Figure 3).

Table S13. Posterior inclusion probabilities for individual phthalate metabolites using Bayesian kernel machine regression analysis.

Table S14. Spearman's coefficients between MnBP and household income at enrollment in the KEEP II study.

Figure S1. Directed acyclic graph (DAG) for representing the potential confounding effect of socioeconomic status (SES) on associations between MnBP and slowness of walking speed in men (*A*) and women (*B*). The potential causal relationships (either protective or adverse) between one variable on another were represented as arrows.