

Supplementary Table 2. Description of primers used in qRT-PCR, Targeted gene deep sequencing, Sanger sequencing and Methylation-Specific PCR.

Primer sequences qRT-PCR

Target gene	Primer sequences (5'-3')	Size
E2F1 ¹	CATCCAGGAGGTCACTTCTG GACAACAGCGGTCTTGCTC	144
P57 ²	GGCCTGTGATCCGATTTCTTC GGGTCTGCTCCACCGAG	126
TAp53 ³	CGCAGTCAGATCCTAGCGTC CTGGACCTGGGTCTTCAGTG	171
Δ133p53 ⁴	GGTTGCAGGAGGTGCTTACAC GTTGAGGGCAGGGGAGTACTG	128
β2μG	CCAGCAGAGAATGGAAAGTC CGAGACATGTAAGCAGCATC	285
β actin	CAGGCACCAGGGCGTG GTGAGGATGCCTCTTGTCT	84

MystiCQ™ microRNA qPCR

Target	SIGMA Reference
Universal PCR Primer	MIRUP
Assay Primer hsa-miR-25-3p	MIRAP00059
Assay Primer hsa-miR-200a-3p	MIRAP00247
Assay Primer hsa-miR-203	MIRAP00255
Assay Primer hsa-miR-205-5p	MIRAP00257
Assay Primer hsa-miR-221-3p	MIRAP00277
Assay Primer hsa-miR-222-3p	MIRAP00279
Assay Primer hsa-miR-375	MIRAP00360
Control Primer SNORD48	MIRCP00007

Primer sequences for Sanger DNA sequencing

Target Gene	Region	Sequence 5'-3'	Size
TP53	Exon 4 ⁵	ACTGCTCTTTTCAACCATCTAC TCATGGAAGCCAGCCCTCAG	332
	Exon 5	GGGTAGACGCCAACTCTCTC CCAGCTGCTCACCATCGCTA	340
	Exon 6 ^{6,7}	TGGTTGCCAGGGTCCCC CGGAGGGCCACTGACAACC	223
	Exon 7 ⁸	TGCCACAGGTCTCCCAAGG AGTGTGCAGGGTGGCAAGTG	196
	Exon 8	GCTTTGAGGTGCGTGTGTTGT CTGAGGCATAACTGCACCT	170

* European Molecular Biology Laboratory (EMBL) Accession No. X54156.1 (p53 gene sequence).

Primer sequences for Methylation-Specific PCR and Bisulfite Genomic Sequencing

Target Gene	Sequence 5'-3'	Size	
P57	MSP-U-F	GTTGAATGTTGAGGATTAGAATTGT	203
	MSP-U-R	CAAAAAACAACTAACAACCACCAC	
	MSP-M-F	AGTTGAACGTCGAGGATTAGAATC	203
	MSP-M-R	GAAAAACGAATAACAACCACCG	
BS-F	GAGGATTAGAATYGTGGGATT	184	
	BS-R		AACTAACAACCACRCRA

References:

1. Wang P, Alvarez-Perez JC, Felsenfeld DP, et al. A high-throughput chemical screen reveals that harmine-mediated inhibition of DYRK1A increases human pancreatic beta cell replication. *Nat Med.* 2015;21(4):383-388.
2. Ueberberg S, Tannapfel A, Schenker P, et al. Differential expression of cell-cycle regulators in human beta-cells derived from insulinoma tissue. *Metabolism.* 2016;65(5):736-746.
3. Nutthasirikul N, Limpai boon T, Leelayuwat C, Patrakitkomjorn S, Jearanaikoon P. Ratio disruption of the 133p53 and TAp53 isoform equilibrium correlates with poor clinical outcome in intrahepatic cholangiocarcinoma. *Int J Oncol.* 2013;42(4):1181-1188.
4. Ohgaki H, Dessen P, Jourde B, et al. Genetic pathways to glioblastoma: a population-based study. *Cancer Res.* 2004;64(19):6892-6899.
5. Bharaj BS, Angelopoulou K, Diamandis EP. Rapid sequencing of the p53 gene with a new automated DNA sequencer. *Clin Chem.* 1998;44(7):1397-1403.
6. Serakinci N, Christensen R, Graakjaer J, et al. Ectopically hTERT expressing adult human mesenchymal stem cells are less radiosensitive than their telomerase negative counterpart. *Exp Cell Res.* 2007;313(5):1056-1067.
7. Millikan R, Hulka B, Thor A, et al. p53 mutations in benign breast tissue. *J Clin Oncol.* 1995;13(9):2293-2300.