

RESEARCH ARTICLE

Open Access

Genetic variation in the tau protein phosphatase-2A pathway is not associated with Alzheimer's disease risk

José L Vázquez-Higuera¹, Ignacio Mateo¹, Pascual Sánchez-Juan¹, Eloy Rodríguez-Rodríguez¹, Ana Pozueta¹, Miguel Calero², José L Dobato³, Ana Frank-García⁴, Fernando Valdivieso⁵, José Berciano¹, Maria J Bullido⁵ and Onofre Combarros^{1*}

Abstract

Background: Tau abnormal hyperphosphorylation and the formation of neurofibrillary tangles in AD brain is the result of upregulation of tau kinases and downregulation of tau phosphatases.

Methods: In a group of 729 Spanish late-onset Alzheimer's disease (AD) patients and 670 healthy controls, we examined variations into a set of candidate genes (PPP2CA, PPP2R2A, ANP32A, LCMT1, PPME1 and PIN1) in the tau protein phosphatase-2A (PP2A) pathway, to address hypotheses of genetic variation that might influence AD risk.

Results: There were no differences in the genotypic, allelic or haplotypic distributions between cases and controls in the overall analysis or after stratification by age, gender or APOE ϵ 4 allele.

Conclusion: Our negative findings in the Spanish population argue against the hypothesis that genetic variation in the tau protein phosphatase-2A (PP2A) pathway is causally related to AD risk

Background

One of the neuropathological hallmarks in Alzheimer's disease (AD) is neurofibrillary tangles (NFTs), which are composed of the microtubule-binding protein tau that is hyperphosphorylated [1]. Tau phosphorylation is catalysed by tau protein kinases and reversed by tau protein phosphatases [2]. It has been reported that the expression and activity of the major tau phosphatase in human brain, protein phosphatase-2A (PP2A), is decreased in the affected areas of AD brain [3,4], suggesting that a downregulation of tau phosphatases in AD brain might underlie the abnormal hyperphosphorylation of tau. The overall PP2A activity is determined by composition of the holoenzyme from the catalytic subunit alpha (PP2CA) and the regulatory subunit B alpha (PP2R2A), the level of PP2A inhibitors such as ANP32A (inhibitor-1 of protein phosphatase-2A), and changes in PP2A methylation regulated by the leucine carboxyl methyltransferase-1 (LCMT1) and the protein


phosphatase methylesterase-1 (PPME1); in addition, the peptidyl-prolyl cis/trans isomerase PIN1 induces conformational changes in tau that can facilitate tau dephosphorylation by PP2A (Figure 1). Consequently, PPP2CA, PPP2R2A, ANP32A, LCMT1, PPME1 and PIN1 are good candidate genes for the analysis of AD susceptibility. The largest genome wide association (GWA) study in AD [5] did not find significant results for all these PP2A-related genes. However, it cannot be discarded that these genes in the tau PP2A pathway are among the genes with significant nominal association but without reaching significance ($p < 10^{-5}$ or less) after adjustment for multiple testing in GWAs; in addition, it is also possible that some of the SNPs analyzed in our study were not present in the arrays used in GWAs. Therefore, we conducted a case-control association study involving genes in the tau PP2A pathway in relation to AD risk, in a Spanish cohort.

Methods

The study included 729 AD patients (67% women; mean age at study 77.2 years; SD 8.0; range 61-103 years; mean age at onset 73.3 years; SD 7.8; range 60-100

* Correspondence: combarro@unican.es

¹Neurology Service and CIBERNED, "Marqués de Valdecilla" University Hospital (University of Cantabria and IFIMAV), Santander, Spain
Full list of author information is available at the end of the article



years) who met NINCDS/ADRDA criteria for probable AD [6]. All AD cases were defined as sporadic because their family history did not mention any first-degree relative with dementia. AD patients were recruited from the Departments of Neurology of University Hospital “Marqués de Valdecilla” (Santander, Spain) and Hospital “La Paz” (Madrid, Spain), and from Alzheimer Center Reina Sofia Foundation (Madrid, Spain). The large majority of patients were living in the community and had been referred by their general practitioner; few had been admitted from hospital wards or nursing home facilities. Control subjects were 670 unrelated individuals (64% women; mean age 78.3 years; SD 9.4; range 60-104 years) randomly selected from nursing homes. These subjects had complete neurologic and medical examinations that showed that they were free of significant illness and had Mini Mental State Examination scores of 28 or more, which were verified by at least one subsequent annual following-up assessment. The controls arose from the same base population as the cases. The AD and control samples were Caucasians originating from a limited geographical area in northern Spain (Santander) and from the central area of Spain (Madrid).

Blood samples were taken after written informed consent had been obtained from the subjects or their representatives. The study was approved by the ethical committees of the University Hospital “Marqués de Valdecilla”, Alzheimer Center Reina Sofia Foundation, and the Hospital “La Paz”. Genotyping of PPP2CA (rs7705319 and rs4958177), PPP2R2A (rs2046225, rs7823021, rs17055172, rs12676426 and rs2046223),

ANP32A (rs2924633, rs1551345, rs1551344, rs1551342, rs11632936, rs2958405 and rs1551343), LCMT1 (rs8062337, rs277898, rs277886 and rs277892), PPME1 (rs10898966, rs2848557, rs500608 and rs544356), and PIN1 (rs2233678, rs1077220, rs2010457 and rs2287838) polymorphisms was performed using the iPLEX Gold assay on the MassArray system (Sequenom Inc., San Diego, USA). We used data from the HapMap project <http://www.hapmap.org> to select htSNPs capturing 100% of PPP2CA genetic variability, 75% of PPP2R2A, 95% of ANP32A, 95% of LCMT1, 75% of PPME1, and 80% of PIN1 genetic variability in Caucasians. SNPs were chosen among those with minor allele frequencies $\geq 5\%$ using Haploview v3.2 software <http://www.broad.mit.edu/mpg/haploview> with an r^2 threshold of 0.8. PPP2R2A rs12676426 and PPME1 rs500608 that were significantly deviated from Hardy-Weinberg equilibrium (HWE) were excluded from the analysis. In addition, we removed PIN1 rs1077220 with a non-homogeneous genotypic distribution between our two control populations (Santander and Madrid) and which were also different from the HapMap CEU distribution.

HWE was calculated for the htSNPs in the control population using Pearson’s χ^2 statistics. We assessed pairwise linkage disequilibrium (LD) between the htSNPs by D' and r^2 statistics. Haplotype reconstruction and their frequencies in cases and controls were estimated by an expectation-maximization algorithm, method implemented in Haploview 3.32. Pearson’s χ^2 statistics were performed to compare genotype, allele, and haplotype distribution of the patients and control for each htSNP. Genotypic and allelic distributions were assessed by logistic regression using SPSS software.

Results

As shown in Table 1, the distribution of the minor allele frequencies of the PP2A-related genes did not differ significantly between AD and control groups. Haplotype distributions were not significantly different between cases and controls in the overall analysis or after stratification by APOE $\epsilon 4$ allele (data not shown). There were no major differences in allele, genotype or haplotype frequencies in our total sample associated to either age or gender subgroups. The data set was analyzed for epistatic interactions between the 6 PP2A-related genes and no significant effects were observed.

Discussion

Genes harbouring markers with only modest evidence of association (nominally significant but not reaching the genome-wide significance threshold) can be identified if they belong to the same biological pathway or mechanism; therefore, pathway-based approaches,

Table 1 Minor allele frequencies distribution of tau phosphatases genes in AD patients and controls

Gene	SNP	MAF, AD/C	P-value	Gene	SNP	MAF, AD/C	P-value
PPP2CA	rs7705319	0.20/0.20	0.99	LCMT1	rs8062337	0.32/0.30	0.34
	rs4958177	0.11/0.10	0.46		rs277898	0.37/0.38	0.35
					rs277886	0.34/0.32	0.16
PPP2R2A	rs2046225	0.38/0.37	0.49		rs277892	0.28/0.27	0.40
	rs7823021	0.28/0.29	0.88	PPME1	rs10898966	0.11/0.09	0.24
	rs17055172	0.15/0.17	0.31		rs2848557	0.34/0.34	0.84
	rs2046223	0.38/0.39	0.82		rs544356	0.20/0.20	0.89
ANP32A	rs2924633	0.40/0.40	0.67	PIN1	rs2233678	0.11/0.12	0.47
	rs1551345	0.09/0.08	0.48		rs2010457	0.33/0.32	0.83
	rs1551344	0.07/0.06	0.31		rs2287838	0.48/0.49	0.84
	rs1551342	0.11/0.12	0.57				
	rs11632936	0.27/0.27	0.97				
	rs2958405	0.31/0.34	0.07				
	rs1551343	0.13/0.11	0.20				

P-values not corrected for multiple comparisons.

which jointly consider multiple variants in interacting or related genes, might complement the most-significant SNPs/genes approach for interpreting genome-wide association (GWA) data on complex diseases [7,8]. In fact, genetic variation in the immune system and in lipid metabolism pathways is a cause of AD susceptibility [9,10]. Although genetic markers of the genes examined in this study (PPP2CA, PPP2R2A, ANP32A, LCMT1, PPME1 and PIN1) were not found associated to AD in the largest GWA study [5], our main hypothesis was that common variation in genes directly related to tau dephosphorylation in the PP2A pathway might underlie individual differences in susceptibility to AD. To our knowledge, this study is the first of its type to be conducted in this pathway. There is a selective and significant neuron-specific reduction in PP2A catalytic subunit and PP2A regulatory B subunit mRNAs in AD hippocampus [3,4], and this reduced neuronal PP2A immunoreactivity closely correlates with NFT load [4], suggesting that PP2A dysfunction contributes to AD tau pathology. In addition, cDNA microarray techniques have revealed downregulated expression of PPP2CA gene in AD brain [11,12]. Conversely, in AD brain has been observed a significant increase in the neocortical levels of PP2A inhibitors [13], which co-localize with abnormally hyperphosphorylated tau. These data suggest the possible involvement of ANP32A (inhibitor-1 of PP2A) in AD neurofibrillary pathology through the inhibition of PP2A activity. LCMT1 promotes PP2A activity by methylating its catalytic subunit, and PPME1 catalyses the removal of the methyl group, thus reversing the activity of LCMT1. Methylated PP2A catalytic subunit levels are reduced in AD frontal/temporal cortex [14],

and LCMT1 levels are also selectively decreased in AD-affected regions and in tangle-bearing neurons [15]. An interaction between tau and PIN1 is thought to facilitate the dephosphorylation of hyperphosphorylated tau by PP2A, promoting microtubule stability [16]. In AD brain, PIN1 expression in the hippocampus and parietal cortex is low [17] and inversely correlated with neurofibrillary degeneration [18].

In an Italian study [19], carriers of PIN1 (-842, rs2233678) C allele had an increased risk of AD, lower age of onset, and reduced PIN1 levels in peripheral mononuclear cells, but subsequent studies [20-23] did not replicate these findings. We failed to detect the association of PIN1 (-842, rs2233678) with AD. We also failed to observe any allele, genotype or haplotype association of PPP2CA, PPP2R2A, ANP32A, LCMT1 and PPME1 genes with AD. Because we studied htSNPs capturing 80% of PIN1 and 75% of PPP2R2A and PPME1 genetic variability, it might be argued that we have missed a hypothetical disease locus, which would have been detected by analysis of extended haplotypes; however, the complete linkage disequilibrium across the PIN1, PPP2R2A and PPME1 regions in our study argues against this possibility. Our negative results with all these PP2A pathway-related genes are probably not due to insufficient statistical power, because our sample size had enough power (94%) to detect and odds ratio of 1.5 at disease allele frequencies of 0.10.

Conclusion

Despite supporting evidence for the biological role of tau phosphatases in AD exists, our negative findings in the Spanish population argue against the hypothesis that

genetic variation in the tau protein phosphatase-2A (PP2A) pathway is causally related to AD risk.

Acknowledgements

C. Sánchez-Quintana was involved in the DNA sample collections from Santander. We also thank Drs. P. Gil and P. Coria for their cooperation in the generation of the case-control samples from Madrid. This work was made possible by the generous participation of the patients, the control subjects, and their families. This study was supported by grants from CIEN Foundation (PI019/09) and CIBERNED (CB06/07/0037).

Author details

¹Neurology Service and CIBERNED, "Marqués de Valdecilla" University Hospital (University of Cantabria and IFIMAV), Santander, Spain. ²Spongiform Encephalopathies Unit, National Microbiology Centre and CIBERNED, Carlos III Health Institute, Madrid, Spain. ³Alzheimer Disease Research Unit, CIEN Foundation, Carlos III Health Institute, Alzheimer Center Reina Sofia Foundation, Madrid, Spain. ⁴Neurology Service and CIBERNED, Hospital Universitario La Paz (U.A.M.), Madrid, Spain. ⁵Molecular Biology Department and CIBERNED, Centro de Biología Molecular Severo Ochoa (C.S.I.C.-U.A.M.), Madrid, Spain.

Authors' contributions

JLVH and ERR performed the genetic studies and reviewed critically the manuscript. PSJ performed the statistical analyses and reviewed critically the manuscript. IM, AP, MC, JLD, AF, FV, JB and MJB reviewed critically the manuscript. OC drafted the manuscript and contributed to its final version. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 1 June 2011 Accepted: 7 September 2011

Published: 7 September 2011

References

- Ballatore C, Lee VMY, Trojanowski JQ: **Tau-mediated neurodegeneration in Alzheimer's disease and related disorders.** *Nat Rev Neurosci* 2007, **8**:663-72.
- Liu F, Grundke-Iqbal I, Iqbal K, Gong CX: **Contributions of protein phosphatases PP1, PP2A, PP2B and PP5 to the regulation of tau phosphorylation.** *Eur J Neurosci* 2005, **22**:1942-50.
- Vogelsberg-Ragaglia V, Schuck T, Trojanowski JQ, Lee VMY: **PP2A mRNA expression is quantitatively decreased in Alzheimer's disease hippocampus.** *Exp Neurol* 2001, **168**:402-12.
- Sontag E, Luangpirom A, Hladik C, Mudrak I, Ogris E, Speciale S, White CL: **Altered expression levels of the protein phosphatase 2A ABαC enzyme are associated with Alzheimer disease pathology.** *J Neuropathol Exp Neurol* 2004, **63**:287-301.
- Hollingsworth P, Harold D, Sims R, Gerrish A, Lambert JC, Carrasquillo MM, Abraham R, Hamshere ML, Pahwa JS, Moskva V, Dowzell K, Jones N, Stretton A, Thomas C, Richards A, Ivanov D, Widdowson C, Chapman J, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Beaumont H, Warden D, Wilcock G, Love S, Kehoe PG, Hooper NM, Vardy ER, Hardy J, Mead S, Fox NC, Rossor M, Collinge J, Maier W, Jessen F, Ruther E, Schürmann B, Heun R, Kölsch H, van den Bussche H, Heuser I, Kornhuber J, Wiltfang J, Dichgans M, Frölich L, Hampel H, Gallacher J, Hüll M, Rujescu D, Giegling I, Goate AM, Kauwe JS, Cruchaga C, Nowotny P, Morris JC, Mayo K, Sleegers K, Bettens K, Engelborghs S, De Deyn PP, Van Broeckhoven C, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Tsolaki M, Singleton AB, Guerreiro R, Mühleisen TW, Nöthen MM, Moebus S, Jöckel KH, Klopp N, Wichmann HE, Pankratz VS, Sando SB, Aasly JO, Barcikowska M, Wszolek ZK, Dickson DW, Graff-Radford NR, Petersen RC, the Alzheimer's Disease Neuroimaging Initiative, van Duijn CM, Breterler MM, Ikram MA, Destefano AL, Fitzpatrick AL, Lopez O, Launer LJ, Seshadri S, CHARGE consortium, Berr C, Campion D, Epelbaum J, Dartigues JF, Tzourio C, Alperovitch A, Lathrop M, EADI1 consortium, Feulner TM, Friedrich P, Riehle C, Krawczak M, Schreiber S, Mayhaus M, Nicolhaus S, Wagenpfeil S, Steinberg S, Stefansson H, Stefansson K, Snædal J, Björnsson S, Jonsson PV, Chouraki V, Genier-Boley B, Hiltunen M, Soininen H, Combarros O, Zelenika D, Delepine M, Bullido MJ, Pasquier F, Mateo I, Frank-Garcia A, Porcellini E, Hanon O, Coto E, Alvarez V, Bosco P, Siciliano G, Mancuso M, Panza F, Solfrizzi V, Nacmias B, Sorbi S, Bossù P, Piccardi P, Arosio B, Annoni G, Seripa D, Pilotto A, Scarpini E, Galimberti D, Brice A, Hannequin D, Licastro F, Jones L, Holmans PA, Jonsson T, Riemenschneider M, Morgan K, Younkin SG, Owen MJ, O'Donovan M, Amouyel P, Williams J: **Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease.** *Nat Genet* 2011, **43**:429-35.
- McKhaan G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM: **Clinical diagnosis of Alzheimer's disease: report of the NINCDA-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease.** *Neurology* 1984, **34**:934-44.
- Wang K, Li M, Bucan M: **Pathway-based approaches for analysis of genome-wide association studies.** *Am J Hum Genet* 2007, **81**:1278-83.
- Zhong H, Yang X, Kaplan LM, Molony C, Schadt EE: **Integrating pathway analysis and genetics of gene expression for genome-wide association studies.** *Am J Hum Genet* 2010, **86**:581-91.
- Jones L, Holmans PA, Hamshere ML, Harold D, Moskva V, Ivanov D, Pocklington A, Abraham R, Hollingsworth P, Sims R, Gerrish A, Pahwa JS, Jones N, Stretton A, Morgan AR, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Morgan K, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Love S, Kehoe PG, Mead S, Fox N, Rossor M, Collinge J, Maier W, Jessen F, Schürmann B, van den Bussche H, Heuser I, Peters O, Kornhuber J, Wiltfang J, Dichgans M, Frölich L, Hampel H, Hüll M, Rujescu D, Goate AM, Kauwe JS, Cruchaga C, Nowotny P, Morris JC, Mayo K, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Singleton AB, Guerreiro R, Mühleisen TW, Nöthen MM, Moebus S, Jöckel KH, Klopp N, Wichmann HE, Ruther E, Carrasquillo MM, Pankratz VS, Younkin SG, Hardy J, O'Donovan MC, Owen MJ, Williams J: **Genetic evidence implicates the immune system and cholesterol metabolism in the aetiology of Alzheimer's disease.** *PLoS ONE* 2010, **5**:e13950.
- Lambert JC, Grenier-Boley B, Chouraki V, Heath S, Zelenika D, Fievet N, Hannequin D, Pasquier F, Hanon O, Brice A, Epelbaum J, Berr C, Dartigues JF, Tzourio C, Campion D, Lathrop M, Amouyel P: **Implication of the immune system in Alzheimer's disease: evidence from genome-wide pathway analysis.** *J Alzheimers Dis* 2010, **20**:1107-18.
- Ginsberg SD, Hemby SE, Lee VMY, Eberwine JH, Trojanowski JQ: **Expresión profile of transcripts in Alzheimer's disease tangle-bearing CA1 neurons.** *Ann Neurol* 2000, **48**:77-87.
- Tan MG, Chua WT, Esiri MM, Smith AD, Vinters HV, Lai MK: **Genome wide profiling of altered gene expression in the neocortex of Alzheimer's disease.** *J Neurosci Res* 2010, **88**:1157-69.
- Tanimukai H, Grundke-Iqbal I, Iqbal K: **Up-regulation of inhibitors of protein phosphatase-2A in Alzheimer's disease.** *Am J Pathol* 2005, **166**:1761-71.
- Zhou XW, Gustafsson JA, Tanila H, Bjorkdahl C, Liu R, Winblad B, Pei JJ: **Tau hyperphosphorylation correlates with reduced methylation of protein phosphatase 2A.** *Neurobiol Dis* 2008, **31**:386-94.
- Sontag E, Hladik C, Montgomery L, Luangpirom A, Mudrak I, Ogris E, White CL: **Downregulation of protein phosphatase 2A carboxyl methylation and methyltransferase may contribute to Alzheimer disease pathogenesis.** *J Neuropathol Exp Neurol* 2004, **63**:1080-91.
- Lu PJ, Wulf G, Zhou XZ, Davies P, Lu KP: **The prolyl isomerase Pin1 restores the function of Alzheimer-associated phosphorylated tau protein.** *Nature* 1999, **399**:784-788.
- Sultana R, Boyd-Kimball D, Poon HF, Cai J, Pierce WM, Klein JB, Markesbery WR, Zhou XZ, Lu KP, Butterfield DA: **Oxidative modification and down-regulation of Pin1 in Alzheimer's disease hippocampus: a redox proteomics analysis.** *Neurobiol Aging* 2006, **27**:918-25.
- Liou YC, Sun A, Ryo A, Zhou XZ, Yu ZX, Huang HK, Uchida T, Bronson R, Bing G, Li X, Hunter T, Lu KP: **Role of the prolyl isomerase Pin1 in protecting against age-dependent neurodegeneration.** *Nature* 2003, **424**:556-61.
- Segat L, Pontillo A, Annoni G, Trabattoni D, Vergani C, Clerici M, Arosio B, Crovella S: **PIN1 promoter polymorphisms are associated with Alzheimer's disease.** *Neurobiol Aging* 2007, **28**:69-74.

20. Lambert JC, Beniseain F, Chapuis J, Cotel D, Amouyel P: **Association study of the PIN1 gene with Alzheimer's disease.** *Neurosci Lett* 2006, **402**:259-61.
21. Nowotny P, Bertelsen S, Hinrichs AL, Kauwe JSK, Mayo K, Jacquart S, Morris JC, Goate A: **Association studies between common variants in Prolyl isomerase Pin1 and the risk for late-onset Alzheimer's disease.** *Neurosci Lett* 2007, **419**:15-7.
22. Maruszak A, Safranow K, Gustaw K, Kijanowska-Haladyna B, Jakubowska K, Olszewska M, Styczynska M, Berdyski M, Tysarowski S, Chlubek D, Siedlecki J, Barcikowska M, Zekanowski C: **PIN1 gene variants in Alzheimer's disease.** *BMC Med Genet* 2009, **10**:115.
23. Ma SL, Tang NL, Tam CW, Cheng Lui VW, Lam LC, Chiu HF, Driver JA, Pastorino L, Lu KP: **A PIN1 polymorphism that prevents its suppression by AP4 associates with delayed onset of Alzheimer's disease.** *Neurobiol Aging* 2010.

doi:10.1186/1756-0500-4-327

Cite this article as: Vázquez-Higuera *et al.*: Genetic variation in the tau protein phosphatase-2A pathway is not associated with Alzheimer's disease risk. *BMC Research Notes* 2011 **4**:327.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

