

PRIMER

Model systems for regeneration: zebrafish

Ines J. Marques¹, Eleonora Lupi^{1,2} and Nadia Mercader^{1,3,*}

ABSTRACT

Tissue damage can resolve completely through healing and regeneration, or can produce permanent scarring and loss of function. The response to tissue damage varies across tissues and between species. Determining the natural mechanisms behind regeneration in model organisms that regenerate well can help us develop strategies for tissue recovery in species with poor regenerative capacity (such as humans). The zebrafish (*Danio rerio*) is one of the most accessible vertebrate models to study regeneration. In this Primer, we highlight the tools available to study regeneration in the zebrafish, provide an overview of the mechanisms underlying regeneration in this system and discuss future perspectives for the field.

KEY WORDS: Regeneration, Zebrafish, Cell progenitors, Injury methods, Genetic tools

Introduction

Most vertebrate tissues undergo continuous cell turnover to maintain tissue homeostasis, and cells that no longer exert their function correctly are eliminated and replaced by new ones. This continuous cellular turnover is not homogenous in all tissues: whereas gut enterocytes and skin keratinocytes show a high turnover (Young and Heath, 2000), cardiomyocytes (Bergmann et al., 2009; Senyo et al., 2013) and neurons are rarely replaced throughout adult life. In the case of neurons, proliferation is limited to specific regions in the brain (Ernst et al., 2014; Lim and Alvarez-Buylla, 2016). Beyond homeostatic tissue maintenance, some tissues and organs are capable of reacting to an injury by replacing the lesioned region. In this scenario, responses can either involve tissue repair, where a fibrotic scar replaces the injury, or regeneration, where new cells proliferate and/or differentiate into the same cell type as the one that was damaged, thus reconstituting the tissue so that it can recover its functional activity. As a general outline, during injury response the damaged area (Fig. 1A,B) is initially infiltrated by immune cells (Fig. 1C), which clean the dead tissue and prevent infections. This is followed by the deposition of extracellular matrix (ECM) (Fig. 1D), which provides support and structure to the injured area. ECM deposition results in the development of a, sometimes permanent, fibrotic scar. Alternatively, the lesioned region can be replaced by new tissue (Fig. 1E) in a multi-step regenerative process that can involve dedifferentiation, proliferation and re-differentiation

of existing cells (as discussed further below). In some cases, the regenerative phase occurs concomitant with the regression of the transient fibrotic tissue. Overall, the resulting regenerated tissue has the same morphological and physiological properties as the lost tissue, and is hence fully functional (Fig. 1F) (Hardy, 1989).

Vertebrates vary widely in their regenerative capacity. Whereas mammals usually have a more limited regenerative capacity – restricted to few tissues or organs (e.g. skin and liver) or a short time window after birth [e.g. the heart (Porrello et al., 2011)] – other vertebrates possess a higher degree of regenerative capacity: e.g. axolotls are capable of regenerating several organs, including the heart (Cano-Martinez et al., 2010) and the limbs (Simon and Tanaka, 2013). Among the vertebrate animal models available for regeneration studies, the zebrafish (*Danio rerio*) – a 2-5 cm long freshwater teleost – is one of the most widely used. Following their establishment as a genetically and experimentally accessible model system for studying development (Haffter et al., 1996; Streisinger et al., 1981), early regeneration studies showed that zebrafish, like many other fish, are able to fully regenerate many tissues and organs (Bernhardt et al., 1996; Johnson and Weston, 1995; Poss et al., 2002; Raya et al., 2003; Rowleson et al., 1997; White et al., 1994). This pioneering research convinced the scientific community to consider this model organism for studying organ regeneration as well as development. Some of the tissues and organs most typically used in regeneration studies include the spinal cord (Becker et al., 1998), brain and cerebellum (Kroehne et al., 2011; Liu et al., 2004), retina (Vihtelic and Hyde, 2000), hair cells (Harris et al., 2003), heart (Poss et al., 2002; Raya et al., 2003), caudal fin (Nechiporuk and Keating, 2002; White et al., 1994), kidney (Reimschuessel, 2001), and liver (Burkhardt-Holm et al., 1999).

Regeneration studies have been greatly facilitated by the ever-increasing power of genetic tools, including transgenesis and genome editing to generate loss-of-function mutations. (Hwang et al., 2013; Kawakami, 2007; Moore et al., 2012; Stuart et al., 1988). In comparison with other highly regenerative animal models such as the axolotl, these were developed much earlier in zebrafish,

Model systems for regeneration

This article is part of a series entitled 'Model systems for regeneration'. This series of articles aims to highlight key model systems and species that are currently being used to study tissue and organ regeneration. Each article provides background information about the phylogenetic position of the species, its life-cycle and habitat, the different organs and tissues that regenerate, and the experimental tools and techniques that are available for studying these organisms in a regenerative context. Importantly, these articles also give examples of how the study of these models has increased our understanding of regenerative mechanisms more broadly, and how some of the open questions in the field of regeneration may be answered using these organisms. To see the full collection as it grows, please visit: https://dev.biologists.org/collection/regeneration_models

¹Institute of Anatomy, University of Bern, Bern 3012, Switzerland. ²Acquifer, Databases, Digital Biomedical Imaging Systems, Pforzheim, Germany. ³Centro Nacional de Investigaciones Cardiovasculares CNIC, Madrid 2029, Spain.

*Author for correspondence (nadia.mercader@ana.unibe.ch)

 I.J.M., 0000-0003-1254-9394; E.L., 0000-0003-0509-3616; N.M., 0000-0002-0905-6399

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.




Fig. 1. Phases of tissue regeneration. (A) Uninjured tissue. (B) Injury: injured area is shaded in gray, representing damaged and dead cells. The blood vessel in direct contact with the lesion is also damaged and is marked in a lighter red color than the healthy vessels. (C) Wound closure and immune response: the injured area is infiltrated by platelets to stop the bleeding resulting from injury to the blood vessels in the damaged area. At the same time, the lesion is also infiltrated by immune cells. (D) Clearing of debris and deposition of extracellular matrix (ECM): the immune cells start to clean the dead tissue and other debris resulting from the injury. Fibroblasts proliferate and produce ECM fibers that fill the cleared injury area to prevent the injury from collapsing. The formation of new blood vessels takes place (thinner red lines). (E) Resorption of ECM and cell proliferation: the ECM starts to be eliminated as the injury is repopulated by new cells through proliferation. (F) Regeneration: the injured area has been completely repopulated and the tissue reconstituted to its homeostatic condition. Cells undergo cell cycle arrest and very few proliferating cells can be seen.

further stimulating its use for regeneration studies. Moreover, excellent platforms for sharing resources and knowledge (summarized in Table 1) are contributing to a steady growth of the research community working with this model organism, also in the context of regeneration. In this Primer, we provide a broad overview of the available tools for regeneration research in the zebrafish, and focus on the challenges and opportunities presented by this model organism.

Tools for studying regeneration in zebrafish

Many of the attributes that make zebrafish an appealing model for developmental biologists also give it significant advantages for regeneration researchers – with the genetic and genomic tools, and accessibility for live imaging and drug screening being particularly important. We discuss these topics further below, but focus first on the range of injury models available to researchers.

Tissue injury models

Several injury models have been established to study regeneration in zebrafish, including physical injury models such as resection,

stabbing or cryoinjury, or, alternatively, genetic procedures to ablate a specific cell type (Table 2 and Fig. 2). Although most studies are performed in adult zebrafish, in some instances larvae are also used to study organ regeneration. Choosing the appropriate injury model, and developmental stage, is an important decision. First, one should take into account the type of injury intended to be mimicked – particularly if the primary aim is to provide a model for a human condition. However, it is not always possible to mimic a mammalian injury process. For example, myocardial infarction is caused by an ischemic event, often due to the occlusion of coronary vessels by an atherosclerotic plaque. This type of injury can be mimicked in mice, but is very challenging to recapitulate in the zebrafish, owing to its small size. Second, the choice of the injury method also depends on the biological question to be answered. To study global processes of regeneration, one probably needs to use a method that lesions the whole tissue, such as surgically induced injuries. If the aim is to understand the regenerative capacity of a specific cell type, then genetic ablation might be a better choice. As a further consideration, different types of injury can lead to changes in the efficiency of cell replacement or speed of regeneration, as

Table 1. Online resources for the zebrafish community

Resource	Description	URL and references
Zebrafish Information Network (ZFIN)	Database of multiple resources available for zebrafish studies, including protocols and available lines	http://www.zfin.org (Howe et al., 2013)
International Zebrafish Society (IZFS)	Promotion of zebrafish research, organization of conferences, travel fellowships and awards	https://www.izfs.org/
EUFishBiomed	Promotes exchange of information within the fish community, support of conferences and travel fellowships	https://www.eufishbiomed.kit.edu/ (Strahle et al., 2012)
Zebrafish International Resource Center (ZIRC)	Zebrafish stock center (USA) collecting existing zebrafish resources	https://zebrafish.org/
Karlsruhe Institute of Technology (KIT)	Zebrafish stock center (Europe) collecting existing zebrafish resources and providing training	https://www.ezrc.kit.edu/ (Geisler et al., 2016)
China Zebrafish Resource Center (CZRC)	China Zebrafish Resource Center collecting existing zebrafish resources, developing new lines and technology	http://en.zfish.cn/

Table 2. Injury models for zebrafish organ regeneration

Organ	Injury method	References
Fin	Resection	Geraudie et al. (1993); Pfefferli and Jaźwińska (2015)
	Cryoinjury	Chassot et al. (2016)
Heart	Resection	Poss et al. (2002)
	Cryoinjury	Chablais et al. (2011); Gonzalez-Rosa et al. (2011); Schnabel et al. (2011)
	Genetic ablation	Wang et al. (2011)
Central nervous system (brain, spinal cord and eye)	Nerve crushing	Becker and Becker (2008); Bernhardt et al. (1996)
	Stabbing	März et al. (2011)
	Transection	Becker and Becker (2008); Becker et al. (1998)
	Visible light	Vihtelic and Hyde (2000)
	Heat	Raymond et al. (2006)
	Chemical	Fimbel et al. (2007)
Pancreas	Genetic ablation	Pisharath et al. (2007)
Liver	Resection	Sadler et al. (2007)
	Chemical	Cox et al. (2014)
Kidney	Chemical	Reimschuessel (2001)
Sensory hair cells	Chemical	Harris et al. (2003)
Skin	Laser	Richardson et al. (2013)

exemplified for the spinal cord (Ohnmacht et al., 2016) and the heart (Gonzalez-Rosa et al., 2011).

Surgical injuries

One of the most common surgical methods is resection (amputation) of the tissue. This technique is commonly used for fin and heart. Following fin amputation, the first step towards regeneration involves the development of an epidermal cap that closes the injury. Underneath, a blastema comprising undifferentiated cells is formed (discussed further below). After initial proliferation, these cells redifferentiate, culminating in the formation of new fin tissue (reviewed by Pfefferli and Jaźwińska, 2015). In the heart, the first step towards regeneration after ventricular resection is the formation of a blood clot, later replaced by a fibrin clot, followed by *de novo* growth to replace the lost tissue, until the damaged myocardium becomes almost indistinguishable from the surrounding tissue (reviewed by Gonzalez-Rosa et al., 2017).

Another popular surgically induced lesion method is cryoinjury, which is frequently used for the heart (Chablais et al., 2011; Gonzalez-Rosa et al., 2011; Schnabel et al., 2011) and has also been described in fin (Chassot et al., 2016). In this method, the tissue is exposed to a super-cooled metal, which effectively kills the cells it touches. In the fin, after the dead tissue falls off, regeneration

proceeds as in the amputation model (Chassot et al., 2016). By contrast, the regenerative process in the heart is a little different: it begins with the replacement of the necrotic tissue with ECM-depositing fibroblasts. Subsequently, the transient scar tissue regresses and is replaced by new cardiac tissue (Gonzalez-Rosa et al., 2012). As well as low temperatures, cell death can be induced by exposure to high temperatures, either using a heated metal rod (Raymond et al., 2006) or by applying a light source in a determined region (Conedera et al., 2017; Vihtelic and Hyde, 2000).

Finally, another method is stabbing, commonly used to damage the central nervous system, including the spinal cord (Becker et al., 1998). In this case, accumulation and proliferation of glial cells in the damaged area is followed by the proliferation of progenitors that ultimately differentiate into new nervous tissue (März et al., 2011).

Chemically induced cell damage

Besides physical stimuli, organ injuries can be induced by chemical compounds, which can either be directly administered to each individual fish by intraperitoneal injection, perorally or by dissolving in the fish water followed by ingestion. The toxicity of gentamicin is used to produce kidney or lateral line injuries (Kamei et al., 2015) and acetaminophen is used to induce liver injuries (Cox et al., 2014). After an initial phase of cell death, proliferation and cell migration lead to the repopulation of the damaged area, as has been demonstrated in studies on zebrafish kidney (Reimschuessel and Williams, 1995), lateral line hair cell (Harris et al., 2003) and liver (Burkhardt-Holm et al., 1999) regeneration. This method, however, presents some disadvantages, such as off-target effects, as well as a less uniform injuries caused by unequal drug distribution (Strahle and Grabher, 2010).

Genetic ablation

This approach allows tissue- or cell type-specific death and can also provide temporal control of cell ablation. In several transgenic zebrafish lines, cell death can be specifically induced in targeted cells. Methods include the regulated expression of diphtheria toxin A chain (Wang et al., 2011) and overexpression of a cytotoxic fluorescent protein, KillerRed – which has been used to ablate specific cell types through phototoxicity upon intense light illumination (Del Bene et al., 2010). Perhaps the most widely used system is based on the ability of the




Fig. 2. Organ regeneration in the zebrafish. Summary of some of the organs and tissues used for regeneration studies in zebrafish. Preferred injury models are annotated for each organ.

Table 3. Genetic tools for zebrafish regeneration studies

Method	Application	Pros	Cons
Fluorescent reporter lines	Cell/tissue imaging	Allow evaluation of gene expression pattern by promoter/enhancer-driven fluorescent reporter expression	Detection limited to promoter activity, not suited for fate mapping Regulatory region might not fully recapitulate endogenous gene expression pattern Fluorescent protein expression might be more stable than the endogenous gene
Photoconvertible proteins (e.g. KAEDE)	Lineage tracing	Activated by light Produces a faster response than chemically induced systems	Difficult to apply in non-superficial adult tissues Transient labelling Leakiness
CreER ^{T2} /loxP	Lineage tracing Gene overexpression	Temporal and spatial conditional gene expression Often used for lineage tracing studies	Cannot be turned off Requires chemical treatment Can be leaky (Cre activity even without tamoxifen induction)
Dre/rox	Lineage tracing Gene overexpression	Conditional gene expression If combined with CreER ^{T2} /loxP, it allows binary recombination studies Dre can be linked to estrogen or progesterone receptor to make it inducible (as with Cre)	Cannot be turned off Not well established in the zebrafish Fewer lines available than for Cre/Lox
TetOn/TetOff	Gene overexpression	Specific gene visualization and/or overexpression in a spatio-temporally controlled manner System can be turned on and off as needed	Few available lines Doxycyclin toxicity not well assessed
Heat-shock promoter	Gene overexpression	Temperature-dependent inducible expression Does not require chemical treatment	Gene expression levels are temperature dependent: precise temperature settings are critical for the method to work Promotor not equally efficient in all cell types
Gal4/UAS	Gene overexpression Cell/tissue imaging	Spatially controlled gene expression	Transcriptional silencing when Gal4 is expressed from very active promoters: mostly limits its use to developmental stages
TALEN	Genome editing and generation of mutant lines	Highly specific and efficient More versatile for sequence recognition than CRISPR	Time consuming Has not been applied to knock-in approaches
CRISPR/Cas9	Genome editing and generation of mutant lines	sgRNA easier to produce than TALEN Targeting of various genomic regions with one injection	Off-target effect possible Use for site-directed insertions still not fully established

bacterial enzyme nitroreductase to convert the prodrug metronidazol into a cytotoxin, leading to the death of nitroreductase-expressing cells. This technique was first described in studies on pancreatic β -cell regeneration in zebrafish (Curado et al., 2007; Pisharath et al., 2007). Recently, a more potent prodrug, nifurpirinol, has been identified as a more efficient and specific compound (Bergemann et al., 2018). In all cases, genetically targeted cells undergo apoptosis. However, in contrast to the cryoinjury model, ECM deposition is limited (Wang et al., 2011). Following cell death, cells from the surrounding healthy tissue begin proliferating to replace the lost tissue.

Genetic tools

As in other fields of zebrafish research, the use of genetic tools is essential to understand the role played by different genes during regeneration. In Table 3 we summarize most of the genetic modification approaches discussed in the ensuing section and include the main advantages and disadvantages of each of them.

Transgenesis

There is a wide range of tools available for the generation of transgenic lines. The most commonly used include the use of Tol2 transposons (Kawakami et al., 2000) and I-SceI meganuclease (Thermes et al., 2002), which facilitate the integration of foreign DNA into the fish genome. Alternatively, BAC recombineering (Bussmann and Schulte-Merker, 2011; Suster et al., 2011) allows

the insertion of larger fragments of DNA that often better recapitulate endogenous gene expression patterns. Of note, insertion into the genome is non-directed in these approaches. The lack of a ROSA26-like landing site has been a caveat; the use of Phi31 Integrase transgenesis (Mosimann et al., 2013) has been reported as a potential way to address this, and the problem will hopefully be solved using CRISPR/Cas9 approaches. Transgenic fluorescent reporter lines used to tag/track a particular cell type or report a gene expression pattern have proven to be essential for investigating regeneration.

Inducible gene expression and genetic fate mapping

Different techniques have been established that allow the use of genetic tools in a spatially and/or temporally controlled manner during regeneration.

Heat shock

Inducible gene expression can be achieved by using a heat-shock driven system (Ádám et al., 2000). The heat-shock promoter hsp70 is used to drive expression of the gene of interest. By maintaining fish at 35–39°C instead of 28°C, gene expression can be activated.

Gal4/UAS

One of the most widely used systems is the Gal4/UAS transactivation system, which was adapted from *Drosophila* to the

zebrafish. The system requires the combination of two separate transgenic lines – one expressing the Gal4 protein in a specific cell type and the other containing a construct in which a reporter (e.g. fluorescent protein) or a gene of interest is placed downstream of the UAS sequence, to which the Gal4 protein binds. When these lines are crossed, offspring are generated in which the UAS-driven gene is expressed in those cells expressing Gal4 (reviewed in greater detail by Halpern et al., 2008). The Gal4 system can also be used in an inducible manner, by fusing Gal4 to an ER^{T2} domain, so that Gal4 is expressed upon the administration of tamoxifen. This approach has been useful to study hair cell regeneration (Pinto-Teixeira et al., 2015). However, the Gal4/UAS system in zebrafish is prone to transcriptional silencing and accumulation of methylation (Akitake et al., 2011) over time, thus precluding its use in most cases to study regeneration in adults.

Cre/Lox

A further genetic tool for spatially controlled gene expression is the Cre/Lox recombination system (Felker and Mosimann, 2016; Hans et al., 2009; Langenau et al., 2005). It can also be used for lineage tracing, to assess the origin and fate of cell types during organ regeneration (Kikuchi et al., 2010). A number of systems allow temporal control – most commonly, through fusion of Cre to ER^{T2} (Feil et al., 1997) or by placing Cre under the control of the hsp70 promoter for heat-shock control (Hesselson et al., 2009). This system has been used for temporal gene expression control during the analysis of adult heart regeneration (González-Rosa et al., 2018). It is also possible to use Cre-based approaches to achieve both spatial and temporal control (Kirchgeorg et al., 2018).

TetOn/TetOff

This system allows temporally defined tissue-specific gene expression (Wehner et al., 2015). Its main advantage is that gene expression can be turned on and off by administration or removal of doxycycline – i.e. highly controlled transient expression is possible. It has been used to study Wnt signaling during fin and spinal cord regeneration (Wehner et al., 2014, 2017).

Photoconvertible fluorescent proteins

Finally, photoconvertible Kaede lines have proven useful when the aim is to trace a specific cell or cluster of cells during regeneration. This fluorescent protein undergoes conversion, from green to red light emission, when irradiated under a specific wave-length (Ando et al., 2002). This method has been used for *in vivo* tracing studies of cardiomyocytes during regeneration (Itou et al., 2012). Other photoconvertible proteins are also available.

Gene knock-down/knockout tools

For many decades, the analysis of gene function in the zebrafish relied on non-directed mutagenesis approaches followed by mapping of the mutation. This was followed by the use of transgenic gain-of-function studies or the use of dominant-negative forms or morpholino-based knockdown approaches. Morpholinos are traditionally injected at the 1-cell stage and so cannot easily be used to study regeneration. However, *in vivo*-Morpholinos have been developed (by GeneTools) that can be injected into the blood stream or electroporated into a specific tissue and pass through cell membranes. However, there are only few reports on their use and even fewer in the context of regeneration (Bednarek et al., 2015; Hyde et al., 2012; Kizil and Brand, 2011; Thummel et al., 2006). A better approach to understand the effect of the downregulation of a specific gene relies on the use of precise targeted gene editing tools such as TALENs and CRISPR/

Cas9. A better approach to understanding the effect of the downregulation of a specific gene relies on the use of precise targeted gene editing tools such as TALENs and CRISPR/Cas9. These have proven useful for generating null mutants in zebrafish (Hwang et al., 2013; Jao et al., 2013) and for assessing gene function during regeneration (Dupret et al., 2017; Dong et al., 2019; Nauroy et al., 2019; Unal Eroglu et al., 2018). The generation of conditional knockouts has also recently been achieved by the directed insertion of loxP sites using CRISPR/Cas9 (Burg et al., 2018). This new approach will be extremely valuable to address gene function during regeneration. Several publications compare the use of all these genetic engineering methods and can provide researchers with the necessary support to choose the best method for their studies (Chatterjee and Lufkin, 2011; Rafferty and Quinn, 2018). Although not yet well established, the possibility of using modified Cas9 versions for transcriptional modulation or induction of epigenetic modifications is also being explored in the zebrafish (e.g. Liu et al., 2019; Long et al., 2015).

Importantly, several screening studies, such as those from the Kawakami (Kawakami et al., 2010) and Brand (Jungke et al., 2013) labs, have led to the generation of large databases with numerous fish lines expressing Gal4 or Cre under specific promoters, which are available to the whole community. These approaches have been recently reviewed (Albadri et al., 2017; Carney and Mosimann, 2018). Similarly, large open source repositories for zebrafish mutants are available (Table 1).

Omics approaches

The different ‘omics tools’ that have become available in recent years are an important asset for helping researchers gain a deeper knowledge into the mechanisms regulating regeneration. Regarding genomic tools, ChIPseq (Kang et al., 2016; Xiao et al., 2016), Tomo-Seq (Wu et al., 2016), single cell RNAseq (Honkoop et al., 2018 preprint) and histone profiling (Goldman et al., 2017) have been successfully applied to study regeneration in the zebrafish. RNAseq data have provided information, for example, on genes involved in axonal regrowth following spinal cord injury (Mokalled et al., 2016). Complementing these approaches, proteomics (Saxena et al., 2012) and metabolomics (Rabinowitz et al., 2017) are now being used in the context of zebrafish organ regeneration. The possibility of sequencing-based fate mapping will significantly contribute to the study of organ regeneration (Spanjaard et al., 2018). Among the current limitations are the amount of tissue that can be selected for these approaches, annotations of the zebrafish genome and bioinformatics analysis platforms mainly oriented to mammalian models.

Live imaging

One of the main advantages of the zebrafish is the possibility of imaging morphogenetic processes. During embryogenesis, the use of fluorescent reporter lines facilitates imaging of tissues and organs at the cellular and subcellular level. Microscopic techniques such as light sheet microscopy allow for prolonged imaging with minimal interference of physiological processes that occur during either regeneration or development (Ding et al., 2018; Power and Huisken, 2017). The zebrafish larva is also sufficiently transparent that live imaging can be used to follow regenerative processes in organs and tissues in up to 5-day-old larvae or beyond, depending on the organ. The preferred injury model during these stages is genetic ablation, being particularly well suited given the small size of the animals at this age. Automated systems for larval imaging are being developed that facilitate the acquisition and analysis of data, and allow high-throughput screening assays (Early et al., 2018; Pandey et al., 2019).

In the case of adult zebrafish, *in vivo* imaging of regenerating internal organs can prove more difficult. Nevertheless, short- and long-term imaging of regeneration of external structures have been achieved using fluorescent light microscopy, as described for caudal fin neangiogenesis (Xu et al., 2015), regeneration of the skin epidermis (Chen et al., 2016) or scale regeneration (Cox et al., 2018). Although more challenging, even internal structures can be imaged and followed through time, as has been shown for adult neural stem cells in homeostasis and during regeneration (Barbosa et al., 2015; Dray et al., 2015). New fish transgenic lines have been developed in the hope of improving *in vivo* imaging of regenerative processes in the adult. Luciferase-based transgenic lines have proven useful for regeneration studies, particularly when looking at internal regenerating structures, as luciferase bioluminescence can be detected even in deep tissues (Chen et al., 2013).

Ultrasound or magnetic resonance imaging have also been used for studying functional recovery after lesion-induced damage (Goessling et al., 2007; González-Rosa et al., 2014; Hein et al., 2015; Koth et al., 2017).

Small molecules and drug screening

Zebrafish are also particularly appropriate for small-molecule and drug-screening studies (Mathew et al., 2007; Matsuda et al., 2018). In this regard, approaches using genetic ablation injuries in larvae are most valuable, as a large number of animals can be screened simultaneously. By using injured zebrafish larvae, multiple compounds can be tested at the same time and used to identify those with a pro-regenerative effect and those that would work to impair regeneration. Screens have identified small molecules with the potential to induce β -cell proliferation and insulin production (Matsuda et al., 2018; Tsuji et al., 2014), and have led to the identification of vitamin D as a potent inducer of cardiomyocyte proliferation in the context of heart regeneration (Han et al., 2019). The use of reporter lines to visualize cell cycle, such as the FUCCI lines (Sugiyama et al., 2009), has proven to be very useful in this regard as it allows one to assess cell proliferation by fluorescent protein expression (Han et al., 2019). The establishment of luciferase promoter lines might also be interesting for screening, in particular for internal structures (Chen et al., 2013).

Cellular and molecular mechanisms of organ regeneration in the zebrafish

So, what has been learned about organ regeneration in the zebrafish using all these tools? The following section aims to provide an overview of different sources of progenitor cells that contribute to rebuild a tissue or organ in this species, and describe some of the common regulatory mechanisms triggering this process.

Cellular origin

Depending on the tissue, and the degree or nature of damage involved, newly generated cells can originate from a number of potential sources (Fig. 3).

Proliferation of existing cell types

In some cases, cells not damaged by the injury can regenerate lost tissue through proliferation – this may involve a partial dedifferentiation such that the cells enter a proliferative state, followed by re-differentiation down the same lineage. This is thought to apply to heart regeneration, where pre-existing cardiomyocytes de-differentiate and proliferate (Jopling et al., 2010; Kikuchi et al., 2010) (Fig. 3Ai). Indeed, genetic lineage

tracing using a cardiomyocyte-specific promoter showed that all newly formed cardiomyocytes at the site of injury are derived from pre-existent myocardium. Regeneration can also proceed by proliferation of pre-existing cells, without the need for a dedifferentiation step before the proliferation phase. This has been described for liver regeneration – at least when the damage is not too severe and appears similar to the situation in mammals (Michalopoulos, 2007), where regeneration proceeds via proliferation of pre-existing hepatocytes (Sadler et al., 2007) (Fig. 3Aii).

Blastema formation

The replacement of a large area of tissue can occur through the formation of a blastema comprising undifferentiated cells, followed by cell proliferation and differentiation. A good illustration of this process is the regeneration of the zebrafish caudal fin. Upon amputation, an epidermal cap forms that covers the injury. Next, cells from the amputation plane dedifferentiate forming a blastema under the apical epidermal cap. During fin regeneration, the dedifferentiated cells proliferate and undergo re-differentiation to regenerate the various different cell types that make up the fin (reviewed by Pfefferli and Jazwińska, 2015). The dedifferentiation and proliferation of osteoblasts close to the amputated area make up the new rays (Stewart and Stankunas, 2012) (Fig. 3B). However, it has also been shown that fin regeneration remains viable in the absence of osteoblasts. In this case, new bone cells can originate from a reserve population of osteoblast precursor cells (Ando et al., 2017; Singh et al., 2012).

Transdifferentiation

In some cases, regeneration proceeds via dedifferentiation and proliferation followed by re-differentiation into a different mature cell type from the starting population. One example of this is the process following severe hepatocyte loss in the liver. Under these circumstances, regeneration proceeds initially via the proliferation of biliary epithelial cells (Fig. 3C), which differentiate into hepatocytes (Choi et al., 2014). Transdifferentiation has also been observed in zebrafish pancreatic regeneration. A study searching for genes that would promote β -cell regeneration identified a protein, *Igfbp1*, that potentiated β -cell regeneration by stimulating α - to β -cell transdifferentiation (Lu et al., 2016).

Differentiation of tissue-resident stem cells/progenitors

Not all organs regenerate through dedifferentiation and proliferation of mature cells. Injuries to kidney, skeletal muscle or neural tissues resolve by the proliferation and differentiation of stem cells. For example, kidney regeneration following chemical damage proceeds via the proliferation of adult nephron progenitor cells (Diep et al., 2011). As described in mammals (Wosczyzna and Rando, 2018; Young and John, 2000), skeletal muscle regeneration in zebrafish requires the proliferation of satellite cells, a population of adult Pax7⁺ muscle stem cells that, after an injury to the muscle, proliferate and fuse with adjacent muscle fibers (Berberoglu et al., 2017). Regeneration of neurons also has been shown to depend on a specific progenitor cell population. In the injured zebrafish telencephalon, new neurons are derived from a subpopulation of radial glial cells (Kroehne et al., 2011) that have been demonstrated to function as a stem cells population in uninjured brain tissue (Rothenaigner et al., 2011) (Fig. 3D).

Environmental cues driving regeneration

Initial wound closure is followed by an inflammatory response mediated by neutrophils and macrophages, which are attracted by




Fig. 3. Cellular origin of the *de novo* formed tissue during organ regeneration in the zebrafish. (A) Dedifferentiation, proliferation and re-differentiation. (Ai) In the heart, cardiomyocytes in close proximity to the injury revert to a less differentiated stage, re-enter the cell cycle and redifferentiate into mature cardiomyocytes. (Aii) During regeneration of minor liver damage, hepatocyte regeneration occurs with no signs of dedifferentiation prior to cell cycle entry and proliferation. (B) Blastema formation as an intermediate step during regeneration. After fin amputation, cells of various lineages – including osteoblasts – dedifferentiate and accumulate under an apical epidermal cap. They then proliferate and redifferentiate to rebuild the missing fin structures. (C) Phenotypic switch or transdifferentiation during regeneration. Example: after extensive liver damage, biliary ductal cells (green) can transdifferentiate into hepatocytes (green hexagonal cells) that then differentiate into mature proliferating hepatocytes. (D) Stem cells as progenitor cells. Neural stem cells/progenitor cells proliferate and differentiate into new neurons during regeneration of the central nervous system. While neuronal regeneration has been well described, less information is available on robust axon regrowth. Yellow, differentiated cells; orange, dedifferentiated cells; purple, non-osteoblast cells within the fin; green hexagonal cells, cells undergoing transdifferentiation; blue, stem cells/progenitor cells. Damaged area is shown in gray.

signals generated from the dying cells and surrounding tissue, and home to the site of damage. Indeed, inflammation is essential for organ regeneration (Mescher, 2017). An inflammatory response precedes

zebrafish brain regeneration (Kyritsis et al., 2012) and has been suggested to be important for heart regeneration (Lai et al., 2017). Furthermore, T lymphocytes are involved in the regeneration of several

organs, such as spinal cord, heart and retina (Hui et al., 2017). Supporting these findings, genetic or chemical ablation of phagocytic macrophages during regeneration results in impaired fin (Li et al., 2012; Petrie et al., 2014), lateral line (Carrillo et al., 2016), heart (Lai et al., 2017) and spinal cord (Tsarouchas et al., 2018) regeneration.

The ECM and the stromal environment also influence regeneration. Not only do they provide structural support for progenitor cells, but they also have an impact on their proliferation and differentiation. Indeed, fibrosis and regeneration are tightly linked processes. For example, the pro-fibrotic molecule TGF β induces not only fibrosis but also cardiomyocyte proliferation (Chablais and Jazwinska, 2012). Moreover, ablation of collagen-producing cells impairs cardiomyocyte proliferation after heart injury (Sánchez-Iranzo et al., 2018), and ECM proteins are required for spinal cord regeneration (Mokalled et al., 2016; Wehner et al., 2017).

The nervous system is also key for adequate regeneration. This has been demonstrated both in the fin, where denervation following amputation results in impaired regeneration (Simões et al., 2014), and in the heart, where regeneration is also impaired following inhibition of peripheral nervous system activity (Mahmoud et al., 2015). Future work is needed to further unravel the role of the nervous system during organ regeneration, which can range from secretion of molecules to establishment of synaptic connections.

Finally, revascularization is also essential for proper regeneration. When this process is impaired during cardiac injury, regeneration is blocked. Studies conducted on this subject suggest that revascularization of the injured area is essential to stimulate cardiomyocyte proliferation (Harrison et al., 2015; Marin-Juez et al., 2016). The exact mechanisms through which revascularization contributes to regeneration remain to be discovered.

Molecular pathways promoting regeneration

The key signaling pathways involved in embryonic development are often also involved in organ regeneration. To study their function, researchers have used transgenic lines for temporal and tissue-specific overexpression of a core gene of a particular pathway and studied the effect on organ regeneration. In addition, analysis of organ regeneration in mutants of a gene involved in the pathway of interest has been informative. Here, we provide a few examples of how such pathways contribute to regenerative processes in zebrafish; we refer the reader to the papers cited for further details. One example of a pathway important for regeneration is the retinoic acid (RA) pathway. RA regulates FGF, Wnt/ β -catenin and Igf signaling during blastema formation in the regenerating caudal fin (Blum and Begemann, 2012), and stimulates cardiomyocyte proliferation, likely in a paracrine manner (Kikuchi et al., 2011). The fibroblast growth factor (FGF) and bone morphogenetic protein (BMP) pathways, which are essential for organ growth and differentiation (Kan et al., 2009; Smith et al., 2006), are also involved in regeneration. BMP signaling has been implicated in the regeneration of several organs, including the liver, the heart and the fin (Kan et al., 2009; Sehring et al., 2016). Similarly, FGF is involved in multiple processes of regeneration, including the control of blastema proliferation during fin regeneration (Lee et al., 2005; Poss et al., 2000) or heart regeneration (Lepilina et al., 2006). One of the most universally used pathways in tissue regeneration throughout the animal kingdom is the Wnt signaling pathway (Slack, 2017). In the zebrafish, it regulates blastema proliferation in amputated caudal fin among many other processes (Stoick-Cooper et al., 2007; Wehner et al., 2017). Juxtacrine signaling pathways, such as the Notch pathway are also associated with proliferation control both in fin (Münch et al., 2013) and heart (Zhao et al., 2014).

Other pathways involved in the control of cell cycle include the Hippo-Yap/TAZ pathway (Flinn et al., 2018; Mateus et al., 2015), which has recently received considerable attention in the field of regeneration because it links the effect of a changing environment and mechanical forces in response to injury with the regenerative response in the surrounding cells. In all cases, specificity and temporal control of these pathways is crucial for promoting regeneration. In addition, depending on the combination of active pathways, different cellular responses are elicited, ranging from cell differentiation or proliferation to migration to epithelial-to-mesenchymal transition, all of which are crucial for organ regeneration.

In sum, during zebrafish regeneration, cells that repopulate injured areas can have different origins, depending not only on the lesioned organ, but also on the extent and the type of injury. Furthermore, for regeneration to succeed there must be a balance between the signals that induce cell proliferation and those that halt cell division and determine cell fate.

Extent and limits of the regenerative capacity in zebrafish

Without doubt, the zebrafish is an organism with an amazing capacity for regeneration. Here, we discuss some of the mechanisms that might underlie this capacity, but also highlight some possible limitations of organ regeneration in this animal.

Possible explanations for the high regenerative capacity

The regenerative capacity in the zebrafish might rely on its capacity to activate specific gene regulatory networks in response to injury that are repressed in other animal models (Yang and Kang, 2018). Indeed, epigenetic modifications are necessary to trigger regeneration. One of the earliest findings in this field reported that, in the fin, histone modifications precede initiation of regeneration (Stewart et al., 2009), and histone methyltransferases are necessary for this process (Dupret et al., 2017). The use of ChIPseq and transgenesis has provided much insight into the landscape of active enhancers during organ regeneration, identifying enhancer elements that are activated in response to injury consistently in different organs and even in different organisms (Kang et al., 2016; Pfefferli and Jaźwińska, 2017).

One important characteristic of zebrafish is its capacity to grow throughout its lifespan, meaning that most of the zebrafish cells, e.g. cardiomyocytes, retain their proliferative capacity (Wills et al., 2008). For heart regeneration, cardiomyocyte ploidy has been shown to be a limiting step. Zebrafish cardiomyocytes are mononuclear and diploid, whereas in adult humans they are mostly polyploid. Indeed, induction of polyploidy in zebrafish cardiomyocytes abrogates heart regeneration (González-Rosa et al., 2018; Patterson et al., 2017).

The immune system might play a key role in defining the regenerative capacity of an organism. As an example, differences in immune response to cardiac injury in two teleost models (medaka and zebrafish) correlate with reduced regenerative capacity of medaka (Lai et al., 2017). Finally, regenerative capacities might also be attributed to differences in the environment that species inhabit – exemplified by the differences in heart regeneration in surface versus cave morphs of the cavefish *Astyanax mexicanus* (Stockdale et al., 2018). Thus, using other fish models with different tissue recovery capacities for regeneration studies (see Table 4) may contribute to a better understanding of the different mechanisms underlying this process.

Unlimited regeneration in the zebrafish?

Regeneration in the zebrafish seems to be nearly unlimited: regeneration of the caudal fin and barbell occurs even after

Table 4. Other teleost fish models of regeneration

Fish model	Organs in which regeneration has been studied
<i>Astyanax mexicanus</i> (Mexican cave fish)	Heart (Stockdale et al., 2018)
<i>Carassius auratus</i> (Goldfish)	Spinal cord (Bernstein and Gelderd, 1970) Eye (Parrilla et al., 2012) Fin (Caskey and O'Brien, 1948)
<i>Oryzias latipes</i> (Medaka)	Heart (Grivas et al., 2014) Kidney (Watanabe et al., 2009) Fin (Nemoto et al., 2007) Pancreatic β -cells (Otsuka and Takeda, 2017)

repetitive amputations (Azevedo et al., 2011; LeClair and Topczewski, 2010), although subtle changes to the newly formed organ, such as variation in the pigmentation patterning of the fin or the position of the bony ray bifurcations, have been observed (Azevedo et al., 2012). Moreover, age does not seem to limit regenerative capacity, at least for the heart (Itou et al., 2012). Nevertheless, there is evidence that organ regeneration can be partially incomplete. For example, although heart morphology is restored after lesion and no scar is visible, the ventricular wall motion of the regenerated area remains partially affected and contracts asynchronously with the remainder of the heart, even after complete tissue regeneration (González-Rosa et al., 2014). Furthermore, following resorption of extracellular fibrotic components during heart healing, accumulated fibroblasts are not fully eliminated after complete regeneration but become inactivated (Sánchez-Iranzo et al., 2018). It is thus important to distinguish between morphological and functional regeneration and to recognize that – even in a system such as zebrafish – regeneration may not be functionally ‘perfect’.

Conclusions and perspectives

In conclusion, using the zebrafish as an animal model in regeneration studies has undoubtedly contributed to the development of the field but, as with any other model, there are advantages and disadvantages to consider (Table 5). Despite the contributions made, there are still unanswered questions. For example, there is little information on how metabolism is linked to regenerative capacity. Furthermore, the interplay between fibrosis and regeneration, as well as the mechanisms of fibrotic tissue regression, are still not fully understood. Why is a scar formed under some circumstances or in

Table 5. Advantages and disadvantages of the zebrafish as a regeneration model

Advantages	Disadvantages
High regenerative capacity	Non-mammalian model
Vertebrate model	Partial genome duplication
Established husbandry protocols	Dearth of antibodies for protein studies
Low-cost maintenance	Difficulty in performing <i>in vivo</i> imaging in adult stages
External development	Difficulty in using some molecular tools (e.g. knock-ins)
Possibility of working with large numbers	Anatomical and physiological differences to humans (e.g. two-chambered heart)
Ease of <i>in vivo</i> imaging during development	
Genetic engineering, annotated genome	
Physiological similarities with humans (e.g. heart rate)	

some organs while in other cases regeneration occurs? To what extent does regeneration rely on a transient fibrotic scar? It is also important to understand how mechanical forces influence regeneration – studies in the zebrafish might help to explain how, for example, stiffness or shear stress regulates organ regeneration. In addition, it is intriguing that regenerating tissues stop growing at the appropriate size; this is poorly understood. Indeed, there is still a dearth of knowledge regarding which cell populations become active during regeneration and whether all the cells have the same regenerative potential. To answer many of these questions, some of the molecular tools such as conditional genome editing will need to be improved and widely established for regeneration studies. Advances in *intra vital* imaging for adult zebrafish will also help to expand our toolbox. With these new technological advances in hand, important discoveries on the mechanisms of organ regeneration using the zebrafish model will certainly be achieved. A deep knowledge on the mechanisms of organ regeneration in species such as the zebrafish with innate regenerative capacity might serve as inspiration for the development of therapeutic strategies in mammals.

Acknowledgements

We thank Andres Sanz for critical reading of the manuscript and the reviewers for their suggestions. We also acknowledge all those authors whose work could not be cited here due to space limitations. Their research keeps providing us with invaluable knowledge in the field of zebrafish regeneration.

Competing interests

The authors declare no competing or financial interests.

Funding

N.M. is supported by a European Research Council starting grant (2013 zebraHeart-337703), E.L. is supported by the European Commission (H2020-MSCA-ITN-2016 European Industrial Doctorate 4DHeart 722427).

References

- Ádám, A., Bártfai, R., Lele, Z., Krone, P. H. and Orbán, L. (2000). Heat-inducible expression of a reporter gene detected by transient assay in zebrafish. *Exp. Cell Res.* **256**, 282-290. doi:10.1006/excr.2000.4805
- Akitake, C. M., Macurak, M., Halpern, M. E. and Goll, M. G. (2011). Transgenerational analysis of transcriptional silencing in zebrafish. *Dev. Biol.* **352**, 191-201. doi:10.1016/j.ydbio.2011.01.002
- Albadri, S., Del Bene, F. and Revenu, C. (2017). Genome editing using CRISPR/Cas9-based knock-in approaches in zebrafish. *Methods* **121-122**, 77-85. doi:10.1016/j.ymeth.2017.03.005
- Ando, R., Hama, H., Yamamoto-Hino, M., Mizuno, H. and Miyawaki, A. (2002). An optical marker based on the UV-induced green-to-red photoconversion of a fluorescent protein. *Proc. Natl Acad. Sci. USA* **99**, 12651. doi:10.1073/pnas.202320599
- Ando, K., Shibata, E., Hans, S., Brand, M. and Kawakami, A. (2017). Osteoblast production by reserved progenitor cells in zebrafish bone regeneration and maintenance. *Dev. Cell* **43**, 643-650.e643. doi:10.1016/j.devcel.2017.10.015
- Azevedo, A. S., Grotek, B., Jacinto, A., Weidinger, G. and Saúde, L. (2011). The regenerative capacity of the zebrafish caudal fin is not affected by repeated amputations. *PLoS ONE* **6**, e22820. doi:10.1371/journal.pone.0022820
- Azevedo, A. S., Sousa, S., Jacinto, A. and Saúde, L. (2012). An amputation resets positional information to a proximal identity in the regenerating zebrafish caudal fin. *BMC Dev. Biol.* **12**, 24. doi:10.1186/1471-213X-12-24
- Barbosa, J. S., Sanchez-Gonzalez, R., Di Giaimo, R., Baumgart, E. V., Theis, F. J., Gotz, M. and Ninkovic, J. (2015). Neurodevelopment. Live imaging of adult neural stem cell behavior in the intact and injured zebrafish brain. *Science* **348**, 789-793. doi:10.1126/science.aaa2729
- Becker, C. G. and Becker, T. (2008). Adult zebrafish as a model for successful central nervous system regeneration. *Restor. Neurol. Neurosci.* **26**, 71-80.
- Becker, T., Wullimann Mario, F., Becker Catherina, G., Bernhardt, R. R. and Schachner, M. (1998). Axonal regrowth after spinal cord transection in adult zebrafish. *J. Comp. Neurol.* **377**, 577-595. doi:10.1002/(SICI)1096-9861(19970127)377:4<577::AID-CNE8>3.0.CO;2-#
- Bednarek, D., González-Rosa, J. M., Guzmán-Martínez, G., Gutiérrez-Gutiérrez, Ó., Aguado, T., Sánchez-Ferrer, C., Marques, I. J., Galardi-Castilla, M., de Diego, I., Gómez, M. J. et al. (2015). Telomerase is essential for zebrafish heart regeneration. *Cell Reports* **12**, 1691-1703. doi:10.1016/j.celrep.2015.07.064
- Berberoglu, M. A., Gallagher, T. L., Morrow, Z. T., Talbot, J. C., Hromowky, K. J., Tenente, I. M., Langenau, D. M. and Amacher, S. L. (2017). Satellite-like cells

- contribute to pax7-dependent skeletal muscle repair in adult zebrafish. *Dev. Biol.* **424**, 162-180. doi:10.1016/j.ydbio.2017.03.004
- Bergemann, D., Massoz, L., Bourdouxhe, J., Carril Pardo, C. A., Voz, M. L., Peers, B. and Manfried, I.** (2018). Nifurpirinol: a more potent and reliable substrate compared to metronidazole for nitroreductase-mediated cell ablations. *Wound Repair. Regen.* **26**, 238-244. doi:10.1111/wrr.12633
- Bergmann, O., Bhardwaj, R. D., Bernard, S., Zdunek, S., Barnabe-Heider, F., Walsh, S., Zupicich, J., Alkass, K., Buchholz, B. A., Druid, H. et al.** (2009). Evidence for cardiomyocyte renewal in humans. *Science* **324**, 98-102. doi:10.1126/science.1164680
- Bernhardt, R. R., Tongiorgi, E., Anzini, P. and Schachner, M.** (1996). Increased expression of specific recognition molecules by retinal ganglion cells and by optic pathway glia accompanies the successful regeneration of retinal axons in adult zebrafish. *J. Comp. Neurol.* **376**, 253-264. doi:10.1002/(SICI)1096-9861(19961209)376:2<253::AID-CNE7>3.0.CO;2-2
- Bernstein, J. J. and Gelder, J. B.** (1970). Regeneration of the long spinal tracts in the goldfish. *Brain Res.* **20**, 33-38. doi:10.1016/0006-8993(70)90151-4
- Blum, N. and Begemann, G.** (2012). Retinoic acid signaling controls the formation, proliferation and survival of the blastema during adult zebrafish fin regeneration. *Development* **139**, 107-116. doi:10.1242/dev.065391
- Burg, L., Palmer, N., Kikhi, K., Miroshnik, E. S., Rueckert, H., Gaddy, E., MacPherson Cunningham, C., Mattonet, K., Lai, S. L., Marin-Juez, R. et al.** (2018). Conditional mutagenesis by oligonucleotide-mediated integration of loxP sites in zebrafish. *PLoS Genet.* **14**, e1007754. doi:10.1371/journal.pgen.1007754
- Burkhardt-Holm, P., Oulmi, Y., Schroeder, A., Storch, V. and Braunbeck, T.** (1999). Toxicity of 4-chloroaniline in early life stages of zebrafish (*Danio rerio*): II. Cytopathology and regeneration of liver and gills after prolonged exposure to waterborne 4-chloroaniline. *Arch. Environ. Contam. Toxicol.* **37**, 85-102. doi:10.1007/s002449900493
- Bussmann, J. and Schulte-Merker, S.** (2011). Rapid BAC selection for tol2-mediated transgenesis in zebrafish. *Development* **138**, 4327-4332. doi:10.1242/dev.068080
- Cano-Martinez, A., Vargas-Gonzalez, A., Guarner-Lans, V., Prado-Zayago, E., Leon-Oleda, M. and Nieto-Lima, B.** (2010). Functional and structural regeneration in the axolotl heart (*Ambystoma mexicanum*) after partial ventricular amputation. *Arch. Cardiol. Mex.* **80**, 79-86.
- Carney, T. J. and Mosimann, C.** (2018). Switch and trace: recombinase genetics in zebrafish. *Trends Genet.* **34**, 362-378. doi:10.1016/j.tig.2018.01.004
- Carrillo, S. A., Anguita-Salinas, C., Peña, O. A., Morales, R. A., Muñoz-Sánchez, S., Muñoz-Montecinos, C., Paredes-Zúñiga, S., Tapia, K. and Allende, M. L.** (2016). Macrophage Recruitment contributes to regeneration of mechanosensory hair cells in the zebrafish lateral line. *J. Cell. Biochem.* **117**, 1880-1889. doi:10.1002/jcb.25487
- Caskey, J. M. and O'Brien, J. P.** (1948). The cellular basis of regeneration of the caudal fin of the goldfish. *Anat. Rec.* **101**, 710.
- Chablais, F. and Jazwinska, A.** (2012). The regenerative capacity of the zebrafish heart is dependent on TGFbeta signaling. *Development* **139**, 1921-1930. doi:10.1242/dev.078543
- Chablais, F., Veit, J., Rainer, G., Rainer, G. and Jaźwińska, A.** (2011). The zebrafish heart regenerates after cryoinjury-induced myocardial infarction. *BMC Dev. Biol.* **11**, 21. doi:10.1186/1471-213X-11-21
- Chassot, B., Pury, D. and Jaźwińska, A.** (2016). Zebrafish fin regeneration after cryoinjury-induced tissue damage. *Biol. Open* **5**, 819-828. doi:10.1242/bio.016865
- Chatterjee, S. and Lufkin, T.** (2011). Fishing for function: zebrafish BAC transgenics for functional genomics. *Mol. Biosyst.* **7**, 2345-2351. doi:10.1039/c1mb05116d
- Chen, C.-H., Durand, E., Wang, J., Zon, L. I. and Poss, K. D.** (2013). zebrafish transgenic lines for in vivo bioluminescence imaging of stem cells and regeneration in adult zebrafish. *Development* **140**, 4988-4997. doi:10.1242/dev.120253
- Chen, C.-H., Puliafito, A., Cox, B. D., Primo, L., Fang, Y., Di Talia, S. and Poss, K. D.** (2016). Multicolor cell barcoding technology for long-term surveillance of epithelial regeneration in zebrafish. *Dev. Cell* **36**, 668-680. doi:10.1016/j.devcel.2016.02.017
- Choi, T.-Y., Ninov, N., Stainier, D. Y. R. and Shin, D.** (2014). Extensive conversion of hepatic biliary epithelial cells to hepatocytes after near total loss of hepatocytes in zebrafish. *Gastroenterology* **146**, 776-788. doi:10.1053/j.gastro.2013.10.019
- Conedera, F. M., Arendt, P., Trepp, C., Tschopp, M., Enzmann, V.** (2017). Muller glia cell activation in a laser-induced retinal degeneration and regeneration model in zebrafish. *J. Vis. Exp.* **128**, e56249. doi:10.3791/56249
- Cox, A. G., Saunders, D. C., Kelsey, P. B., Jr, Conway, A. A., Tesmenitsky, Y., Marchini, J. F., Brown, K. K., Stamlir, J. S., Colagiovanni, D. B., Rosenthal, G. J. et al.** (2014). S-nitrosothiol signaling regulates liver development and improves outcome following toxic liver injury. *Cell Reports* **6**, 56-69. doi:10.1016/j.celrep.2013.12.007
- Cox, B. D., De Simone, A., Tornini, V. A., Singh, S. P., Di Talia, S. and Poss, K. D.** (2018). In toto imaging of dynamic osteoblast behaviors in regenerating skeletal bone. *Curr. Biol.* **28**, 3937-3947.e3934. doi:10.1016/j.cub.2018.10.052
- Curado, S., Anderson, R. M., Jungblut, B., Mumm, J., Schroeter, E. and Stainier, D. Y.** (2007). Conditional targeted cell ablation in zebrafish: a new tool for regeneration studies. *Dev. Dyn.* **236**, 1025-1035. doi:10.1002/dvdy.21100
- Del Bene, F., Wyart, C., Robles, E., Tran, A., Looger, L., Scott, E. K., Isacoff, E. Y. and Baier, H.** (2010). Filtering of visual information in the tectum by an identified neural circuit. *Science* **330**, 669-673. doi:10.1126/science.1192949
- Diep, C. Q., Ma, D., Deo, R. C., Holm, T. M., Naylor, R. W., Arora, N., Wingert, R. A., Bollig, F., Djordjevic, G., Lichman, B. et al.** (2011). Identification of adult nephron progenitors capable of kidney regeneration in zebrafish. *Nature* **470**, 95. doi:10.1038/nature09669
- Ding, Y., Lee, J., Hsu, J. J., Chang, C.-C., Baek, K. I., Ranjbarvaziri, S., Ardehali, R., Packard, R. R. S. and Hsiai, T. K.** (2018). Light-sheet imaging to elucidate cardiovascular injury and repair. *Curr. Cardiol. Rep.* **20**, 35. doi:10.1007/s11886-018-0979-6
- Dong, Z., Wu, S., Zhu, C., Wang, X., Li, Y., Chen, X., Liu, D., Qiang, L., Baas, P. W. and Liu, M.** (2019). Clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9-mediated kif15 mutations accelerate axonal outgrowth during neuronal development and regeneration in zebrafish. *Traffic* **20**, 71-81. doi:10.1111/tra.12621
- Dray, N., Bedu, S., Vuillemin, N., Alunni, A., Coolen, M., Krecsmarik, M., Supatto, W., Beaupaire, E. and Bally-Cuif, L.** (2015). Large-scale live imaging of adult neural stem cells in their endogenous niche. *Development* **142**, 3592-3600. doi:10.1242/dev.123018
- Dupret, B., Volkel, P., Vennin, C., Toillon, R. A., Le Bourhis, X. and Angrand, P. O.** (2017). The histone lysine methyltransferase Ezh2 is required for maintenance of the intestine integrity and for caudal fin regeneration in zebrafish. *Biochim. Biophys. Acta. Gene Regul. Mech.* **1860**, 1079-1093. doi:10.1016/j.bbagr.2017.08.011
- Early, J. J., Cole, K. L. H., Williamson, J. M., Swire, M., Kamadurai, H., Muskavitch, M. and Lyons, D. A.** (2018). An automated high-resolution in vivo screen in zebrafish to identify chemical regulators of myelination. *eLife* **7**, e35136. doi:10.7554/eLife.35136
- Ernst, A., Alkass, K., Bernard, S., Salehpour, M., Perl, S., Tisdale, J., Possner, G., Druid, H. and Frisé, J.** (2014). Neurogenesis in the striatum of the adult human brain. *Cell* **156**, 1072-1083. doi:10.1016/j.cell.2014.01.044
- Feil, R., Wagner, J., Metzger, D. and Chambon, P.** (1997). Regulation of Cre recombinase activity by mutated estrogen receptor ligand-binding domains. *Biochem. Biophys. Res. Commun.* **237**, 752-757. doi:10.1006/bbrc.1997.7124
- Felker, A. and Mosimann, C.** (2016). Contemporary zebrafish transgenesis with Tol2 and application for Cre/lox recombination experiments. *Methods Cell Biol.* **135**, 219-244. doi:10.1016/bs.mcb.2016.01.009
- Fimbel, S. M., Montgomery, J. E., Burket, C. T., Burket, C. T., Hyde, D. R. and Hyde, D. R.** (2007). Regeneration of inner retinal neurons after intravitreal injection of ouabain in zebrafish. *J. Neurosci.* **27**, 1712-1724. doi:10.1523/JNEUROSCI.5317-06.2007
- Flinn, M. A., Jeffery, B. E., O'Meara, C. C. and Link, B. A.** (2018). Yap is required for scar formation but not myocyte proliferation during heart regeneration in zebrafish. *Cardiovasc. Res.* **115**, 570-577. doi:10.1093/cvr/cvy243
- Geisler, R., Borel, N., Ferg, M., Maier, J. V. and Strahle, U.** (2016). Maintenance of zebrafish lines at the european zebrafish resource center. *Zebrafish* **13** Suppl. 1, S19-S23. doi:10.1089/zeb.2015.1205
- Geraudie, J., Monnot, M. J., Ridet, A., Thorogood, P. and Ferretti, P.** (1993). Is exogenous retinoic acid necessary to alter positional information during regeneration of the fin in zebrafish? *Prog. Clin. Biol. Res.* **383B**, 803-814.
- Goessling, W., North, T. E. and Zon, L. I.** (2007). Ultrasound biomicroscopy permits in vivo characterization of zebrafish liver tumors. *Nat. Methods* **4**, 551-553. doi:10.1038/nmeth1059
- Goldman, J. A., Kuzu, G., Lee, N., Karasik, J., Gemberling, M., Foglia, M. J., Karra, R., Dickson, A. L., Sun, F., Tolstorukov, M. Y. et al.** (2017). Resolving heart regeneration by replacement histone profiling. *Dev. Cell* **40**, 392-404.e395. doi:10.1016/j.devcel.2017.01.013
- Gonzalez-Rosa, J. M., Martin, V., Peralta, M., Torres, M. and Mercader, N.** (2011). Extensive scar formation and regression during heart regeneration after cryoinjury in zebrafish. *Development* **138**, 1663-1674. doi:10.1242/dev.060897
- Gonzalez-Rosa, J. M., Peralta, M. and Mercader, N.** (2012). Pan-epicardial lineage tracing reveals that epicardium derived cells give rise to myofibroblasts and perivascular cells during zebrafish heart regeneration. *Dev. Biol.* **370**, 173-186. doi:10.1016/j.ydbio.2012.07.007
- González-Rosa, J. M., Guzmán-Martínez, G., Marques, I. J., Sánchez-Iranzo, H., Jiménez-Borreguero, L. J. and Mercader, N.** (2014). Use of echocardiography reveals reestablishment of ventricular pumping efficiency and partial ventricular wall motion recovery upon ventricular cryoinjury in the zebrafish. *PLoS ONE* **9**, e115604. doi:10.1371/journal.pone.0115604
- Gonzalez-Rosa, J. M., Burns, C. E. and Burns, C. G.** (2017). Zebrafish heart regeneration: 15 years of discoveries. *Regeneration* **4**, 105-123. doi:10.1002/reg.2.83
- González-Rosa, J. M., Sharpe, M., Field, D., Soonpaa, M. H., Field, L. J., Burns, C. E. and Burns, C. G.** (2018). Myocardial polyploidization creates a barrier to heart regeneration in zebrafish. *Dev. Cell* **44**, 433-446.e7. doi:10.1016/j.devcel.2018.01.021

- Grivas, J., Haag, M., Johnson, A., Manalo, T., Roell, J., Das, T. L., Brown, E., Burns, A. R. and Lafontant, P. J. (2014). Cardiac repair and regenerative potential in the goldfish (*Carassius auratus*) heart. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* **163**, 14-23. doi:10.1016/j.cbpc.2014.02.002
- Haffter, P., Granato, M., Brand, M., Mullins, M. C., Hammerschmidt, M., Kane, D. A., Odenthal, J., van Eeden, F. J., Jiang, Y. J., Heisenberg, C. P. et al. (1996). The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**, 1-36. doi:10.1016/0736-5748(96)80313-3
- Halpern, M. E., Rhee, J., Goll, M. G., Akitake, C. M., Parsons, M. and Leach, S. D. (2008). Gal4/UAS transgenic tools and their application to zebrafish. *Zebrafish* **5**, 97-110. doi:10.1089/zeb.2008.0530
- Han, Y., Chen, A., Umansky, K. B., Oonk, K. A., Choi, W. Y., Dickson, A. L., Ou, J., Cigliola, V., Yifa, O., Cao, J. et al. (2019). Vitamin D stimulates cardiomyocyte proliferation and controls organ size and regeneration in zebrafish. *Dev. Cell* **48**, 853-863.e855. doi:10.1016/j.devcel.2019.01.001
- Hans, S., Kaslin, J., Freudenreich, D. and Brand, M. (2009). Temporally-controlled site-specific recombination in zebrafish. *PLoS ONE* **4**, e4640-e4640. doi:10.1371/journal.pone.0004640
- Hardy, M. A. (1989). The biology of scar formation. *Phys. Ther.* **69**, 1014-1024. doi:10.1093/ptj/69.12.1014
- Harris, J. A., Cheng, A. G., Cunningham, L. L., MacDonald, G., Raible, D. W. and Rubel, E. W. (2003). Neomycin-induced hair cell death and rapid regeneration in the lateral line of zebrafish (*Danio rerio*). *J. Assoc. Res. Otolaryngol.* **4**, 219-234. doi:10.1007/s10162-002-3022-x
- Harrison, M. R., Bussmann, J., Huang, Y., Zhao, L., Osorio, A., Burns, C. G., Burns, C. E., Sucov, H. M., Siekmann, A. F. and Lien, C. L. (2015). Chemokine-guided angiogenesis directs coronary vasculature formation in zebrafish. *Dev. Cell* **33**, 442-454. doi:10.1016/j.devcel.2015.04.001
- Hein, S. J., Lehmann, L. H., Kossack, M., Juergensen, L., Fuchs, D., Katus, H. A. and Hassel, D. (2015). Advanced echocardiography in adult zebrafish reveals delayed recovery of heart function after myocardial cryoinjury. *PLoS ONE* **10**, e0122665. doi:10.1371/journal.pone.0122665
- Hesselson, D., Anderson, R. M., Beinat, M. and Stainier, D. Y. R. (2009). Distinct populations of quiescent and proliferative pancreatic beta-cells identified by HOTcre mediated labeling. *Proc. Natl. Acad. Sci. USA* **106**, 14896-14901. doi:10.1073/pnas.0906348106
- Honkoop, H., de Bakker, D., Aharonov, A., Kruse, F., Shakked, A., Nguyen, P., de Heus, C., Garric, L., Muraro, M., Shoffner, A. et al. (2018). A metabolic switch from OXPHOS to glycolysis is essential for cardiomyocyte proliferation in the regenerating heart. *bioRxiv* 498899. doi:10.1101/498899
- Howe, D. G., Bradford, Y. M., Conlin, T., Eagle, A. E., Fashena, D., Frazer, K., Knight, J., Mani, P., Martin, R., Moxon, S. A. et al. (2013). ZFIN, the zebrafish model organism database: increased support for mutants and transgenics. *Nucleic Acids Res.* **41**, D854-D860. doi:10.1093/nar/gks938
- Hui, S. P., Sheng, D. Z., Sugimoto, K., Gonzalez-Rajal, A., Nakagawa, S., Hesselson, D. and Kikuchi, K. (2017). Zebrafish regulatory T cells mediate organ-specific regenerative programs. *Dev. Cell* **43**, 659-672.e655. doi:10.1016/j.devcel.2017.11.010
- Hwang, W. Y., Fu, Y., Reyon, D., Maeder, M. L., Tsai, S. Q., Sander, J. D., Peterson, R. T., Yeh, J. R. and Joung, J. K. (2013). Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 227-229. doi:10.1038/nbt.2501
- Hyde, D. R., Godwin, A. R., Thummel, R. (2012). In vivo electroporation of morpholinos into the regenerating adult zebrafish tail fin. *J. Vis. Exp.* **61**, 3632. doi:10.3791/3632
- Itou, J., Oishi, I., Kawakami, H., Glass, T. J., Richter, J., Johnson, A., Lund, T. C. and Kawakami, Y. (2012). Migration of cardiomyocytes is essential for heart regeneration in zebrafish. *Development* **139**, 4133. doi:10.1242/dev.079756
- Jao, L.-E., Wente, S. R. and Chen, W. (2013). Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. USA* **110**, 13904-13909. doi:10.1073/pnas.1308335110
- Johnson, S. L. and Weston, J. A. (1995). Temperature-sensitive mutations that cause stage-specific defects in Zebrafish fin regeneration. *Genetics* **141**, 1583-1595.
- Jopling, C., Sleep, E., Raya, M., Marti, M., Raya, A. and Belmonte, J. C. I. (2010). Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature* **464**, 606. doi:10.1038/nature08899
- Jungke, P., Hans, S. and Brand, M. (2013). The zebrafish CreZoo: an easy-to-handle database for novel CreER(T2)-driver lines. *Zebrafish* **10**, 259-263. doi:10.1089/zeb.2012.0834
- Kamei, C. N., Liu, Y. and Drummond, I. A. (2015). Kidney regeneration in adult zebrafish by gentamicin induced injury. *J. Vis. Exp.* **102**, e51912. doi:10.3791/51912
- Kan, N. G., Junghans, D. and Izpisua Belmonte, J. C. (2009). Compensatory growth mechanisms regulated by BMP and FGF signaling mediate liver regeneration in zebrafish after partial hepatectomy. *FASEB J.* **23**, 3516-3525. doi:10.1096/fj.09-131730
- Kang, J., Hu, J., Karra, R., Dickson, A. L., Tornini, V. A., Nachtrab, G., Gemberling, M., Goldman, J. A., Black, B. L. and Poss, K. D. (2016). Modulation of tissue repair by regeneration enhancer elements. *Nature* **532**, 201-206. doi:10.1038/nature17644
- Kawakami, K. (2007). Tol2: a versatile gene transfer vector in vertebrates. *Genome Biol.* **8**, S7-S7. doi:10.1186/gb-2007-8-s1-s7
- Kawakami, K., Shima, A. and Kawakami, N. (2000). Identification of a functional transposase of the Tol2 element, an Ac-like element from the Japanese medaka fish, and its transposition in the zebrafish germ lineage. *Proc. Natl. Acad. Sci. USA* **97**, 11403-11408. doi:10.1073/pnas.97.21.11403
- Kawakami, K., Abe, G., Asada, T., Asakawa, K., Fukuda, R., Ito, A., Lal, P., Mouri, N., Muto, A., Suster, M. L. et al. (2010). zTrap: zebrafish gene trap and enhancer trap database. *BMC Dev. Biol.* **10**, 105. doi:10.1186/1471-213X-10-105
- Kikuchi, K., Holdway, J. E., Werdich, A. A., Anderson, R. M., Fang, Y., Egnaczyk, G. F., Evans, T., MacRae, C. A., Stainier, D. Y. R. and Poss, K. D. (2010). Primary contribution to zebrafish heart regeneration by gata4(+) cardiomyocytes. *Nature* **464**, 601-605. doi:10.1038/nature08804
- Kikuchi, K., Holdway, J. E., Major, R. J., Blum, N., Dahn, R. D., Begemann, G. and Poss, K. D. (2011). Retinoic acid production by endocardium and epicardium is an injury response essential for zebrafish heart regeneration. *Dev. Cell* **20**, 397-404. doi:10.1016/j.devcel.2011.01.010
- Kirchgeorg, L., Felker, A., van Oostrom, M., Chiavacci, E. and Mosimann, C. (2018). Cre/lox-controlled spatiotemporal perturbation of FGF signaling in zebrafish. *Dev. Dyn.* **247**, 1146-1159. doi:10.1002/dvdy.24668
- Kizil, C. and Brand, M. (2011). Cerebroventricular microinjection (CVMI) into adult zebrafish brain is an efficient misexpression method for forebrain ventricular cells. *PLoS ONE* **6**, e27395. doi:10.1371/journal.pone.0027395
- Koth, J., Maguire, M. L., McClymont, D., Diffley, L., Thornton, V. L., Beech, J., Patient, R. K., Riley, P. R. and Schneider, J. E. (2017). High-resolution magnetic resonance imaging of the regenerating adult zebrafish heart. *Sci. Rep.* **7**, 2917. doi:10.1038/s41598-017-03050-y
- Kroehne, V., Freudenreich, D., Hans, S., Kaslin, J. and Brand, M. (2011). Regeneration of the adult zebrafish brain from neurogenic radial glia-type progenitors. *Development* **138**, 4831. doi:10.1242/dev.072587
- Kyritsis, N., Kizil, C., Zocher, S., Kroehne, V., Kaslin, J., Freudenreich, D., Iltzsche, A. and Brand, M. (2012). Acute inflammation initiates the regenerative response in the adult zebrafish brain. *Science* **338**, 1353. doi:10.1126/science.1228773
- Lai, S.-L., Marín-Juez, R., Moura, P. L., Kuenen, C., Lai, J. K. H., Tseke, A. T., Guenther, S., Looso, M. and Stainier, D. Y. R. (2017). Reciprocal analyses in zebrafish and medaka reveal that harnessing the immune response promotes cardiac regeneration. *eLife* **6**, e25605. doi:10.7554/eLife.25605
- Langenau, D. M., Feng, H., Berghmans, S., Kanki, J. P., Kutok, J. L. and Look, A. T. (2005). Cre/lox-regulated transgenic zebrafish model with conditional myc-induced T cell acute lymphoblastic leukemia. *Proc. Natl. Acad. Sci. USA* **102**, 6068-6073. doi:10.1073/pnas.0408708102
- LeClair, E. E. and Topczewski, J. (2010). Development and regeneration of the zebrafish maxillary barbel: a novel study system for vertebrate tissue growth and repair. *PLoS ONE* **5**, e8737. doi:10.1371/journal.pone.0008737
- Lee, Y., Grill, S., Sanchez, A., Murphy-Ryan, M. and Poss, K. D. (2005). Fgf signaling instructs position-dependent growth rate during zebrafish fin regeneration. *Development* **132**, 5173. doi:10.1242/dev.02101
- Lepilina, A., Coon, A. N., Kikuchi, K., Holdway, J. E., Roberts, R. W., Burns, C. G. and Poss, K. D. (2006). A dynamic epicardial injury response supports progenitor cell activity during zebrafish heart regeneration. *Cell* **127**, 607-619. doi:10.1016/j.cell.2006.08.052
- Li, L., Yan, B., Shi, Y. Q., Zhang, W. Q. and Wen, Z. L. (2012). Live imaging reveals differing roles of macrophages and neutrophils during zebrafish tail fin regeneration. *J. Biol. Chem.* **287**, 25353-25360. doi:10.1074/jbc.M112.349126
- Lim, D. A. and Alvarez-Buylla, A. (2016). The adult ventricular-subventricular zone (V-SVZ) and olfactory bulb (OB) neurogenesis. *Cold Spring Harbor Perspect. Biol.* **8**, a018820. doi:10.1101/cshperspect.a018820
- Liu, K., Petree, C., Requena, T., Varshney, P. and Varshney, G. K. (2019). Expanding the CRISPR toolbox in zebrafish for studying development and disease. *Front. Cell Dev. Biol.* **7**, 13. doi:10.3389/fcell.2019.00013
- Liu, Q., Azodi, E., Kerstetter, A. E. and Wilson, A. L. (2004). Cadherin-2 and cadherin-4 in developing, adult and regenerating zebrafish cerebellum. *Brain Res. Dev. Brain Res.* **150**, 63-71. doi:10.1016/j.devbrainres.2004.03.002
- Long, L., Guo, H., Yao, D., Xiong, K., Li, Y., Liu, P., Zhu, Z. and Liu, D. (2015). Regulation of transcriptionally active genes via the catalytically inactive Cas9 in *C. elegans* and *D. rerio*. *Cell Res.* **25**, 638-641. doi:10.1038/cr.2015.35
- Lu, J., Liu, K.-C., Schulz, N., Karampelias, C., Charbord, J., Hilding, A., Rautio, L., Bertolino, P., Östenson, C.-G., Brismar, K. et al. (2016). IGFBP1 increases beta-cell regeneration by promoting alpha-to beta-cell transdifferentiation. *EMBO J.* **35**, 2026-2044. doi:10.15252/embj.201592903
- Mahmoud, A. I., O'Meara, C. C., Gemberling, M., Zhao, L., Bryant, D. M., Zheng, R., Gannon, J. B., Cai, L., Choi, W. Y., Egnaczyk, G. F. et al. (2015). Nerves regulate cardiomyocyte proliferation and heart regeneration. *Dev. Cell* **34**, 387-399. doi:10.1016/j.devcel.2015.06.017
- Marín-Juez, R., Marass, M., Gauvrit, S., Rossi, A., Lai, S.-L., Materna, S. C., Black, B. L. and Stainier, D. Y. R. (2016). Fast revascularization of the injured

- area is essential to support zebrafish heart regeneration. *Proc. Natl. Acad. Sci. USA* **113**, 11237-11242. doi:10.1073/pnas.1605431113
- März, M., Schmidt, R., Rastegar, S. and Strähle, U. (2011). Regenerative response following stab injury in the adult zebrafish telencephalon. *Dev. Dyn.* **240**, 2221-2231. doi:10.1002/dvdy.22710
- Mateus, R., Lourenco, R., Fang, Y., Brito, G., Farinho, A., Valerio, F. and Jacinto, A. (2015). Control of tissue growth by Yap relies on cell density and F-actin in zebrafish fin regeneration. *Development* **142**, 2752-2763. doi:10.1242/dev.119701
- Mathew, L. K., Sengupta, S., Kawakami, A., Andreassen, E. A., Löhr, C. V., Loynes, C. A., Renshaw, S. A., Peterson, R. T. and Tanguay, R. L. (2007). Unraveling tissue regeneration pathways using chemical genetics. *J. Biol. Chem.* **282**, 35202-35210. doi:10.1074/jbc.M706640200
- Matsuda, H., Mullapudi, S. T., Carol Yang, Y. H., Masaki, H., Hesselson, D. and Stainier, D. Y. R. (2018). Whole organism chemical screening identifies modulators of pancreatic beta cell function. *Diabetes* **67**, 2268-2279. doi:10.2337/db17-1223
- Mescher, A. L. (2017). Macrophages and fibroblasts during inflammation and tissue repair in models of organ regeneration. *Regeneration* **4**, 39-53. doi:10.1002/reg.2.77
- Michalopoulos, G. K. (2007). Liver regeneration. *J. Cell. Physiol.* **213**, 286-300. doi:10.1002/jcp.21172
- Mokalled, M. H., Patra, C., Dickson, A. L., Endo, T., Stainier, D. Y. R. and Poss, K. D. (2016). Injury-induced ctgfa directs glial bridging and spinal cord regeneration in zebrafish. *Science* **354**, 630-634. doi:10.1126/science.aaf2679
- Moore, F. E., Reyon, D., Sander, J. D., Martinez, S. A., Blackburn, J. S., Khayter, C., Ramirez, C. L., Joung, J. K. and Langenau, D. M. (2012). Improved somatic mutagenesis in zebrafish using transcription activator-like effector nucleases (TALENs). *PLoS ONE* **7**, e37877. doi:10.1371/journal.pone.0037877
- Mosimann, C., Puller, A. C., Lawson, K. L., Tschopp, P., Amsterdam, A. and Zon, L. I. (2013). Site-directed zebrafish transgenesis into single landing sites with the phiC31 integrase system. *Dev. Dyn.* **242**, 949-963. doi:10.1002/dvdy.23989
- Münch, J., González-Rajal, A. and de la Pompa, J. L. (2013). Notch regulates blastema proliferation and prevents differentiation during adult zebrafish fin regeneration. *Development* **140**, 1402. doi:10.1242/dev.087346
- Nauroy, P., Guiraud, A., Chlasta, J., Malbouyres, M., Gillet, B., Hughes, S., Lambert, E. and Ruggiero, F. (2019). Gene profile of zebrafish fin regeneration offers clues to kinetics, organization and biomechanics of basement membrane. *Matrix Biol.* **75-76**, 82-101. doi:10.1016/j.matbio.2018.07.005
- Nechiporuk, A. and Keating, M. T. (2002). A proliferation gradient between proximal and msxb-expressing distal blastema directs zebrafish fin regeneration. *Development* **129**, 2607.
- Nemoto, Y., Higuchi, K., Baba, O., Kudo, A. and Takano, Y. (2007). Multinucleate osteoclasts in medaka as evidence of active bone remodeling. *Bone* **40**, 399-408. doi:10.1016/j.bone.2006.08.019
- Ohnmacht, J., Yang, Y., Maurer, G. W., Barreiro-Iglesias, A., Tsarouchas, T. M., Wehner, D., Sieger, D., Becker, C. G. and Becker, T. (2016). Spinal motor neurons are regenerated after mechanical lesion and genetic ablation in larval zebrafish. *Development* **143**, 1464. doi:10.1242/dev.129155
- Otsuka, T. and Takeda, H. (2017). Targeted ablation of pancreatic beta cells in medaka. *Zool. Sci.* **34**, 179-184. doi:10.2108/zs170004
- Pandey, G., Westhoff, J. H., Schaefer, F. and Gehrig, J. (2019). A smart imaging workflow for organ-specific screening in a cystic kidney zebrafish disease model. *Int. J. Mol. Sci.* **20**, E1290. doi:10.3390/ijms20061290
- Parrilla, M., Lillo, C., Herrero-Turrión, M. J., Arévalo, R., Aijón, J., Lara, J. M. and Velasco, A. (2012). Characterization of Pax2 expression in the goldfish optic nerve head during retina regeneration. *PLoS ONE* **7**, e32348. doi:10.1371/journal.pone.0032348
- Patterson, M., Barske, L., Van Handel, B., Rau, C. D., Gan, P., Sharma, A., Parikh, S., Denholtz, M., Huang, Y., Yamaguchi, Y. et al. (2017). Frequency of mononuclear diploid cardiomyocytes underlies natural variation in heart regeneration. *Nat. Genet.* **49**, 1346-1353. doi:10.1038/ng.3929
- Petrie, T. A., Strand, N. S., Tsung-Yang, C., Rabinowitz, J. S. and Moon, R. T. (2014). Macrophages modulate adult zebrafish tail fin regeneration. *Development* **141**, 2581-2591. doi:10.1242/dev.098459
- Pfefferli, C. and Jaźwińska, A. (2015). The art of fin regeneration in zebrafish. *Regeneration* **2**, 72-83. doi:10.1002/reg.2.33
- Pfefferli, C. and Jaźwińska, A. (2017). The careg element reveals a common regulation of regeneration in the zebrafish myocardium and fin. *Nat. Commun.* **8**, 15151. doi:10.1038/ncomms15151
- Pinto-Teixeira, F., Viader-Llagues, O., Torres-Mejía, E., Turan, M., Gonzalez-Gualda, E., Pola-Morell, L. and Lopez-Schier, H. (2015). Inexhaustible hair-cell regeneration in young and aged zebrafish. *Biol. Open* **4**, 903-909. doi:10.1242/bio.012112
- Pisharath, H., Rhee, J. M., Swanson, M. A., Leach, S. D. and Parsons, M. J. (2007). Targeted ablation of beta cells in the embryonic zebrafish pancreas using E. coli nitroreductase. *Mech. Dev.* **124**, 218-229. doi:10.1016/j.mod.2006.11.005
- Porrello, E. R., Mahmoud, A. I., Simpson, E., Hill, J. A., Richardson, J. A., Olson, E. N. and Sadek, H. A. (2011). Transient regenerative potential of the neonatal mouse heart. *Science* **331**, 1078. doi:10.1126/science.1200708
- Poss, K. D., Shen, J., Nechiporuk, A., McMahon, G., Thisse, B., Thisse, C. and Keating, M. T. (2000). Roles for Fgf signaling during zebrafish fin regeneration. *Dev. Biol.* **222**, 347-358. doi:10.1006/dbio.2000.9722
- Poss, K. D., Wilson, L. G. and Keating, M. T. (2002). Heart regeneration in zebrafish. *Science* **298**, 2188. doi:10.1126/science.1077857
- Power, R. M. and Huisken, J. (2017). A guide to light-sheet fluorescence microscopy for multiscale imaging. *Nat. Methods* **14**, 360-373. doi:10.1038/nmeth.4224
- Rabinowitz, J. S., Robitaille, A. M., Wang, Y., Ray, C. A., Thummel, R., Gu, H., Djukovic, D., Rafferty, D., Berndt, J. D. and Moon, R. T. (2017). Transcriptomic, proteomic, and metabolomic landscape of positional memory in the caudal fin of zebrafish. *Proc. Natl. Acad. Sci. USA* **114**, E717. doi:10.1073/pnas.1620755114
- Rafferty, S. A. and Quinn, T. A. (2018). A beginner's guide to understanding and implementing the genetic modification of zebrafish. *Prog. Biophys. Mol. Biol.* **138**, 3-19. doi:10.1016/j.pbiomolbio.2018.07.005
- Raya, A., Koth, C. M., Buscher, D., Kawakami, Y., Itoh, T., Raya, R. M., Sternik, G., Tsai, H.-J., Rodriguez-Esteban, C. and Izpisua-Belmonte, J. C. (2003). Activation of Notch signaling pathway precedes heart regeneration in zebrafish. *Proc. Natl. Acad. Sci. USA* **100** Suppl. 1, 11889-11895. doi:10.1073/pnas.1834204100
- Raymond, P. A., Barthel, L. K., Bernardos, R. L. and Perkowski, J. J. (2006). Molecular characterization of retinal stem cells and their niches in adult zebrafish. *BMC Dev. Biol.* **6**, 36. doi:10.1186/1471-213X-6-36
- Reimschuessel, R. (2001). A fish model of renal regeneration and development. *ILAR J.* **42**, 285-291. doi:10.1093/ilar.42.4.285
- Reimschuessel, R. and Williams, D. (1995). Development of new nephrons in adult kidneys following gentamicin-induced nephrotoxicity. *Ren. Fail.* **17**, 101-106. doi:10.3109/08860229509026246
- Richardson, R., Slanchev, K., Kraus, C., Knyphausen, P., Eming, S. and Hammerschmidt, M. (2013). Adult zebrafish as a model system for cutaneous wound-healing research. *J. Invest. Dermatol.* **133**, 1655-1665. doi:10.1038/jid.2013.16
- Rothenaigner, I., Krecsmarik, M., Hayes, J. A., Bahn, B., Lepier, A., Fortin, G., Götz, M., Jagasia, R. and Bally-Cuif, L. (2011). Clonal analysis by distinct viral vectors identifies bona fide neural stem cells in the adult zebrafish telencephalon and characterizes their division properties and fate. *Development* **138**, 1459. doi:10.1242/dev.058156
- Rowlerson, A., Radaelli, G., Mascarello, F. and Veggetti, A. (1997). Regeneration of skeletal muscle in two teleost fish: Sparus aurata and Brachydanio rerio. *Cell Tissue Res.* **289**, 311-322. doi:10.1007/s004410050878
- Sadler, K. C., Krahn, K. N., Gaur, N. A. and Ukomadu, C. (2007). Liver growth in the embryo and during liver regeneration in zebrafish requires the cell cycle regulator, uhrf1. *Proc. Natl. Acad. Sci. USA* **104**, 1570-1575. doi:10.1073/pnas.0610774104
- Sánchez-Iranzo, H., Galardi-Castilla, M., Sanz-Morejón, A., González-Rosa, J. M., Costa, R., Ernst, A., Sainz de Aja, J., Langa, X. and Mercader, N. (2018). Transient fibrosis resolves via fibroblast inactivation in the regenerating zebrafish heart. *Proc. Natl. Acad. Sci. USA* **115**, 4188. doi:10.1073/pnas.1716713115
- Saxena, S., Singh, S. K., Lakshmi, M. G. M., Meghah, V., Bhatti, B., Swamy, C. V. B., Sundaram, C. S. and Idris, M. M. (2012). Proteomic analysis of zebrafish caudal fin regeneration. *Mol. Cell. Proteomics* **11**, M1111.014118. doi:10.1074/mcp.M111.014118
- Schnabel, K., Wu, C.-C., Kurth, T. and Weidinger, G. (2011). Regeneration of cryoinjury induced necrotic heart lesions in zebrafish is associated with epicardial activation and cardiomyocyte proliferation. *PLoS ONE* **6**, e18503. doi:10.1371/journal.pone.0018503
- Sehring, I. M., Jahn, C. and Weidinger, G. (2016). Zebrafish fin and heart: what's special about regeneration? *Curr. Opin. Genet. Dev.* **40**, 48-56. doi:10.1016/j.gde.2016.05.011
- Senyo, S. E., Steinhauser, M. L., Pizzimenti, C. L., Yang, V. K., Cai, L., Wang, M., Wu, T.-D., Guerquin-Kern, J.-L., Lechene, C. P. and Lee, R. T. (2013). Mammalian heart renewal by pre-existing cardiomyocytes. *Nature* **493**, 433-436. doi:10.1038/nature11682
- Simões, M. G., Bensimon-Brito, A., Fonseca, M., Farinho, A., Valério, F., Sousa, S., Afonso, N., Kumar, A. and Jacinto, A. (2014). Denervation impairs regeneration of amputated zebrafish fins. *BMC Dev. Biol.* **14**, 49. doi:10.1186/s12861-014-0049-2
- Simon, A. and Tanaka, E. M. (2013). Limb regeneration. *Wiley Interdiscip. Rev. Dev. Biol.* **2**, 291-300. doi:10.1002/wdev.73
- Singh, S. P., Holdway, J. E. and Poss, K. D. (2012). Regeneration of amputated zebrafish fin rays from de novo osteoblasts. *Dev. Cell* **22**, 879-886. doi:10.1016/j.devcel.2012.03.006
- Slack, J. M. W. (2017). Animal regeneration: ancestral character or evolutionary novelty? *EMBO Rep.* **18**, 1497-1508. doi:10.15252/embr.201643795
- Smith, A., Avaron, F., Guay, D., Padhi, B. K. and Akimenko, M. A. (2006). Inhibition of BMP signaling during zebrafish fin regeneration disrupts fin growth and scleroblast differentiation and function. *Dev. Biol.* **299**, 438-454. doi:10.1016/j.ydbio.2006.08.016
- Spanjaard, B., Hu, B., Mitic, N., Olivares-Chauvet, P., Janjuha, S., Ninov, N. and Junker, J. P. (2018). Simultaneous lineage tracing and cell-type identification

- using CRISPR-Cas9-induced genetic scars. *Nat. Biotechnol.* **36**, 469-473. doi:10.1038/nbt.4124
- Stewart, S. and Stankunas, K.** (2012). Limited dedifferentiation provides replacement tissue during zebrafish fin regeneration. *Dev. Biol.* **365**, 339-349. doi:10.1016/j.ydbio.2012.02.031
- Stewart, S., Tsun, Z.-Y. and Izpissua Belmonte, J. C. I.** (2009). A histone demethylase is necessary for regeneration in zebrafish. *Proc. Natl. Acad. Sci. USA* **106**, 19889-19894. doi:10.1073/pnas.0904132106
- Stockdale, W. T., Lemieux, M. E., Killen, A. C., Zhao, J., Hu, Z., Riepsaame, J., Hamilton, N., Kudoh, T., Riley, P. R., van Aerle, R. et al.** (2018). Heart regeneration in the mexican cavefish. *Cell Rep.* **25**, 1997-2007. doi:10.1016/j.celrep.2018.10.072
- Stoick-Cooper, C. L., Weidinger, G., Riehle, K. J., Hubbert, C., Major, M. B., Fausto, N. and Moon, R. T.** (2007). Distinct Wnt signaling pathways have opposing roles in appendage regeneration. *Development* **134**, 479-489. doi:10.1242/dev.001123
- Strahle, U. and Grabher, C.** (2010). The zebrafish embryo as a model for assessing off-target drug effects. *Dis. Model. Mech.* **3**, 689-692. doi:10.1242/dmm.006312
- Strahle, U., Bally-Cuif, L., Kelsh, R., Beis, D., Mione, M., Panula, P., Figueras, A., Gothif, Y., Brösamle, C., Geisler, R. et al.** (2012). EuFishBioMed (COST Action BM0804): a European network to promote the use of small fishes in biomedical research. *Zebrafish* **9**, 90-93. doi:10.1089/zeb.2012.0742
- Streisinger, G., Walker, C., Dower, N., Knauber, D. and Singer, F.** (1981). Production of clones of homozygous diploid zebra fish (*Brachydanio rerio*). *Nature* **291**, 293-296. doi:10.1038/291293a0
- Stuart, G. W., McMurray, J. V. and Westerfield, M.** (1988). Replication, integration and stable germ-line transmission of foreign sequences injected into early zebrafish embryos. *Development* **103**, 403-412.
- Sugiyama, M., Sakaue-Sawano, A., Iimura, T., Fukami, K., Kitaguchi, T., Kawakami, K., Okamoto, H., Higashijima, S. and Miyawaki, A.** (2009). Illuminating cell-cycle progression in the developing zebrafish embryo. *Proc. Natl. Acad. Sci. USA* **106**, 20812-20817. doi:10.1073/pnas.0906464106
- Suster, M. L., Abe, G., Schouw, A. and Kawakami, K.** (2011). Transposon-mediated BAC transgenesis in zebrafish. *Nat. Protoc.* **6**, 1998-2021. doi:10.1038/nprot.2011.416
- Thermes, V., Grabher, C., Ristoratore, F., Bourrat, F., Choulika, A., Wittbrodt, J. and Joly, J.-S.** (2002). I-SceI meganuclease mediates highly efficient transgenesis in fish. *Mech. Dev.* **118**, 91-98. doi:10.1016/S0925-4773(02)00218-6
- Thummel, R., Bai, S., Sarras, M. P., Jr, Song, P., McDermott, J., Brewer, J., Perry, M., Zhang, X., Hyde, D. R. and Godwin, A. R.** (2006). Inhibition of zebrafish fin regeneration using in vivo electroporation of morpholinos against fgfr1 and msxb. *Dev. Dyn.* **235**, 336-346. doi:10.1002/dvdy.20630
- Tsarouchas, T. M., Wehner, D., Cavone, L., Munir, T., Keatinge, M., Lambertus, M., Underhill, A., Barrett, T., Kassapis, E., Ogryzko, N. et al.** (2018). Dynamic control of proinflammatory cytokines Il-1beta and Tnf-alpha by macrophages in zebrafish spinal cord regeneration. *Nat. Commun.* **9**, 4670. doi:10.1038/s41467-018-07036-w
- Tsuji, N., Ninov, N., Delawary, M., Osman, S., Roh, A. S., Gut, P. and Stainier, D. Y. R.** (2014). Whole organism high content screening identifies stimulators of pancreatic beta-cell proliferation. *PLoS ONE* **9**, e104112. doi:10.1371/journal.pone.0104112
- Unal Eroglu, A., Mulligan, T. S., Zhang, L., White, D. T., Sengupta, S., Nie, C., Lu, N. Y., Qian, J., Xu, L., Pei, W. et al.** (2018). Multiplexed CRISPR/Cas9 targeting of genes implicated in retinal regeneration and degeneration. *Front. Cell Dev. Biol.* **6**, 88. doi:10.3389/fcell.2018.00088
- Vihelic, T. S. and Hyde, D. R.** (2000). Light-induced rod and cone cell death and regeneration in the adult albino zebrafish (*Danio rerio*) retina. *J. Neurobiol.* **44**, 289-307. doi:10.1002/1097-4695(20000905)44:3<289::AID-NEU1>3.0.CO;2-H
- Wang, J., Panáková, D., Kikuchi, K., Holdway, J. E., Gemberling, M., Burris, J. S., Singh, S. P., Dickson, A. L., Lin, Y.-F., Sabeh, M. K. et al.** (2011). The regenerative capacity of zebrafish reverses cardiac failure caused by genetic cardiomyocyte depletion. *Development* **138**, 3421. doi:10.1242/dev.068601
- Watanabe, N., Kato, M., Suzuki, N., Inoue, C., Fedorova, S., Hashimoto, H., Maruyama, S., Matsuo, S. and Wakamatsu, Y.** (2009). Kidney regeneration through nephron neogenesis in medaka. *Dev. Growth Differ.* **51**, 135-143. doi:10.1111/j.1440-169X.2009.01090.x
- Wehner, D., Cizelsky, W., Vasudevaro, M. D., Özhan, G., Haase, C., Kagermeier-Schenk, B., Röder, A., Dorsky, R. I., Moro, E., Argenton, F. et al.** (2014). Wnt/beta-catenin signaling defines organizing centers that orchestrate growth and differentiation of the regenerating zebrafish caudal fin. *Cell Rep.* **6**, 467-481. doi:10.1016/j.celrep.2013.12.036
- Wehner, D., Jahn, C. and Weidinger, G.** (2015). Use of the TetON system to study molecular mechanisms of zebrafish regeneration. *J. Vis. Exp.* **100**, e52756. doi:10.3791/52756
- Wehner, D., Tsarouchas, T. M., Michael, A., Haase, C., Weidinger, G., Reimer, M. M., Becker, T. and Becker, C. G.** (2017). Wnt signaling controls pro-regenerative Collagen XII in functional spinal cord regeneration in zebrafish. *Nat. Commun.* **8**, 126. doi:10.1038/s41467-017-00143-0
- White, J. A., Boffa, M. B., Jones, B. and Petkovich, M.** (1994). A zebrafish retinoic acid receptor expressed in the regenerating caudal fin. *Development* **120**, 1861.
- Wills, A. A., Holdway, J. E., Major, R. J. and Poss, K. D.** (2008). Regulated addition of new myocardial and epicardial cells fosters homeostatic cardiac growth and maintenance in adult zebrafish. *Development* **135**, 183-192. doi:10.1242/dev.010363
- Wosczyzna, M. N. and Rando, T. A.** (2018). A muscle stem cell support group: coordinated cellular responses in muscle regeneration. *Dev. Cell* **46**, 135-143. doi:10.1016/j.devcel.2018.06.018
- Wu, C.-C., Kruse, F., Vasudevarao, M. D., Junker, J. P., Zebrowski, D. C., Fischer, K., Noël, E. S., Grün, D., Berezikov, E., Engel, F. B. et al.** (2016). Spatially resolved genome-wide transcriptional profiling identifies BMP signaling as essential regulator of zebrafish cardiomyocyte regeneration. *Dev. Cell* **36**, 36-49. doi:10.1016/j.devcel.2015.12.010
- Xiao, C., Gao, L., Hou, Y., Xu, C., Chang, N., Wang, F., Hu, K., He, A., Luo, Y., Wang, J. et al.** (2016). Chromatin-remodelling factor Brg1 regulates myocardial proliferation and regeneration in zebrafish. *Nat. Commun.* **7**, 13787. doi:10.1038/ncomms13787
- Xu, C., Volkery, S. and Siekmann, A. F.** (2015). Intubation-based anesthesia for long-term time-lapse imaging of adult zebrafish. *Nat. Protoc.* **10**, 2064-2073. doi:10.1038/nprot.2015.130
- Yang, K. and Kang, J.** (2018). Tissue regeneration enhancer elements: a way to unlock endogenous healing power. *Dev. Dyn.* **248**, 34-42. doi:10.1002/dvdy.24676
- Young, B. H. and Heath, J. W.** (2000). *Wheater's Functional Histology: A Text and Colour Atlas*, 4th edn. Churchill Livingstone.
- Zhao, L., Borikova, A. L., Ben-Yair, R., Guner-Ataman, B., MacRae, C. A., Lee, R. T., Burns, C. G. and Burns, C. E.** (2014). Notch signaling regulates cardiomyocyte proliferation during zebrafish heart regeneration. *Proc. Natl. Acad. Sci. USA* **111**, 1403. doi:10.1073/pnas.1311705111