

Complete Genomic Sequence of European Bat Lyssavirus 1, Isolated from *Eptesicus isabellinus* in Spain

Denise A. Marston,^a Sonia Vázquez-Morón,^{b,9} Richard J. Ellis,^c Emma L. Wise,^a Lorraine M. McElhinney,^{a,d} Xavier de Lamballerie,^e Anthony R. Fooks,^{a,d,f} Juan E. Echevarría^{b,9}

Wildlife Zoonoses & Vector-Borne Diseases Research Group,^a Specialist Scientific Support Department,^c Animal and Plant Health Agency (APHA), Surrey, United Kingdom; Centro Nacional de Microbiología, Instituto de Salud Carlos III, Majadahonda, Madrid, Spain^b; National Consortium for Zoonosis Research, University of Liverpool, Liverpool, United Kingdom^d; UMR_D 190, "Emergence de Pathologies Virales," Aix Marseille University, Marseille, France^e; Department of Clinical Infection, Microbiology and Immunology, University of Liverpool, Liverpool, United Kingdom^f; Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain⁹

All members of the lyssavirus genus cause the disease rabies. European bat lyssavirus 1 (EBLV-1) viruses are divided genetically into three groups according to geographic location and host reservoir. We report here the first genome sequence for an EBLV-1 isolated from *Eptesicus isabellinus* in the Iberian Peninsula, Spain.

Received 15 December 2014 Accepted 5 January 2015 Published 12 February 2015

Citation Marston DA, Vázquez-Morón S, Ellis RJ, Wise EL, McElhinney LM, de Lamballerie X, Fooks AR, Echevarría JE. 2015. Complete genomic sequence of European bat lyssavirus 1, isolated from *Eptesicus isabellinus* in Spain. *Genome Announc* 3(1):e01518-14. doi:10.1128/genomeA.01518-14.

Copyright © 2015 Marston et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Denise A. Marston, denise.marston@apha.gsi.gov.uk.

There are 14 classified lyssavirus species. In Europe, bat rabies cases are dominated by two lyssavirus species: European bat lyssavirus 1 (EBLV-1) and European bat lyssavirus 2 (EBLV-2) (1). EBLV-1 molecular epidemiological investigations separated the species into 2 sublineages (1a and 1b) (2). Analysis of EBLV-1 viruses from *Eptesicus isabellinus* in the Iberian Peninsula has demonstrated that these viruses are genetically similar, yet distinct from other EBLV-1 circulating in *Eptesicus serotinus* (3). Full-genome sequences are available for five EBLV-1a and a single EBLV-1b virus. Here, we describe full-genome sequencing of an EBLV-1 virus (292R07, RV2416), isolated from an infected *E. isabellinus* bat from the Iberian peninsula (Granada), Spain, in 2007.

RNA from mouse brain homogenate (passage 1) was prepared for next-generation sequencing on the MiSeq platform. Briefly, TRIzol extracted viral RNA was depleted of host genomic DNA and rRNA as described previously (4). Double-stranded (ds) cDNA was synthesized from 50 ng RNA by use of a random cDNA synthesis system (Roche), according to the manufacturer's instructions. The ds cDNA was purified using Ampure XP magnetic beads (Beckman Coulter), and 1 ng was used for the Nextera XT DNA sample preparation kit (Illumina). A sequencing library was prepared according to the manufacturers' instructions and sequenced on an Illumina MiSeq with 2 × 150-bp paired-end reads following standard Illumina protocols. The total reads (2,032,092) were mapped to a reference sequence (EU293112) using BWA (v0.7.5a-r405) (5) and was visualized in Tablet (6). A modified samtools/vcfutils (7) script was used to generate an intermediate consensus sequence in which any indels relative to the original reference sequence were appropriately called. The intermediate consensus was used as the reference for subsequent iteration of mapping and consensus calling. The total number of assembled viral reads was 5,600 (0.27% of the total reads). Despite the low proportion of viral sequence detected within the total data set, adequate coverage of the entire genome was obtained (average read depth of 58.16), with the exception of the highly conserved com-

plementary untranslated regions (UTRs). The genomic termini sequence was obtained by using primers described previously (4).

The genetic organization of the *E. isabellinus* EBLV-1 genome was similar to those of other EBLV-1 genomes, with a complete genome size of 11,964 nucleotides. The coding region lengths were conserved; however, indels were observed in a number of intergenic regions. A 6-bp indel in the N-P intergenic region (1478 to 1483), which was previously identified within the EBLV-1 species (8), was observed. Interestingly, only the EBLV-1b full genome from *E. serotinus* from France had the 6-bp insertion; all other EBLV-1 genomes, including the *E. isabellinus* EBLV-1, did not. Two further indels were identified, (i) A²⁴²⁰ in the P-M region and (ii) C⁵²⁴⁹ in the G-L region. Both nucleotides were absent in the *E. isabellinus* EBLV-1 in comparison to the six EBLV-1 genomes available. It is unlikely that the observed indels have any biological influence; however, they support the independent evolution of the Iberian peninsula EBLV-1 viruses to other EBLV-1 viruses, due to geographical and host separation.

Nucleotide sequence accession number. The complete genomic sequence of RV2416 (292R07) has been deposited in GenBank under the accession number [KP241939](https://www.ncbi.nlm.nih.gov/nuclseq/KP241939).

ACKNOWLEDGMENTS

This work was supported by grants from the Department for Environment, Food and Rural Affairs (SE0423 and SE0427) and by the European Union FP7-funded Research Infrastructure Grant European Virus Archive (19 228292).

REFERENCES

- McElhinney LM, Marston DA, Leech S, Freuling CM, van der Poel WH, Echevarría J, Vázquez-Morón S, Horton DL, Müller T, Fooks AR. 2013. Molecular epidemiology of bat lyssaviruses in Europe. *Zoonoses Public Health* 60:35–45. <http://dx.doi.org/10.1111/zph.12003>.
- Davis PL, Holmes EC, Larrous F, Van der Poel WH, Tjørnehøj K, Alonso WJ, Bourhy H. 2005. Phylogeography, population dynamics, and molecular evolution of European bat lyssaviruses. *J Virol* 79:10487–10497. <http://dx.doi.org/10.1128/JVI.79.16.10487-10497.2005>.

3. Vázquez-Moron S, Juste J, Ibáñez C, Berciano JM, Echevarria JE. 2011. Phylogeny of European bat lyssavirus 1 in *Eptesicus isabellinus* bats, Spain. *Emerg Infect Dis* 17:520–523. <http://dx.doi.org/10.3201/eid1703.100894>.
4. Marston DA, McElhinney LM, Ellis RJ, Horton DL, Wise EL, Leech SL, David D, de Lamballerie X, Fooks AR. 2013. Next generation sequencing of viral RNA genomes. *BMC Genomics* 14:444. <http://dx.doi.org/10.1186/1471-2164-14-444>.
5. Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <http://dx.doi.org/10.1093/bioinformatics/btp698>.
6. Milne I, Stephen G, Bayer M, Cock PJ, Pritchard L, Cardle L, Shaw PD, Marshall D. 2013. Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform* 14:193–202. <http://dx.doi.org/10.1093/bib/bbs012>.
7. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079. <http://dx.doi.org/10.1093/bioinformatics/btp352>.
8. Johnson N, Freuling C, Marston DA, Tordo N, Fooks AR, Müller T. 2007. Identification of European bat lyssavirus isolates with short genomic insertions. *Virus Res* 128:140–143. <http://dx.doi.org/10.1016/j.virusres.2007.04.012>.