








Antibodies against the *Mycobacterium tuberculosis* complex and *Brucella* spp. in captive and free-living European bison (*Bison bonasus*) in Poland

Anna Didkowska¹  | Elisa Ferreras Colino²  | Wanda Olech³  |
 Huguette Gloddy^{2,4}  | Krzysztof Anusz¹  | José Antonio Infantes-Lorenzo⁵  |
 Christian Gortázar² 

¹Department of Food Hygiene and Public Health Protection, Institute of Veterinary Medicine, Warsaw University of Life Sciences (SGGW), Warsaw, Poland

²SaBio Instituto de Investigación en Recursos Cinegéticos IREC, Consejo Superior de Investigaciones Científicas, Universidad de Castilla – La Mancha, Ciudad Real, Spain

³Department of Animal Genetics and Conservation, Institute of Animal Sciences, University of Life Sciences (SGGW), Warsaw, Poland

⁴Centre de Recherche en Sciences Naturelles de Lwiro (CRSN Lwiro), Lwiro, South Kivu, Democratic Republic of Congo

⁵Unidad de Inmunología Microbiana, Centro Nacional de Microbiología, Instituto de Salud Carlos III, Majadahonda, Madrid, Spain

Correspondence

Anna Didkowska, Department of Food Hygiene and Public Health Protection, Institute of Veterinary Medicine, Warsaw University of Life Sciences (SGGW), Nowoursynowska 159, 02-776 Warsaw, Poland.
 Email: anna_didkowska@sggw.edu.pl

Funding information

Complex project of European bison conservation by State Forests, Forest Fund (Poland), Grant/Award Number: contract no OR.271.3.10.2017

Abstract

Background: The European bison (*Bison bonasus*), a symbol of Polish nature, is a protected species that requires active health monitoring. However, conservation efforts are made difficult by the zoonotic diseases such as brucellosis and tuberculosis.

Objective: The aim of this study was to screen the Polish European bison population for exposure to the *Mycobacterium tuberculosis* complex (MTC) and *Brucella* spp.

Methods: A total of 323 free-living and captive European bison from 13 localities were tested serologically for antibodies against the *M. bovis* P22 multi-protein complex (in-house ELISA) and against *Brucella* spp. (commercial ELISA).

Results: Antibodies against the MTC (P22) were detected in 7% (22/323) of the tested European bison. Anti-MTC antibody positivity was not significantly different by sex, age, and captive/free range status. Anti-MTC antibodies were found in six of 13 populations sampled, always in populations with larger sample sizes including the four free-living ones. Antibodies against *Brucella* spp. were detected in 36% (116/323) of the tested bison. While *Brucella* spp. antibody prevalence was not different by sex, it was significantly different by age (lower in adults) and captive/free-living status. *Brucella* spp. seroprevalence decreased with sample size and seropositive bison were found in 12 of 13 sampling populations.

Conclusions: Our findings identify potential emerging threats to the European bison population and confirm the first serological response to P22 in European bison. As Poland is currently officially free of brucellosis and bovine tuberculosis, our results require careful interpretation. Further studies are needed to establish the presence of cross-reactions with atypical mycobacteria in the case of MTC and other bacteria (e.g. *Yersinia enterocolitica* O:9) in the case of *Brucella* spp.

KEYWORDS

Brucella spp, ELISA, European bison, P22, serology, tuberculosis

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Veterinary Medicine and Science* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

The European bison (*Bison bonasus*) is an iconic Polish wildlife species, with more than a quarter of the global European bison population living in Poland. The species is currently safe from extinction, having recently changed its conservation status to 'Near Threatened', according to the International Union for Conservation of Nature (IUCN) (Plumb et al., 2020).

The number of European bison in Poland at the end of 2021 was 206 in captivity (most herds are of a low number) and 2223 in free-living populations located in eight different herds (Raczynski & Bołbot, 2022). The European bison in Poland is undergoing dynamic population development (Olech & Perzanowski, 2022) that results in high local population densities; therefore, it is important to monitor infectious diseases and emerging threats (Klich et al., 2020, 2021).

Tuberculosis (TB), caused by members of the *Mycobacterium tuberculosis* complex (MTC), and brucellosis, caused by the genus *Brucella*, not only have a considerable impact on cattle herds worldwide but also are bacterial zoonoses with significant potential impact on public health. Although Poland is officially free of TB and brucellosis in cattle (Regulation [(EU) 2021/620]) both are occasionally noted in Polish wildlife (Szulowski et al., 2013a; Krajewska et al., 2014). It has been suggested that European bison are highly sensitive to *Mycobacterium* spp. infections (Didkowska, Orłowska, et al., 2021); indeed, TB has been confirmed in both captive (Didkowska, Orłowska, et al., 2021) and free-living herds of European bison in Poland (Welz et al., 2005). Although paratuberculosis does not seem to be an emerging problem in Polish European bison herds (Didkowska, Ptak et al., 2021), atypical mycobacteria (*Mycobacterium avium* and *Mycobacterium xenopi*) have been isolated from them (Didkowska, Orłowska, et al., 2021).

TB has also been identified in European bison in a Brazilian zoo (Zimpel et al., 2017) and is endemic in American bison (*Bison bison*) in some North American populations (Himsworth et al., 2010; Nishi et al., 2002; Shury et al., 2015; Tessaro et al., 1990). In recent years, due to the emerging cases of TB in bison, diagnostic procedures have been extended to incorporate new methods, including serological tests (Didkowska, Dziekan, et al., 2021).

One antigen that has recently been studied for use in serological tests in various mammalian species is P22 (Bezoz et al., 2018; Casal et al., 2017; Infantes-Lorenzo et al., 2018; Infantes-Lorenzo, Dave et al., 2019; Infantes-Lorenzo, Moreno et al., 2019; Thomas et al 2019; Arrieta-Villegas et al., 2020; Ferreras-Colino et al., 2022). P22 is a multi-protein complex obtained from *Mycobacterium bovis* purified protein derivative (B-PPD) by affinity chromatography (Infantes-Lorenzo et al., 2017). P22 has demonstrated high sensitivity (Se) and specificity (Sp), depending on the animal species and epidemiological situation (Infantes-Lorenzo, Moreno et al., 2019). In the case of cattle Se was 87% (Casal et al., 2017) and Sp was 92.5-99.4 depending on cut-off and epidemiological situation (Infantes-Lorenzo, Moreno et al., 2019). However, no data about the antibody response to P22 in European bison are available.

Another disease of livestock and wildlife that has significant consequences for animal and public health, as well as international trade,

is brucellosis (Khurana et al., 2021). *Brucella abortus* is the primary cause of brucellosis in cattle, but it also affects wildlife, including bison (Mackintosh et al., 2002; Rhyan et al., 2001; Tessaro et al., 1990). A recent meta-analysis found that among wild Bovidae, the highest prevalence of brucellosis was noted in American bison (*Bison bison*) (39.5%) (Dadar et al., 2021). Highly similar *B. abortus* variable number tandem repeat patterns have been identified in cattle and American bison, indicating possible transmission between wildlife and livestock (O'Brien et al., 2017; Rhyan et al., 2013). As *B. abortus* was also noted in American bison in northern Canada (Wobeser 2009; Shury et al., 2016), it would be advisable to monitor European bison for brucellosis.

Although only a few studies have been performed on brucellosis in European bison, their findings indicate that antibodies to *B. abortus* are generally absent or only present in small numbers (Kita & Anusz, 1991; Krzyśiak et al., 2018). It is important that endangered species are monitored for diseases causing abortions. To this end, recent studies have attempted to evaluate the potential of serological tests against brucellosis (e.g. Sánchez-Sarmiento et al., 2020).

The aim of this article was to describe the serum antibody response of European bison to the P22 protein complex and to *Brucella* spp. and related factors: gender, age and locality.

2 | MATERIALS AND METHODS

2.1 | Materials

Between October 2017 and April 2022, a total of 323 blood samples were collected from captive ($n = 152$) and free-living ($n = 159$) European bison. Captive European bison were those living in zoos/breeding centres, and free-living, living in different herds in the wild. The animals were derived from different locations, presented in Figure 1. The age of the sampled bison ranged from three months to 25 years (mean 6.5 years). The age of the bison was assessed by experienced veterinarians based on the appearance of horns and tooth eruption (Kraśnińska & Kraśniński, 2017). Age was categorized into three groups: calves (<1-year old), juveniles (1-3-year old), and adults (>3-year old) as described previously (Kraśnińska & Kraśniński, 2017). The samples originated from males ($n = 136$) and females ($n = 175$). The material was collected from living animals following immobilization, or dead ones that had been culled or found dead. Immobilization was performed pharmacologically with the use of a Palmer Cap-Chur tranquilization gun (Palmer Cap-Chur Equipment, Inc.) as described previously (Didkowska et al., 2022). For some animals, it was not possible to acquire data regarding age, locality, or sex; therefore, these animals were not considered when performing statistics.

The blood was collected into 6-9 mL sterile tubes with a clot activator with a needle of a diameter of 1.2 mm from a jugular vein (*vena jugularis externa*), or sometimes from the tail vein (*vena caudalis mediana*). In the case of dead animals, blood was collected from the heart and body cavities. The tubes containing the blood were refrigerated and transported immediately to the laboratory. In the laboratory, the



FIGURE 1 Map of Poland indicating the distribution of European bison (*Bison bonasus*) sampling localities included in the study. The Białowieża Forest, Bieszczady Mountains and Borecka Forest locations include both captive and free-living herds, whereas Knyszyńska Forest only includes a free-living herd. All other localities include only captive herds.

blood was centrifuged (3000 g, 10 min) and the serum was separated. To minimize the risk of false positive results, all extensively hemolysed samples were excluded from the study. The samples were stored at -20°C until analysis.

2.2 | Enzyme-linked immunosorbent assays (ELISA)

Serum samples were brought to room temperature (RT) and then subjected to two serological tests: an in-house ELISA for anti-P22 antibodies and a commercial test for detecting antibodies against *Brucella* spp.

To detect anti-P22 antibodies, the sera were tested with an in-house indirect P22-ELISA as described previously (Thomas et al., 2019). Positive controls were derived from TB-positive cervids (compatible lesions, *M. bovis* positive culture, OD > 1 at P22-ELISA). Negative controls were derived from TB-negative cervids (the absence of compatible lesions, negative culture, OD < 0.2 at P22-ELISA). Briefly, the ELISA plate was coated with P22 at 10 $\mu\text{g}/\text{mL}$ in phosphate-buffered saline solution (PBS; Panreac Química S.L.U.) and incubated overnight at 4°C . After washing with PBS solution containing 0.05% Tween-20 (PBST) (Tween 20; Sigma-Aldrich Química SA), the wells were blocked with 5% skimmed milk powder solution in PBS (SM) for 1 h at RT. Then tested sera and controls were added in 1:10 dilution in SM. After 1-h incubation at 37°C , the plates were washed three times with PBST. Next, protein G horseradish peroxidase (HRP)-conjugated (Sigma-Aldrich Química SA) at a concentration of 0.002 mg/mL in PBS

was added. After incubation for 1 h at RT, the plates were washed with PBST, and o-phenylenediamine-dihydrochloride substrate (FAST OPD; Sigma-Aldrich Química SA) was added. After incubation for 20 min at RT, the reaction was stopped by adding $\text{H}_2\text{SO}_4 \cdot 3\text{N}$. The optical density (OD) was read in a spectrophotometer at a wavelength of 492 nm. The result for each sample was expressed as ELISA percentage ($E\%$). $E\%$ was calculated according to the following formula: $[\text{sample } E\% = (\text{sample OD}/2 \times \text{mean of negative control OD}) \times 100]$. The cut-off value of 110 was chosen based on the plotted results (Figure 2). A conservative cut-off of 110 $E\%$ was chosen to maximize specificity (Infantes-Lorenzo, Moreno et al., 2019).

To detect antibodies against *Brucella* spp., a commercial ELISA R.10.BRU.K3 INgezim BRUCELLA Compacis (Ingenasa) test based on a blocking enzyme immunoassay was performed according to the manufacturer's instructions. The test allows the detection of specific antibodies to the LPS of *Brucella* spp. in multiple animal species (Muñoz et al. 2010). The cut-off value was 40, as instructed. Absorbance was read with a spectrophotometer at 450 nm. According to manufacturers' instructions, the sensitivity and specificity of the test for cattle are 98% and 99.9%.

2.3 | Statistical analysis

The effects of sex, age, captive/free, year of collecting material and locality were assessed by homogeneity tests (two-tailed Fisher's exact test and chi-square test where needed). In addition, 95% confidence




FIGURE 2 ELISA percentage (E%) results of Polish European bison (*Bison bonasus*) antibodies against protein complex P22 by age category (calves, juveniles and adults). The solid red line indicates the E% value above which the result was considered positive (i.e. cut-off); E% = 110.

intervals (95%CI) were calculated for seroprevalence using Sterne's exact method.

2.4 | Ethical statement

No animal was culled or immobilized specifically for this study. *Ante-mortem* sampling (e.g. during translocation, putting the collar) was performed as part of standard veterinary care. The procedures did not require ethical approval according to II Local Ethical Committee for Animal Experiments (Warsaw, Poland).

3 | RESULTS

3.1 | Antibodies against the *M. tuberculosis* complex

Antibodies against P22 were detected in 6.81% (95% CI: 4.44–10.17; 22/323; Table 1) of the tested European bison. The anti-MTC antibody positivity was not significantly different by sex (Fisher's exact test, $p = 0.085$), nor age category (chi-square test, chi-square value = 3.45, degree of freedom = 2, $p = 0.18$). E% results had a similar distribution for each age category. For each age category, a few individuals showed high antibody responses (6 of 51 calves, 11.8%; 7 of 83 juveniles, 8.4%; 8 of 170 adults, 4.7%) (Figure 2). The MTC antibody positivity was not significantly different by captive/free-living status (Fisher's exact test, $p = 1$). Data on seroprevalence by year are presented in Table 1.

The seropositive bison were found in 6 of the 13 sampling localities (46.15%), including the four free-living ones (Table 1). In these six localities, at least 19 bison samples were tested (range 19–72); in contrast, in the remaining eight localities, only 1–8 samples were tested. Thus, comparisons among localities were only performed for localities with >19 samples (chi-square test, chi-square value = 3.3, degree of freedom = 5, $p = 0.65$). Seroprevalence slightly increased with sample size (Figure 3).

3.2 | *Brucella* spp. ELISA results


The general seroprevalence for *Brucella* spp. was 35.91% (95% CI: 30.79–41.32; 116/323; Table 2). Anti-*Brucella* spp. antibody prevalence was not significantly different by sex (Fisher's exact test, $p = 0.64$); however, it was significantly different by both captive/free-living status (Fisher's exact test, $p < 0.001$) and age (chi square test, chi-square value = 16.18, degree of freedom = 2, $p = 0.003$). All age categories demonstrated a similar E% distribution, as shown in Figure 4. For each age category, a significant number of individuals were positive: 23 of 51 calves (45.1%), 42 of 83 juveniles (50.6%), and 45 of 170 adults (26.5%). Data on seroprevalence by year are presented in Table 2.

Seropositive bison were found in 12 of the 13 sampling localities (92.31%) (Table 2). The comparisons among localities only included the localities where at least 19 bison samples, that is larger groups, were

TABLE 1 The frequencies of seropositive to MTC European bison (*Bison bonasus*) regarding their different characteristics.

Variable	Number seropositive to all samples tested
Population type	21/311 (6.8%; 95% CI: 4.37–10.08)
Captive	10/152 (6.6%; 95% CI: 3.52–11.75)
Free-living	11/159 (6.9%, 95% CI: 3.62–12.16)
Sex	18/311 (5.8%, 95% CI: 3.66–8.96)
Female	14/175 (8%, 95% CI: 4.76–13.06)
Male	4/136 (2.9%, 95% CI: 1.02–7.25)
Age group	21/304 (6.9%, 95% CI: 4.47–10.32)
Calves (<1-year old)	6/51 (11.8%, 95% CI: 5.25–23.36)
Juveniles (1–3-year old)	7/83 (8.4%, 95% CI: 4.03–16.69)
Adults (>3-year old)	8/170 (4.7%, 95% CI: 2.22–9.03)
Sampling year	
2017	0/12 (0%)
2018	10/97 (10.3%, 95% CI: 5.51–17.93)
2019	4/47 (8.5%, 95% CI: 2.96–19.99)
2020	2/45 (4.4%, 95% CI: 0.8–15.20)
2021	2/84 (2.4%, 95% CI: 0.43–8.16)
2022	2/38 (5.3%, 95% CI: 0.95–17.99)
Localities	
Bałtów (captive)	0/2 (0%)
Białowiska Forest	5/72 (6.9%, 95% CI: 2.78–15.13)
Captive	3/38 (7.9%, 95% CI: 2.19–20.79)
Free-living	2/34 (5.9%, 95% CI: 1.06–20.11)
Bieszczady Mountains	5/67 (7.5%, 95% CI: 2.99–16.25)
Captive	2/16 (12.5%, 95% CI: 2.27–37.16)
Free-living	3/51 (5.9%, 95% CI: 1.63–16.36)
Borecka Forest	2/45 (4.4%, 95% CI: 0.8–15.2)
Captive	0/8 (0%)
Free-living	2/37 (5.4%, 95% CI: 0.97–18.48)
Gdańsk zoo (captive)	0/2 (0%)
Gołuchów (captive)	0/7 (0%)
Kiermusy (captive)	0/8 (0%)
Międzyzdroje (captive)	0/1 (0%)
Niepołomice (captive)	3/19 (15.8%, 95% CI: 4.45–39.19)
Pszczyna (captive)	2/38 (5.3%, 95% CI: 0.95–17.99)
Knyszyńska Forest (free-living)	4/37 (10.8%, 95% CI: 3.78–25.38)
Ustroń (captive)	0/4 (0%)
Warszawa zoo (captive)	0/8 (0%)

tested (range 19–72): If the sample size was small, it was impossible to draw reliable conclusions. The anti-*Brucella* spp. seroprevalence was significantly different by locality (chi-square test, chi-square value = 16.383, degree of freedom = 5, $p = 0.006$). The seroprevalence relation to sample size is shown in Figure 3. *Brucella* spp. seroprevalence decreased with sample size.



Blue: *Brucella* spp.; Red: *Mycobacterium tuberculosis* complex; Triangle: captive; Square: free-ranging.

FIGURE 3 The *Mycobacterium tuberculosis* complex (MTC) and *Brucella* spp. seroprevalence in Polish European bison (*Bison bonasus*) in relation to sample sizes. Blue colour: *Brucella* spp. seroprevalence; red colour: MTC (protein complex P22) seroprevalence; triangle: captive herd; square: free-living herd. Dashed regression lines are plotted for MTC and *Brucella* spp. antibody prevalence.

4 | DISCUSSION

Our findings indicate a 6.81% prevalence of anti-MTC antibodies and a 35.91% prevalence of anti-*Brucella* spp. antibodies in European bison from Poland. This is the first confirmation of an antibody response to P22 in bison. The correct diagnosis of TB still presents a challenge in this species, so every new diagnostic tool is desirable (Didkowska, Krajewska-Wędzina et al., 2021). In protected species, there is also a greater need to monitor diseases that can cause reproductive alterations, such as brucellosis, as these can reduce the size of the population and thus its genetic variability, which is already low in European bison (Druet et al., 2020). It should be emphasized that our findings are particularly valuable as they were obtained from a large group of animals over several years; this was necessary as the material was taken from a protected species. This study indicates the current trends regarding the diagnostic value of P22 in European bison and microbiological tests toward *Brucella* spp., highlighting the need for further research.

To confront the problem posed by TB in Polish European bison herds, efforts have been made to find cheap, convenient, and highly sensitive diagnostic tools by improving sample collection (Didkowska et al., 2020) and imaging methods (Didkowska et al., 2021d). Even though culture remains the gold standard for TB diagnosis, it is not a convenient tool for monitoring wildlife (Thomas et al. 2021). Microbiological tests are quite expensive, require long wait times and entail

difficulties in collecting suitable antemortem material due to periodic shedding (Cousins & Florisson, 2005). Therefore, some studies suggest using serological TB monitoring as an alternative (Thomas et al., 2021). However, few have evaluated the potential of TB serological assays in European bison. Didkowska, Krajewska-Wędzina et al. (2021) found *Mycobacterium caprae*-infected European bison to demonstrate responses to the antibodies ESAT-6, CFP-10, MPB70, MPB83, CFP10/ESAT-6, MPB70/MPB83, DID65 and B-PPD in dual-path platform, multi-antigen print immunoassay (MAPIA) and IDEXX *M. bovis* Ab ELISA (IDEXX Laboratories, Inc.); however, this study (Didkowska, Krajewska-Wędzina, et al., 2021) was performed on a limited number of samples, including only four positives. More recently, serological monitoring based on MPB70 and MPB83 has shown much lower MTC antibody seroprevalence in European bison (1.96%; Krzysiak et al., 2022).

This is the first serological study to use P22 in European bison. Considering Poland that is generally free of brucellosis and TB in cattle, we used caution when interpreting the results. We also considered the lack of bison reference sera and that field samples, as opposed to experimental infection derived ones, can yield lower specificity (Ferrerias-Colino et al., 2022). Therefore, we used a conservative cut-off value of 110 instead of the commonly used cut-off value of 100 (Thomas, Infantes-Lorenzo, Moreno, Cano-Terriza, et al., 2019). The cut-off value of 110 was chosen based on the plotted results (Figure 2). With a cut-off of 110, the MTC seroprevalence in European bison

TABLE 2 The frequencies of seropositive to *Brucella* spp. European bison (*Bison bonasus*) regarding their different characteristics.

Variable	Number seropositive to all samples tested
Population type	113/311 (36.3%; 95% CI: 31.84–40.99)
Captive	75/152 (49.3%; 95% CI: 41.44–57.52)
Free-living	38/159 (23.9%; 95% CI: 17.87–31.09)
Sex	112/311 (36%, 95% CI: 30.86–41.63)
Female	61/175 (34.9%; 95% CI: 27.96–42.27)
Male	51/136 (37.5%; 95% CI: 29.74–45.94)
Age group	110/304 (36.2%, 95% CI: 30.91–41.76)
Calves (≤ 1 -year old)	23/51 (45.1%; 95% CI: 31.82–58.87)
Juveniles (2–3-year old)	42/83 (50.6%; 95% CI: 39.72–61.49)
Adults (≥ 4 -year old)	45/170 (26.5%; 95% CI: 20.24–33.78)
Year of collecting blood	
2017	7/12 (58.3%, 95% CI: 29.40–81.89)
2018	45/97 (46.4%, 95% CI: 36.56–56.72)
2019	15/47 (31.9%, 95% CI: 20–46.78)
2020	10/45 (22.2%, 95% CI: 11.84–36.56)
2021	28/84 (33.3%, 95% CI: 26.39–44.02)
2022	11/38 (28.9%, 95% CI: 16.67–45.29)
Localities	
Bałtów (captive)	2/2 (100%)
Białowieża Forest	35/72 (48.6%, 95% CI: 36.95–60.74)
Captive	24/38 (63.2%, 95% CI: 46.53–77.38)
Free-living	11/34 (32.4%, 95% CI: 18.85–50)
Bieszczady Mountains	17/67 (25.4%, 95% CI: 16.26–37.24)
Captive	8/16 (50%, 95% CI: 27.20–72.80)
Free-living	9/51 (17.5%, 95% CI: 9.26–30.27)
Borecka Forest	9/45 (20%, 95% CI: 10.57–34.30)
Captive	3/8 (37.5%, 95% CI: 11.12–71.07)
Free-living	6/37 (16.2%, 95% CI: 7.31–32.19)
Gdańsk-zoo (captive)	0/2 (0%)
Gołuchów (captive)	0/2 (0%)
Kiermsy (captive)	1/8 (12.5%, 95% CI: 0.64–50)
Międzyzdroje (captive)	1/1 (100%)
Niepołomice (captive)	8/19 (42.1%, 95% CI: 22.19–65.51)
Pszczyna (captive)	18/38 (47.4%, 95% CI: 31.35–63.28)
Knyszyńska Forest (free-living)	12/37 (32.4%, 95% CI: 18.49–48.64)
Ustroń (captive)	2/4 (50%, 95% CI: 9.77 to 90.23)
Warszawa-zoo (captive)	2/8 (25%, 95% CI: 4.64–63.53)

was 6.81% (95% CI: 4.44–10.17; 22/323) and varied from 0% to 8.8% depending on the sampling year (Table 1).

A previous study on TB microbiological monitoring carried out from 2017 to 2019 isolated *M. caprae* from 4.44% (4/90) European bison;

however, all animals in the study derived from a single captive herd in central Poland (Didkowska, Orłowska, et al., 2021). Animals from this herd have not been tested in the present study. In another region of Poland where TB is being confirmed in wildlife—Bieszczady Mountains (Krajewska-Wędzina et al., 2023), there was no higher seroprevalence than in other free-living sites (Table 2). As such, our findings are higher than expected, and the occurrence of MTC in different localizations is surprising. This can be explained in several ways. The first explanation may be that the P22-ELISA was characterized by high Se; however, it would be necessary to examine the selected animals microbiologically to confirm this. If the test turned out to be very sensitive and specific, it would be a great diagnostic tool for European bison.

Alternatively, a cross-reaction may be present with atypical mycobacteria, a well-known phenomenon in TB diagnostics (Jenkins et al., 2018; Raffo et al., 2020). However, Didkowska, Orłowska et al. (2021) reported that atypical mycobacteria were isolated from only 2.22% (2/90) of tested European bison. It should be highlighted that P22-ELISA offers fair to good specificity (Sp) in detecting antibodies against MTC, depending on the species, and so it also would not necessarily be perfect for European bison. In other ruminants, the tested Sp was 92.5%–99.4% in cattle, 30.9%–78% in goats, 94.4%–100% in sheep and 99.0% in red deer (Thomas, Infantes-Lorenzo, Moreno, Romero, et al., 2019). Another hypothesis is that bison could have contact with other mycobacteria belonging to the MTC, such as *Mycobacterium microti*. *M. microti* infections are increasingly reported in free-living wild animals in Europe (Ghielmetti et al., 2021; Michelet et al., 2021; Pérez de Val et al., 2019; Tagliapietra et al., 2021).

In contrast to most previous studies (Joly & Messier, 2004; Varela-Castro et al., 2020), our findings indicate that the MTC seroprevalence is not significantly different by sex and age. This may be influenced by the behaviour of bison and their gregarious nature (Kraśnińska & Kraśniński, 2017). In addition, the MTC seroprevalence is not significantly different by status and locality. In the case of locality, this is surprising as in previous years, TB was microbiologically confirmed only in two captive herds, namely one herd in Smardzewice, central Poland and another (Wolisko) in Borecka Forest, and in two free-living herds, namely in the Bieszczady Mountains and Borecka Forest (Krajewska et al., 2011, 2016; Welz et al., 2005); therefore, it was anticipated that most positive results would be identified in these locations. In addition, localities with the highest seroprevalence (Białowieża Forest, Bieszczady Mountains, Niepołomice and Pszczyna) are not adjacent to each other (Figure 1, Table 1). In Poland, free-living bison populations are rarely spatially connected with each other (Perzanowski et al., 2019), and new areas are rarely colonised (Ziółkowska et al., 2016); in such cases, therefore, there is probably no bison-driven pathogen transmission between localities. Another important consideration regarding the effect of locality is that, in the case of free-living herds in Poland, the carrying capacity of forest complexes is often exceeded, and so captive herds generally have higher densities (Olech & Perzanowski, 2014). Alternatively, the increase of seroprevalence with sample size (Figure 3) might just point at a higher probability of finding cross-reactions in larger samples.

Previous reports confirmed a sporadic occurrence of *B. abortus* antibodies in European bison (Kita & Anusz, 1991; Krzysiak et al., 2018).




FIGURE 4 ELISA percentage (E%) results of *Brucella* spp. antibodies by age category (calves, juveniles and adults) in European bison (*Bison bonasus*) from Poland. The solid red line indicates the E% value above which the result was considered positive (i.e. cut-off): E% = 40.

The 35.91% (95% CI: 30.79–41.32; 116/323) seroprevalence of *Brucella* spp. identified herein suggests that the Polish European bison population has had contact with these bacteria. The annual seroprevalence has varied significantly over time. The period from 2017 to 2020 saw a downward trend, and this was then followed by a relatively constant level (Table 2). Due to the epizootic situation in Poland, with no case of *B. abortus* being reported since its eradication in 1980, it is likely that any such had been with other species of *Brucella*. Indeed, *Brucella suis* biovar 2 has been isolated from Polish cattle with positive serological results (Szulowski et al., 2013a), and this variant has also been confirmed in Polish wildlife, mainly wild boar (*Sus scrofa*) (Szulowski et al., 2013b). It would be advisable that microbiological monitoring is continued, as *Brucella* spp. can also be a potential threat to European bison, especially in the localities where seroprevalence was high, such as Pszczyna and Białowieża (Table 2).

Our findings indicate that the *Brucella* spp. seroprevalence was significantly different by age; however, unlike most studies, (Alhamada et al., 2017; Asgedom et al., 2016; Chaka et al., 2018), the highest seropositivity was noted in juveniles, not in adults (Table 2). It is possible that in such cases, the animal may have made first contact with the pathogen without the protection offered by colostrum antibodies and these would demonstrate the highest serological response. In American bison, most calves were seronegative by 5 months, and the highest seroconversion was noted at 2 years (Rhyan et al., 2009). In the present study, seropositivity was also more likely in captive than free ranging bison (Table 2). This may be due to the higher animal densities characteristic of captive herds. Localities were also found to have a significant effect, which might be related to a greater presence of bacteria in the areas concerned. Finally, sex was found to have no such effect, and this is in-line with previous studies on American bison, where sex had little influence on the prevalence of *B. abortus* antibodies (Scurlock & Edwards, 2010).

This study has some limitations. First, random sampling was not possible. In addition, the tests were not specially designed for European bison. Furthermore, as the true status of the bison remains unknown, further microbiological testing should be included in later studies. In addition, it should be considered that the controls for MTC seroprevalence testing were taken from a different species (deer). What is more, even though following the recommendations of the OIE-World Organisation for Animal Health, complement fixation tests and serum agglutination tests are being replaced by more sensitive and specific tools for screening brucellosis in mammalian species, including indirect ELISA and fluorescence polarization assay (Greiner et al., 2009; Muñoz et al., 2010; Ragan et al., 2013); most brucellosis serological tests intended for use in wildlife still need validation and standardization (Dadar et al., 2021). One further limitation for *Brucella* sp. results are cross-reactions with *Yersinia enterocolitica* O:9, *Stenotrophomonas maltophilia*, *Salmonella urbana*, *Escherichia coli* O:157, or non-specific reactions associated with the condition of the animal (Ducrottoy et al., 2016; Nielsen et al., 2004).

The obtained results for the MTC and *Brucella* spp. serological surveys raise another important problem. As European bison can be a

reservoir of other diseases, care should be taken when introducing them into new areas, as in Spain (Castillo et al., 2019) or Romania (Dănilă et al., 2022).

5 | CONCLUSIONS

This study is the first to report an antibody response to the P22 multi-protein complex in European bison. MTC serological survey showed that 6.81% of the examined European bison are seropositive, which is higher than noted in previous studies using specific *M. bovis* antigens. In addition, more than 1/3 of the tested European bison were seropositive against *Brucella* spp., which shows another direction for future research. Our findings suggest that brucellosis and TB may represent an emerging threat to the largest European bison population. As Poland is TB and brucellosis free, obtaining many positive results might be due to cross-reactions. In the future, it would be worth testing European bison in Poland for pathogens that can cause false positive results.

AUTHOR CONTRIBUTIONS

Conceptualization; data curation; formal analysis; investigation; project administration; resources; writing—original draft: Anna Didkowska. *Data curation; formal analysis; investigation; resources; visualization; writing—review & editing:* Elisa Ferreras. *Data curation; funding acquisition; project administration; resources; writing—review & editing:* Wanda Olech. *Investigation:* Huggette Gloddy. *Supervision; writing—review & editing:* Krzysztof Anusz. *Methodology; supervision; writing—review & editing:* Jose Antonio Infantes-Lorenzo. *K conceptualization; funding acquisition; methodology; supervision; visualization; writing—original draft; writing—review & editing:* Christian Gortazar.

ACKNOWLEDGEMENTS

We thank 'Wildlife Conservation Network Grants' for supporting HG.

CONFLICT OF INTEREST STATEMENT

Authors declared no conflicts of interest.

FUNDING INFORMATION

Complex project of European bison conservation by State Forests, Forest Fund (Poland), contract no OR.271.3.10.2017

DATA AVAILABILITY STATEMENT

Data are available in Department of Food Hygiene and Public Health Protection, Institute of Veterinary Medicine, Warsaw University of Life Sciences (SGGW).

ETHICS STATEMENT

No animal was culled or immobilized specifically for this study. *Ante-mortem* sampling (e.g. during translocation, putting the collar) was performed as part of standard veterinary care. The procedures did not require ethical approval according to II Local Ethical Committee for Animal Experiments (Warsaw, Poland).

ORCID

Anna Didkowska  <https://orcid.org/0000-0003-1073-174X>

Elisa Ferreras Colino  <https://orcid.org/0000-0002-4790-5259>

Wanda Olech  <https://orcid.org/0000-0002-6166-3954>

Krzysztof Anusz  <https://orcid.org/0000-0002-8116-8491>

José Antonio Infantes-Lorenzo  <https://orcid.org/0000-0002-0446-7682>

Christian Gortázar  <https://orcid.org/0000-0003-0012-4006>

REFERENCES

- Alhamada, A. G., Habib, I., Barnes, A., & Robertson, I. (2017). Risk factors associated with *Brucella* seropositivity in sheep and goats in Duhok province. *Iraqi Journal of Veterinary Sciences*, 4, 65. <https://doi.org/10.3390/vetsci4040065>
- Arrieta-Villegas, C., Infantes-Lorenzo, J. A., Bezos, J., Grasa, M., Vidal, E., Mercader, I., Singh, M., Domingo, M., de Juan, L., & Pérez de Val, B. (2020). Evaluation of P22 antigenic complex for the immuno-diagnosis of tuberculosis in BCG vaccinated and unvaccinated goats. *Frontiers in Veterinary Science*, 7, 374. <https://doi.org/10.3389/fvets.2020.00374>
- Asgedom, H., Damena, D., & Duguma, R. (2016). Seroprevalence of bovine brucellosis and associated risk factors in and around Alage district, Ethiopia. *SpringerPlus*, 5(1), 851. <https://doi.org/10.1186/s40064-016-2547-0>
- Bezos, J., Roy, Á., Infantes-Lorenzo, J. A., González, I., Venteo, Á., Romero, B., Grau, A., Mínguez, O., Domínguez, L., & de Juan, L. (2018). The use of serological tests in combination with the intradermal tuberculin test maximizes the detection of tuberculosis infected goats. *Veterinary Immunology and Immunopathology*, 199, 43–52. <https://doi.org/10.1016/j.vetimm.2018.03.006>
- Casal, C., Infantes, J. A., Rivalde, M. A., Díez-Guerrier, A., Domínguez, M., Moreno, I., Romero, B., de Juan, L., Sáez, J. L., Juste, R., Gortázar, C., Domínguez, L., & Bezos, J. (2017). Antibody detection tests improve the sensitivity of tuberculosis diagnosis in cattle. *Research in Veterinary Science*, 112, 214–221. <https://doi.org/10.1016/j.rvsc.2017.05.012>
- Castillo, M. F. (2019). European bison restoration in Spain—9 Years of the work. *European Bison Conservation Newsletter*, 12, 45–52.
- Chaka, H., Aboset, G., Garoma, A., Gumi, B., & Thys, E. (2018). Cross-sectional survey of brucellosis and associated risk factors in the livestock–wildlife interface area of Nechisar National Park, Ethiopia. *Tropical Animal Health and Production*, 50(5), 1041–1049. <https://doi.org/10.1007/s11250-018-1528-4>
- European Union. (2021). Commission Implementing Regulation (EU) 2021/620 of 15 April 2021 laying down rules for the application of Regulation (EU) 2016/429 of the European Parliament and of the Council as regards the approval of the disease-free and non-vaccination status of certain Member States or zones or compartments thereof as regards certain listed diseases and the approval of eradication programmes for those listed diseases (Text with EEA relevance). *OJL*, 131, 78–119.
- Cousins, D. V., & Florisson, N. (2005). A review of tests available for use in the diagnosis of tuberculosis in non-bovine species. *Revue Scientifique et Technique (International Office of Epizootics)*, 24(3), 1039–1059.
- Dadar, M., Shahali, Y., Fakhri, Y., & Godfroid, J. (2021). The global epidemiology of *Brucella* infections in terrestrial wildlife: A meta-analysis. *Transboundary and Emerging Diseases*, 68(2), 715–729. <https://doi.org/10.1111/tbed.13735>
- Dănilă, G., Cătănoiu, S., Simioniu, V., & Roșca, S. (2022). The reintroduction analysis of European bison (*Bison bonasus* L., 1758) in the North of Romania and the identification of the most favourable locations. *Forests*, 13, 920. <https://doi.org/10.3390/f13060920>
- Didkowska, A., Dziekan, P., Czujkowska, A., Bereznowski, A., Witkowski, L., Orłowska, B., Wiśniewski, J., Krzysiak, M., Krajewska-Wędzina, M., Bruczyńska, M., Żychska, M., Olech, W., & Anusz, K. (2021). The first visually-guided bronchoscopy in European bison (*Bison bonasus*)—An additional tool in the diagnosis of bovine tuberculosis? *Veterinary and Animal Science*, 12, 100174. <https://doi.org/10.1016/j.vas.2021.100174>
- Didkowska, A., Krajewska-Wędzina, M., Bielecki, W., Brzezińska, S., Augustynowicz-Kopeć, E., Olech, W., Anusz, K., Sridhara, A. A., Johnathan-Lee, A., Elahi, R., Miller, M. A., Ray Waters, W., & Lyashchenko, K. P. (2021). Antibody responses in European bison (*Bison bonasus*) naturally infected with *Mycobacterium caprae*. *Veterinary Microbiology*, 253, 108952. <https://doi.org/10.1016/j.vetmic.2020.108952>
- Didkowska, A., Orłowska, B., Krajewska-Wędzina, M., Augustynowicz-Kopeć, E., Brzezińska, S., Żygowska, M., Wiśniewski, J., Kaczor, S., Welz, M., Olech, W., & Anusz, K. (2021). Microbiological and molecular monitoring for bovine tuberculosis in the Polish population of European bison (*Bison bonasus*). *Annals of Agricultural and Environmental Medicine: AAEM*, 28(4), 575–578. <https://doi.org/10.26444/aaem/130822>
- Didkowska, A., Orłowska, B., Krajewska-Wędzina, M., Krzysiak, M., Bruczyńska, M., Wiśniewski, J., Klich, D., Olech, W., & Anusz, K. (2022). Intra-palpebral tuberculin skin test and interferon gamma release assay in diagnosing tuberculosis due to *Mycobacterium caprae* in European bison (*Bison bonasus*). *Pathogens*, 11(2), 260. <https://doi.org/10.3390/pathogens11020260>
- Didkowska, A., Orłowska, B., Witkowski, L., Olbrych, K., Brzezińska, S., Augustynowicz-Kopeć, E., Krajewska-Wędzina, M., Bereznowski, A., Bielecki, W., Krzysiak, M., Rakowska, A., Olech, W., Miller, M. A., Waters, W. R., Lyashchenko, K. P., & Anusz, K. (2020). Biopsy and tracheo-bronchial aspirates as additional tools for the diagnosis of bovine tuberculosis in living European bison (*Bison bonasus*). *Animals: An Open Access Journal from MDPI*, 10(11), 2017. <https://doi.org/10.3390/ani10112017>
- Didkowska, A., Ptak, M., Opalczyńska, K., Wojciechowska, M., Olech, W., & Anusz, K. (2021). A serological survey of paratuberculosis in the Polish European bison (*Bison bonasus*) population in 2018–2021. *Animals: An Open Access Journal from MDPI*, 11(7), 2094. <https://doi.org/10.3390/ani11072094>
- Druet, T., Oleński, K., Flori, L., Bertrand, A. R., Olech, W., Tokarska, M., Kaminski, S., & Gautier, M. (2020). Genomic footprints of recovery in the European bison. *The Journal of Heredity*, 111(2), 194–203. <https://doi.org/10.1093/jhered/esaa002>
- Ducrottoy, M. J., Conde-Álvarez, R., Blasco, J. M., & Moriyón, I. (2016). A review of the basis of the immunological diagnosis of ruminant brucellosis. *Veterinary Immunology and Immunopathology*, 171, 81–102. <https://doi.org/10.1016/j.vetimm.2016.02.002>
- Ferreras-Colino, E., Moreno, I., Arnal, M. C., Balseiro, A., Acevedo, P., Domínguez, M., Fernández de Luco, D., Gortázar, C., & Rivalde, M. A. (2022). Is serology a realistic approach for monitoring red deer tuberculosis in the field? *Preventive Veterinary Medicine*, 202, 105612. <https://doi.org/10.1016/j.prevetmed.2022.105612>
- Ghielmetti, G., Kupca, A. M., Hanczaruk, M., Friedel, U., Weinberger, H., Revilla-Fernández, S., Hofer, E., Riehm, J. M., Stephan, R., & Glawischig, W. (2021). *Mycobacterium microti* infections in free-ranging red deer (*Cervus elaphus*). *Emerging Infectious Diseases*, 27(8), 2025–2032. <https://doi.org/10.3201/eid2708.210634>
- Greiner, M., Verloo, D., & de Massis, F. (2009). Meta-analytical equivalence studies on diagnostic tests for bovine brucellosis allowing assessment of a test against a group of comparative tests. *Preventive Veterinary Medicine*, 92(4), 373–381. <https://doi.org/10.1016/j.prevetmed.2009.07.014>
- Himsworth, C. G., Elkin, B. T., Nishi, J. S., Neimanis, A. S., Wobeser, G. A., Turcotte, C., & Leighton, F. A. (2010). An outbreak of bovine tuberculosis in an intensively managed conservation herd of wild bison in the Northwest Territories. *The Canadian Veterinary Journal = La Revue Veterinaire Canadienne*, 51(6), 593–597.
- Infantes-Lorenzo, J. A., Dave, D., Moreno, I., Anderson, P., Lesellier, S., Gormley, E., Domínguez, L., Balseiro, A., Gortázar, C., Domínguez, M., & Salguero, F. J. (2019). New serological platform for detecting antibodies against *Mycobacterium tuberculosis* complex in European badgers. *Veterinary Medicine and Science*, 5(1), 61–69. <https://doi.org/10.1002/vms3.134>

- Infantes-Lorenzo, J. A., Moreno, I., Rivalde, M., Roy, Á., Villar, M., Romero, B., Ibarrola, N., de la Fuente, J., Puentes, E., de Juan, L., Gortázar, C., Bezos, J., Domínguez, L., & Domínguez, M. (2017). Proteomic characterisation of bovine and avian purified protein derivatives and identification of specific antigens for serodiagnosis of bovine tuberculosis. *Clinical Proteomics*, 14, 36. <https://doi.org/10.1186/s12014-017-9171-z>
- Infantes-Lorenzo, J. A., Moreno, I., Roy, A., Rivalde, M. A., Balseiro, A., de Juan, L., Romero, B., Bezos, J., Puentes, E., Åkerstedt, J., Tessema, G. T., Gortázar, C., Domínguez, L., & Domínguez, M. (2019). Specificity of serological test for detection of tuberculosis in cattle, goats, sheep and pigs under different epidemiological situations. *BMC Veterinary Research*, 15(1), 70. <https://doi.org/10.1186/s12917-019-1814-z>
- Infantes-Lorenzo, J. A., Whitehead, C. E., Moreno, I., Bezos, J., Roy, A., Domínguez, L., Domínguez, M., & Salguero, F. J. (2018). Development and evaluation of a serological assay for the diagnosis of tuberculosis in alpacas and llamas. *Frontiers in Veterinary Science*, 5, 189. <https://doi.org/10.3389/fvets.2018.00189>
- Jenkins, A. O., Gormley, E., Gcebe, N., Fosgate, G. T., Conan, A., Aagaard, C., Michel, A. L., & Rutten, V. (2018). Cross reactive immune responses in cattle arising from exposure to *Mycobacterium bovis* and non-tuberculous mycobacteria. *Preventive Veterinary Medicine*, 152, 16–22. <https://doi.org/10.1016/j.prevetmed.2018.02.003>
- Joly, D. O., & Messier, F. (2004). Factors affecting apparent prevalence of tuberculosis and brucellosis in wood bison. *Journal of Animal Ecology*, 73(4), 623–631. <http://www.jstor.org/stable/3505273>
- Khurana, S. K., Sehrawat, A., Tiwari, R., Prasad, M., Gulati, B., Shabbir, M. Z., Chhabra, R., Karthik, K., Patel, S. K., Pathak, M., Iqbal Yatoo, M., Gupta, V. K., Dhama, K., Sah, R., & Chaicumpa, W. (2021). Bovine brucellosis—A comprehensive review. *The Veterinary Quarterly*, 41(1), 61–88. <https://doi.org/10.1080/01652176.2020.1868616>
- Kita, J., & Anusz, K. (1991). Serologic survey for bovine pathogens in free-ranging European bison from Poland. *Journal of Wildlife Diseases*, 27(1), 16–20. <https://doi.org/10.7589/0090-3558-27.1.16>
- Klich, D., Kitowski, I., Łopucki, R., Wiącek, D., & Olech, W. (2021). Essential differences in the mineral status of free-ranging European bison *Bison bonasus* populations in Poland: The effect of the anthroposphere and lithosphere. *The Science of the Total Environment*, 757, 143926. <https://doi.org/10.1016/j.scitotenv.2020.143926>
- Klich, D., Łopucki, R., Stachniuk, A., Sporek, M., Fornal, E., Wojciechowska, M., & Olech, W. (2020). Pesticides and conservation of large ungulates: Health risk to European bison from plant protection products as a result of crop depredation. *PLoS ONE*, 15(1), e0228243. <https://doi.org/10.1371/journal.pone.0228243>
- Krajewska, M., Lipiec, M., & Szulowski, K. (2011). Bovine tuberculosis in bison (*Bison bonasus caucasicus*) located in Poland. *Postępy Nauk Medycznych*, 10, 842–845.
- Krajewska, M., Lipiec, M., Zabost, A., Augustynowicz-Kopeć, E., & Szulowski, K. (2014). Bovine tuberculosis in a wild boar (*Sus scrofa*) in Poland. *Journal of Wildlife Diseases*, 50(4), 1001–1002. <https://doi.org/10.7589/2013-07-187>
- Krajewska, M., Kozińska, M., Orłowska, B., Augustynowicz-Kopeć, E., Anusz, K., Welz, M., & Szulowski, K. (2016). *Mycobacterium bovis* u żubra w Puszczy Boreckiej. Materiały konferencyjne "Żubry w Krainie Dinozaurów", 8–9 września 2016.
- Krajewska-Wędzina, M., Krzysiak, M. K., Bruczyńska, M., Orłowska, B., Didkowska, A., Radulski, Ł., Wiśniewski, J., Olech, W., Nowakiewicz, A., Welz, M., Kaczor, S., Weiner, M., & Anusz, K. (2023). Ten years of animal tuberculosis monitoring in free-living European bison (*Bison bonasus*) in Poland. *Animals*, 13, 1205. <https://doi.org/10.3390/ani13071205>
- Krasińska, M., & Krasiński, Z. A. (2017). *Żubr. Monografia Przyrodnicza*. Wydawnictwo Chyra.pl.
- Krzysiak, M., Larska, M., Zabost, A., Didkowska, A., Krajewska-Wędzina, M., Anusz, K., & Augustynowicz-Kopeć, E. (2022). Is serological monitoring a fit-for-purpose tool to assess the epidemiological situation of tuberculosis in the sylvatic species of European bison (*Bison bonasus*) in Poland? *Journal of Veterinary Research*, 66(3), 333–344. <https://doi.org/10.2478/jvetres-2022-0038>
- Krzysiak, M. K., Jabłoński, A., Iwaniak, W., Krajewska, M., Kęsik-Maliszewska, J., & Larska, M. (2018). Seroprevalence and risk factors for selected respiratory and reproductive tract pathogen exposure in European bison (*Bison bonasus*) in Poland. *Veterinary Microbiology*, 215, 57–65. <https://doi.org/10.1016/j.vetmic.2018.01.005>
- Mackintosh, C., Haigh, J. C., & Griffin, F. (2002). Bacterial diseases of farmed deer and bison. *Revue Scientifique et Technique (International Office of Epizootics)*, 21(2), 249–263. <https://doi.org/10.20506/rst.21.2.1341>
- Michelet, L., Richomme, C., Réveillaud, E., De Cruz, K., Moyen, J. L., & Boschirolu, M. L. (2021). *Mycobacterium microti* Infection in Red Foxes in France. *Microorganisms*, 9(6), 1257. <https://doi.org/10.3390/microorganisms9061257>
- Muñoz, P. M., Boadella, M., Arnal, M., de Miguel, M. J., Revilla, M., Martínez, D., Vicente, J., Acevedo, P., Oleaga, A., Ruiz-Fons, F., Marín, C. M., Prieto, J. M., de la Fuente, J., Barral, M., Barberán, M., de Luco, D. F., Blasco, J. M., & Gortázar, C. (2010). Spatial distribution and risk factors of Brucellosis in Iberian wild ungulates. *BMC Infectious Diseases*, 10, 46. <https://doi.org/10.1186/1471-2334-10-46>
- Nielsen, K., Smith, P., Widdison, J., Gall, D., Kelly, L., Kelly, W., & Nicoletti, P. (2004). Serological relationship between cattle exposed to *Brucella abortus*, *Yersinia enterocolitica* O:9 and *Escherichia coli* O157:H7. *Veterinary Microbiology*, 100(1–2), 25–30. <https://doi.org/10.1016/j.vetmic.2003.12.010>
- Nishi, J. S., Stephen, C., & Elkin, B. T. (2002). Implications of agricultural and wildlife policy on management and eradication of bovine tuberculosis and brucellosis in free-ranging wood bison of northern Canada. *Annals of the New York Academy of Sciences*, 969, 236–244. <https://doi.org/10.1111/j.1749-6632.2002.tb04385.x>
- O'Brien, M. P., Beja-Pereira, A., Anderson, N., Ceballos, R. M., Edwards, W. H., Harris, B., Wallen, R. L., & Costa, V. (2017). Brucellosis transmission between wildlife and livestock in the Greater Yellowstone Ecosystem: Inferences from DNA genotyping. *Journal of Wildlife Diseases*, 53(2), 339–343. <https://doi.org/10.7589/2015-12-329>
- Olech, W., & Perzanowski, K. (Eds.). (2022). *European bison (Bison bonasus) strategic species status review 2020*. IUCN SSC Bison Specialist Group and European Bison Specialist Group. <https://www.iucn.org/commissions/ssc-groups/mammals/mammals-a-e/bison>
- Olech, W., & Perzanowski, K. (2014). *Podręcznik Najlepszych Praktyk Ochrony Żubra*. Centrum Koordynacji Projektów Środowiskowych.
- Pérez de Val, B., Sanz, A., Soler, M., Allepuz, A., Michelet, L., Boschirolu, M. L., & Vidal, E. (2019). *Mycobacterium microti* infection in free-ranging wild boar, Spain, 2017–2019. *Emerging Infectious Diseases*, 25(11), 2152–2154. <https://doi.org/10.3201/eid2511.190746>
- Perzanowski, K., Bleyhl, B., Olech, W., & Kuemmerle, T. (2019). Connectivity or isolation? Identifying reintroduction sites for multiple conservation objectives for wisents in Poland. *Animal Conservation*, 23, 212–221.
- Plumb, G., Kowalczyk, R., & Hernandez-Blanco, J. A. (2020). *Bison bonasus*. The IUCN Red List of Threatened Species. <https://doi.org/10.2305/IUCN.UK.2020-3.RLTS.T2814A45156279.en>
- Raczynski, J., & Bołbot, M. (2022). *European bison pedigree book 2021*. Białowieża National Park.
- Raffo, E., Steuer, P., Tomckowiack, C., Tejeda, C., Collado, B., & Salgado, M. (2020). More insights about the interfering effect of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) infection on *Mycobacterium bovis* (*M. bovis*) detection in dairy cattle. *Tropical Animal Health and Production*, 52(3), 1479–1485. <https://doi.org/10.1007/s11250-019-02151-2>
- Ragan, V., Vroegindewey, G., & Babcock, S. (2013). International standards for brucellosis prevention and management. *Revue Scientifique et Technique (International Office of Epizootics)*, 32(1), 189–198. <https://doi.org/10.20506/rst.32.1.2203>
- Rhyan, J. C., Gidlewski, T., Roffe, T. J., Aune, K., Philo, L. M., & Ewalt, D. R. (2001). Pathology of brucellosis in bison from Yellowstone National Park.

- Journal of Wildlife Diseases*, 37(1), 101–109. <https://doi.org/10.7589/0090-3558-37.1.101>
- Rhyan, J. C., Aune, K., Roffe, T., Ewalt, D., Hennager, S., Gidlewski, T., Olsen, S., & Clarke, R. (2009). Pathogenesis and epidemiology of brucellosis in Yellowstone bison: Serologic and culture results from adult females and their progeny. *Journal of Wildlife Diseases*, 45(3), 729–739. <https://doi.org/10.7589/0090-3558-45.3.729>
- Rhyan, J. C., Nol, P., Quance, C., Gertonson, A., Belfrage, J., Harris, L., Straka, K., & Robbe-Austerman, S. (2013). Transmission of brucellosis from elk to cattle and bison, Greater Yellowstone area, U.S.A., 2002–2012. *Emerging Infectious Diseases*, 19(12), 1992–1995. <https://doi.org/10.3201/eid1912.130167>
- Sánchez-Sarmiento, A. M., Ruoppolo, V., Muelbert, M. M., Ferreira Neto, J. S., & Catão-Dias, J. L. (2020). Serological screening for *Brucella* spp. and *Leptospira* spp. antibodies in southern elephant seals *Mirounga leonina* from Elephant Island, Antarctica, in 2003 and 2004. *Diseases of Aquatic Organisms*, 142, 161–170. <https://doi.org/10.3354/dao03548>
- Scurlock, B. M., & Edwards, W. H. (2010). Status of brucellosis in free-ranging elk and bison in Wyoming. *Journal of Wildlife Diseases*, 46(2), 442–449. <https://doi.org/10.7589/0090-3558-46.2.442>
- Shury, T. K., Nishi, J. S., Elkin, B. T., & Wobeser, G. A. (2015). Tuberculosis and brucellosis in wood bison (bison bison athabascae) in northern Canada: A renewed need to develop options for future management. *Journal of wildlife diseases*, 51(3), 543–554. <https://doi.org/10.7589/2014-06-167>
- Shury, T. K., Nishi, J. S., Elkin, B. T., & Wobeser, G. A. (2016). Tuberculosis and brucellosis in wood bison (bison bison athabascae) in northern Canada: A renewed need to develop options for future management. *Journal of wildlife diseases*, 51(3), 543–554. <https://doi.org/10.7589/2014-06-167>
- Szulowski, K., Iwaniak, W., Weiner, M., & Złotnicka, J. (2013a). *Brucella suis* biovar 2 isolations from cattle in Poland. *Annals of Agricultural and Environmental Medicine: AAEM*, 20(4), 672–675.
- Szulowski, K., Iwaniak, W., Weiner, M., & Złotnicka, J. (2013b). Characteristics of *Brucella* strains isolated from animals in Poland. *Polish Journal of Veterinary Sciences*, 16(4), 757–758. <https://doi.org/10.2478/pjvs-2013-0109>
- Tagliapietra, V., Boniotti, M. B., Mangeli, A., Karaman, I., Alborali, G., Chiari, M., D'Incau, M., Zanon, M., Rizzoli, A., & Pacciarini, M. L. (2021). *Mycobacterium microti* at the environment and wildlife interface. *Microorganisms*, 9(10), 2084. <https://doi.org/10.3390/microorganisms9102084>
- Tessaro, S. V., Forbes, L. B., & Turcotte, C. (1990). A survey of brucellosis and tuberculosis in bison in and around Wood Buffalo National Park, Canada. *The Canadian Veterinary Journal = La Revue Veterinaire Canadienne*, 31(3), 174–180.
- Thomas, J., Infantes-Lorenzo, J. A., Moreno, I., Cano-Terriza, D., de Juan, L., García-Bocanegra, I., Domínguez, L., Domínguez, M., Gortázar, C., & Rivalde, M. A. (2019). Validation of a new serological assay for the identification of *Mycobacterium tuberculosis* complex-specific antibodies in pigs and wild boar. *Preventive Veterinary Medicine*, 162, 11–17. <https://doi.org/10.1016/j.prevetmed.2018.11.004>
- Thomas, J., Infantes-Lorenzo, J. A., Moreno, I., Romero, B., Garrido, J. M., Juste, R., Domínguez, M., Domínguez, L., Gortázar, C., & Rivalde, M. A. (2019). A new test to detect antibodies against *Mycobacterium tuberculosis* complex in red deer serum. *Veterinary Journal (London, England: 1997)*, 244, 98–103. <https://doi.org/10.1016/j.tvjl.2018.12.021>
- Thomas, J., Balseiro, A., Gortázar, C., & Rivalde, M. A. (2021). Diagnosis of tuberculosis in wildlife: A systematic review. *Veterinary Research*, 52(1), 31. <https://doi.org/10.1186/s13567-020-00881-y>
- Varela-Castro, L., Alvarez, V., Sevilla, I. A., & Barral, M. (2020). Risk factors associated to a high *Mycobacterium tuberculosis* complex seroprevalence in wild boar (*Sus scrofa*) from a low bovine tuberculosis prevalence area. *PLoS ONE*, 15(4), e0231559. <https://doi.org/10.1371/journal.pone.0231559>
- Welz, M., Anusz, K., Salwa, A., Zaleska, M., Bielecki, W., Osinska, B., Kaczor, S., & Kita, J. (2005). Bovine tuberculosis in European bison in the Bieszczady region. *Medycyna Weterynaryjna*, 61, 441–444.
- Wobeser, G. (2009). Bovine tuberculosis in Canadian wildlife: An updated history. *The Canadian Veterinary Journal = La Revue Veterinaire Canadienne*, 50(11), 1169–1176.
- Zimpel, C. K., Brum, J. S., de Souza Filho, A. F., Biondo, A. W., Perotta, J. H., Dib, C. C., Bonat, M., Neto, J. S., Brandão, P. E., Heinemann, M. B., & Guimaraes, A. M. (2017). *Mycobacterium bovis* in a European bison (*Bison bonasus*) raises concerns about tuberculosis in Brazilian captive wildlife populations: A case report. *BMC Research Notes*, 10(1), 91. <https://doi.org/10.1186/s13104-017-2413-3>
- Ziółkowska, E., Perzanowski, K., Bleyhl, B., Ostapowicz, K., & Kuemmerle, T. (2016). Understanding unexpected reintroduction outcomes: Why aren't European bison colonizing suitable habitat in the Carpathians? *Biological Conservation*, 195, 106–117.

How to cite this article: Didkowska, A., Colino, E. F., Olech, W., Gloddy, H., Anusz, K., Infantes-Lorenzo, J. A., & Gortázar, C. (2024). Antibodies against the *Mycobacterium tuberculosis* complex and *Brucella* spp. in captive and free-living European bison (*Bison bonasus*) in Poland. *Veterinary Medicine and Science*, 10, e1314. <https://doi.org/10.1002/vms3.1314>