

This is the peer reviewed version of the following article:

Mehrkens D, Dohr J, Nettersheim FS, Ballmann F, Bastigkeit J, Brückner A, Geissen S, De Vore L, Fitsch C, Diekmann C, de la Fuente-Alonso A, Schelemei P, Picard FR, Kochen M, Arkenberg P, Rappenecker A, Ahdab M, Nemade H, Narayan S, Braumann S, Kreuzberg W, Hof A, Guthoff H, Mensah BQ, Lechner S, Guala A, Evangelista A, Teixido-Tura G, Nistal JF, Campanero MR, Kaemmerer H, Wolf Z, Holdenrieder S, Groenink M, van Andel M, Mieremet A, Pfeiler S, Gerdes N, Flögel U, Zimmermann LM, Sengle G, Eich ML, Schömig-Markiefka B, Adam M, Fleischmann BK, Wenzel D, Redondo JM, de Waard V, Klinke A, Baldus S, Mollenhauer M, Winkels H. Myeloperoxidase aggravates thoracic aortic aneurysm formation in Marfan disease. **Cardiovasc Res.** 2025 Dec 17:cvaf248. doi: 10.1093/cvr/cvaf248. Epub ahead of print. PMID: 41403006.

which has been published in final form at <https://doi.org/10.1093/cvr/cvaf248>