

Table 2. Clinical and immunohistochemical characteristics of FL patients

Variable	Nc*	CBX7 expression							p value†
		1	2	3	4	5	6	7	
Pathological grade‡	158								0.0004
1		14	7	6	3	1	1	2	
2		20	14	14	13	4	7	0	
3		8	6	13	11	8	4	0	
4		0	0	0	1	0	3	0	
t(14;18) translocation	141								0.045
Yes		29	23	15	21	6	8	2	
No		11	3	13	3	4	3	0	
Ki67	156								0.015
1		12	10	8	7	2	3	2	
2		26	11	14	6	4	6	0	
3		4	6	9	13	6	4	0	
4		0	0	1	1	1	2	0	
BCL6	156								0.29
Positive		20	17	19	19	9	11	1	
Negative		22	10	13	8	2	4	1	
CD10	158								0.31
Positive		31	23	21	21	10	8	2	
Negative		11	4	11	6	3	7	0	
ZBTB7	160								0.007
0		7	2	0	2	2	0	0	
1		26	14	20	6	6	8	2	
2		9	11	13	20	5	7	0	
cMYC	107								0.007
0		20	9	13	9	0	5	0	
1		14	11	13	8	2	4	0	
2		0	1	0	3	2	3	0	

*Number of cases.

†Statistical analyses were performed to evaluate the correlation between CBX7 expression and the indicated variables. p value was generated by Fisher's exact test.

‡Pathological grade was determined according to World Health Organization classification.