

ORIGINAL RESEARCH



Wnt5a Promotes Lysosomal Cholesterol Egress and Protects Against Atherosclerosis

Sara Awan¹*, Magalie Lambert¹*, Ali Imtiaz¹, Fabien Alpy¹, Catherine Tomasetto¹, Mustapha Oulad-Abdelghani, Christine Schaeffer¹, Chloé Moritz, Diane Julien-David¹, Souad Najib¹, Laurent O. Martinez¹, Rachel L. Matz, Xavier Collet, Roberto Silva-Rojas¹, Johann Böhm, Joachim Herz¹, Jérôme Terrand,[†] Philippe Boucher¹†

BACKGROUND: Impairment of cellular cholesterol trafficking is at the heart of atherosclerotic lesions formation. This involves egress of cholesterol from the lysosomes and 2 lysosomal proteins, the NPC1 (Niemann–Pick C1) and NPC2 that promotes cholesterol trafficking. However, movement of cholesterol out the lysosome and how disrupted cholesterol trafficking leads to atherosclerosis is unclear. As the Wnt ligand, Wnt5a inhibits the intracellular accumulation of cholesterol in multiple cell types, we tested whether Wnt5a interacts with the lysosomal cholesterol export machinery and studied its role in atherosclerotic lesions formation.

METHODS: We generated mice deleted for the *Wnt5a* gene in vascular smooth muscle cells. To establish whether Wnt5a also protects against cholesterol accumulation in human vascular smooth muscle cells, we used a CRISPR/Cas9 guided nuclease approach to generate human vascular smooth muscle cells knockout for Wnt5a.

RESULTS: We show that Wnt5a is a crucial component of the lysosomal cholesterol export machinery. By increasing lysosomal acid lipase expression, decreasing metabolic signaling by the mTORC1 (mechanistic target of rapamycin complex 1) kinase, and through binding to NPC1 and NPC2, Wnt5a senses changes in dietary cholesterol supply and promotes lysosomal cholesterol egress to the endoplasmic reticulum. Consequently, loss of Wnt5a decoupled mTORC1 from variations in lysosomal sterol levels, disrupted lysosomal function, decreased cholesterol content in the endoplasmic reticulum, and promoted atherosclerosis.

CONCLUSIONS: These results reveal an unexpected function of the Wnt5a pathway as essential for maintaining cholesterol homeostasis in vivo.

GRAPHIC ABSTRACT: A [graphic abstract](#) is available for this article.

Key Words: atherosclerosis ■ homeostasis ■ ligands ■ lysosomes

In This Issue, see p 163 | Meet the First Author, see p 164

Wnt signaling has pleiotropic effects in diverse tissues. It plays an important role in cell differentiation, proliferation, and migration during embryogenesis.¹ It has been associated with diverse human diseases¹ and has been recently recognized as signaling involved in hyperlipemia syndrome and atherosclerosis. In mice, the mutation R611C of the Wnt co-receptor LRP6 (LDL [low-density lipoprotein]

receptor-related protein 6), a member of the LDLR (LDL receptor) gene family causes elevated plasma LDL and TG levels and fatty liver.² In humans, a missense mutation in *LRP6* causes autosomal-dominant early onset cardiovascular disease and metabolic syndrome, including hyperlipidemia, diabetes, osteoporosis, and hypertension.³ *LRP6*_{R611C} mutation carriers exhibit impaired LDL clearance and internalization with elevated serum

Correspondence to: Philippe Boucher, PhD, UMR-S INSERM 1109, University of Strasbourg, 1, place de l'Hôpital, 67000 Strasbourg, France. Email philippe.boucher@unistra.fr

*S. Awan and M. Lambert are joint first authors.

†J. Terrand and P. Boucher are joint senior authors.

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/CIRCRESAHA.121.318881>.

For Sources of Funding and Disclosures, see page 198.

© 2021 American Heart Association, Inc.

Circulation Research is available at www.ahajournals.org/journal/res

Novelty and Significance

What Is Known?

- Disrupted cholesterol trafficking leads to atherosclerosis.
- In humans, missense mutation in the Wnt co-receptor *LRP6* causes autosomal-dominant early onset cardiovascular disease.
- The *LRP6* (LDL [low-density lipoprotein] receptor-related protein 6) ligand, *Wnt5a* is expressed in murine and human atherosclerotic lesions.
- *Wnt5a* deficiency in mouse embryonic fibroblasts promotes cholesterol accumulation within perinuclear cytosolic vesicles.
- *Wnt5a* increases cholesterol export and decreases cholesterol biosynthesis.

What New Information Does This Article Contribute?

- Mice deficient for *Wnt5a* in vascular smooth muscle cells developed twice as much atherosclerosis as controls.
- *Wnt5a* binds to cholesterol-rich membranes and to the lysosomal NPC1 (Niemann–Pick C1) and NPC2 proteins and promotes lysosomal cholesterol egress.

- The loss of *Wnt5a* in vascular smooth muscle cells reduces the lysosomal acid lipase expression and its capacity to catabolize atherogenic lipoproteins.
- *Wnt5a* promotes cholesterol trafficking by down regulating the PI3K/Akt (protein kinase B)/mTORC1 (mechanistic target of rapamycin complex 1) pathway in human vascular smooth muscle cells.

The present studies provide evidence that *Wnt5a* is essential for cholesterol homeostasis. It activates lysosomal function, promotes cholesterol trafficking, and protects against atherosclerosis.

Nonstandard Abbreviations and Acronyms

ACAT1	acyl-CoA cholesterolacyltransferase 1
ACTA	α -smooth muscle actin
AKT	protein kinase B
ATP6V	V-type proton ATPase
CE	cholesteryl ester
CTSF	cathepsin F
DGAT2	acyl-CoA:diacylglycerol acyltransferase 2
ER	endoplasmic reticulum
GAL3	galectin-3
GFP-D4	D4 fragment of perfringolysin O fused to GFP
KLF4	Kruppel-like transcription factor 4
LAL	lysosomal acid lipase
LAMP1	lysosome-associated membrane protein 1
LDL	low-density lipoprotein
LDLR	LDL receptor
LELs	late endosomes/lysosomes
LRP1	LDL receptor-related protein 1
LRP6	LDL receptor-related protein 6
MAPK	mitogen-activated protein kinase
mTORC1	mechanistic target of rapamycin complex 1
NF-κB	nuclear factor κ B
NPC1/2	Niemann–Pick C1/2

p-4EBP1	phospho-eukaryotic translation initiation factor 4E-binding protein 1
p-Akt	phospho-protein kinase B
PDGF	platelet-derived growth factor
PI3K	phosphatidylinositol 3-kinase
SGSH	N-sulphoglucosamine sulphohydrolase
SREBP-2	sterol regulatory element-binding protein-2
TFEB	transcription factor EB
TGF-β	transforming growth factor β
TPP1	tripeptidyl peptidase 1
VSMCs	vascular smooth muscle cells

LDL cholesterol, implicating *LRP6* and Wnt signaling as drivers of cholesterol trafficking.³ More recently, Wnt signaling was also reported to promote resolution of atherosclerosis in mice and human.⁴ However, how Wnt signaling mechanistically causes hyperlipidemia and impaired cholesterol trafficking remains an enigma.

Cholesterol uptake begins when LDL-derived cholesteryl esters are endocytosed through clathrin-coated pits into late endosomes/lysosomes (LELs) where they are hydrolyzed into free cholesterol and fatty acids by LAL (lysosomal acid lipase).⁵ Free cholesterol then binds 2 lysosomal proteins essential for cholesterol egress, the soluble NPC2 (Niemann–Pick C2) which is concentrated in LELs, and the polytopic membrane-bound NPC1 (Niemann–Pick C1).⁵ Upon binding to NPC2 in the lumen

of the lysosome, cholesterol is handed off to NPC1 and then delivered to other cellular compartments including the plasma membrane and eventually the endoplasmic reticulum (ER) where regulation of cholesterol biosynthesis occurs. Heritable mutations in the *NPC1* genes lead to massive cholesterol accumulation within LELs of many organs, including the brain. In macrophages, it causes impaired cholesterol efflux and atherosclerosis.⁶ Whereas the functions of NPC1 and NPC2 are well understood at the molecular levels,⁷ there are other crucial pieces of cholesterol export machinery we know very little about. For instance, *in vitro* binding experiments showed that whereas cholesterol binds rapidly to NPC2, handing it over to NPC1 is a slow process, especially at 4°C suggesting that other components beside NPC proteins are needed to perform efficient LELs cholesterol egress.⁷

The nutrient/energy/stress sensor mTORC1 (mechanistic target of rapamycin complex 1) was also suggested to participate in cholesterol trafficking. Accumulation of cholesterol activates the PI3K (phosphatidylinositol 3-kinase) and AKT (protein kinase B) which promotes translocation of mTORC1 from the cytosol to the lysosomal limiting membrane. Active mTORC1 then suppresses autophagy, steering the cell toward cholesterol accumulation.⁸ Conversely, when mTORC1 activity is low, lysosomal exit of cholesterol raises ER cholesterol and suppresses activity of the SREBP-2 (sterol regulatory element-binding protein-2), a main transcriptional driver of cholesterol biosynthesis.⁹ In NPC1-defective cells, mTORC1 signaling is elevated suggesting that NPCs, possibly in complex with other proteins function as sensors, regulating the mTORC1-scaffolding complex via cholesterol-dependent physical interaction.

We reported that mice deficient in the transmembrane receptor, the LRP1 (LDL receptor-related protein 1) in vascular smooth muscle cells (VSMCs) rapidly develop atherosclerosis.¹⁰ LRP1 protects against atherosclerosis by integrating TGF- β (transforming growth factor β) and PDGF (platelet-derived growth factor) signaling and by stimulating the expression of a Wnt signaling protein, Wnt5a, an LRP6 ligand which we identified as an inhibitor of intracellular cholesterol accumulation.^{11,12} LRP1 deficient mouse embryonic fibroblasts expressed very low endogenous Wnt5a.¹¹ When treated with a standard adipogenic mixture containing insulin, dexamethasone, and IBMX (3-Isobutyl-1-methylxanthine),¹¹ these cells accumulated large amounts of cholesterol within perinuclear cytosolic vesicles,¹¹ whereas their treatments with recombinant Wnt5a inhibited cholesterol accumulation. Similarly, mice overexpressing Wnt5a in adipose tissue display decreased cholesterol biosynthesis and accumulation in adipocytes.¹²

Wnt5a is also expressed in murine and human atherosclerotic lesions. Its expression positively correlates with the severity of the lesions.^{13–15} Wnt5a upregulates the reverse cholesterol transport in fibroblasts, macrophages, and

VSMCs.^{11,15} However, it also stimulates endothelial cell proliferation and decreased endothelial NO production *in vitro*,¹⁶ both of which being proatherogenic. Thus, its role in atherosclerosis lacks a solid mechanistic understanding. Here, we investigated whether, as previously reported with LRP1, augmenting Wnt5a in VSMCs protects against atherosclerosis in mice. We uncovered that lysosomal composition and function are profoundly disrupted in the absence of Wnt5a. We show that Wnt5a interacts with NPCs to down regulate mTORC1 signaling and lysosomal cholesterol egress. This mechanism is essential for maintaining cholesterol homeostasis and to protect against atherosclerosis.

METHODS

Data Availability

The data and study material related to this study are available to other researchers on reasonable request. Methods are expanded in the [Supplemental Material](#).

RESULTS

Accelerated Atherosclerosis in Mice Deleted for Wnt5a in VSMCs

We achieved smooth muscle-specific Wnt5a inactivation by crossing SM22Cre transgenic mice with Wnt5a^{fllox/fllox} animals. To increase susceptibility to spontaneous atherosclerosis, these mice were crossed to LDL receptor knockout animals to generate SM22Cre+/Wnt5a^{fllox/fllox}/LDLR^{-/-} mice (smWnt5a-) and littermate controls SM22Cre-/Wnt5a^{fllox/fllox}/LDLR^{-/-} mice (smWnt5a+; [Figure S1A](#)). Only 20% of the smWnt5a- mice were born alive ([Figure S1B](#)). Death was most likely due to heart defect as suggested by histology analysis showing cardiac tissue heavily disrupted and disorganized in dead mutants (data not shown). Surviving animals looked healthy. They were subjected to a cholesterol-rich (Paigen) diet for 5 months. As evaluated by p-NF- κ B (phospho-nuclear-factor- κ B) protein levels in the liver, we did not observe any toxicity of the diet ([Figure S1E](#)). No significant difference was found in plasma cholesterol levels between mutants and controls ([Figure S1C](#)). As shown by en face analysis ([Figure 1A](#)) and quantification of the total size of the lesions in male and female animals ([Figure 1C](#)), mutant mice developed twice as much atherosclerosis as controls. Interestingly, Oil-Red-O staining and histological analysis of thoracic and abdominal aortas showed lipid accumulation within lesions and in nonlesion areas in mutant mice, whereas in controls, lipid accumulated within lesions only ([Figure 1B](#), [Figure S2B](#)), indicating that lysosomal function is impaired in the absence of Wnt5a. Thus, Wnt5a in VSMCs not only limits cholesterol accumulation, but also protects against atherosclerosis in mice, and in its absence lysosomal function is impaired.




Figure 1. Accelerated formation of atherosclerotic lesions in smWnt5a- mice.

A, Opened and Sudan IV-stained whole aortas from cholesterol-fed LDL (low-density lipoprotein) receptor deficient mice that express (smWnt5a+) or lack Wnt5a (smWnt5a-) in vascular smooth muscle cells (VSMCs). Scale bar: 0.5 cm (n=6 mice/group). **B**, Oil Red O and immunohistochemical staining of galectin-3 (GAL3), Acta-2, CD45, and Wnt5a in thoracic aortas from mice lacking Wnt5a in VSMCs or controls fed a cholesterol-rich diet. The subpanels are higher magnification images (2.5x) of the areas outlined in white. The dashed line demarcates the media. Arrow heads show macrophages staining (Gal-3 and CD45). Scale bars 50 μm (n=6 mice/group). (Continued)

Downloaded from <http://ahajournals.org> by on January 27, 2025

Macrophage Phenotypic Switching in the Absence of Wnt5a

Because in advanced atherosclerosis 40% to 50% of VSMCs express macrophage markers,¹⁷ we next evaluated whether the loss of Wnt5a affected VSMCs plasticity. In control mice, we found low expression of the macrophage marker galectin-3 (GAL3) in the media, whereas the VSMCs marker Acta-2 (α -smooth muscle actin) is expressed in the media and at the periphery of plaques. By contrast in mutant mice, we found higher expression of GAL3 (Figure 1B and 1D) and decreased expression of Acta-2 (Figure 1B) within lesions and in the media of the aortas. The endothelial cell markers CD31 expression was not affected (Figure S2A). In addition, CD45 a leukocyte/macrophage specific marker never expressed by VSMCs is expressed in the media in control mice, but not in the absence of Wnt5a indicating that these GAL3-positive cells are VSMCs in origin (Figure 1B). Together, these data suggested that Wnt5a protects against macrophage phenotypic switching of VSMCs. To confirm this in human VSMCs, we used a CRISPR/Cas9 guided nuclease approach to generate human VSMCs knockout for Wnt5a (Wnt5a^{-/-} VSMCs) and nontargeted controls (Wnt5a^{+/+} VSMCs). In human, Wnt5a gene produces 2 protein isoforms.¹⁸ Both isoforms were deleted in human VSMCs (Figure S3A and S3B). Cells were treated for 10 days with the cholesterol accumulation cocktail we previously reported, that consisted of a standard adipogenic mixture containing insulin, dexamethasone, IBMX, and the PPAR γ agonist rosiglitazone.¹¹ In human Wnt5a^{-/-} VSMCs western blot analysis revealed increased expression of the macrophage marker GAL3 (Figure S1D) and decreased expression of the VSMCs marker myocardin (Figure 1F) upon cholesterol loading. In addition, mRNA expressions of the KLF4 (Kruppel-like transcription factor 4), a transcription factor known to silence VSMC marker genes,¹⁷ and to promote formation of cholesterol-induced VSMC-derived foam cells strongly increased in untreated cells and upon cholesterol loading in the absence of Wnt5a (Figure 1F). Thus, Wnt5 contribute to protect against macrophage phenotypic switching of VSMCs in vitro and during vascular injury.

Lipid Accumulation in Wnt5a^{-/-} Human VSMCs

We next tested whether Wnt5a protects against cholesterol accumulation in human VSMCs. When cultured

in normal conditions, total cholesterol increased slightly but non significantly in human Wnt5a^{-/-} VSMCs when compared with controls (Figure 2A). However, cholesteryl ester (CE) concentrations were significantly higher in Wnt5a^{-/-} VSMCs (Figure 2B). Upon cholesterol loading, Wnt5a protein levels increased over time in wild-type human VSMCs (Figure S3C) and neutral lipids accumulated modestly as shown by Oil-Red-O staining and microscopy analysis (Figure S3E). By contrast, neutral lipids strongly accumulated in Wnt5a^{-/-} VSMCs compared with controls, whereas their treatment with human recombinant Wnt5a rescued neutral lipids accumulation back to control levels (Figure 2D). At day 10, CE level was strongly increased in Wnt5a^{-/-} VSMCs (Figure 2B), and total cholesterol concentration was elevated as well (Figure 2A). In both treated and untreated cells, triglyceride levels (Figure S3D) as well as transcript levels of the DGAT2 (acyl-CoA:diacylglycerol acyltransferase 2) enzyme that is required for triglyceride synthesis in adipocytes,¹⁹ were similar in Wnt5a^{-/-} VSMCs and controls (Figure S3G). To confirm cholesterol accumulation in Wnt5a^{-/-} VSMCs we used oxidized low-density lipoproteins (ox-LDL) as a lipoprotein vehicle for cholesterol loading. Previous reports indicated that upon ox-LDL treatments, RAW264.7 macrophages and VSMCs knockdown for Wnt5a accumulated cholesterol.¹⁵ In agreement, we now show that when treated with ox-LDL for 12 days, Wnt5a^{-/-} human VSMCs accumulated more neutral lipids than controls as evaluated by Oil Red O staining and microscopic analysis (Figure S4A).

We next tested whether loss of Wnt5a activates cholesterol biosynthesis genes in human VSMCs. We found that mRNA levels of SREBP-2, and mRNA and protein levels of its target gene HMG CoA reductase were increased in the absence of Wnt5a in untreated and treated human VSMCs (Figure 2C, Figure S3F). Thus, loss of Wnt5a in human VSMCs constitutively increased the SREBP-2 pathway. These data are consistent with our previous report indicating that overexpression of Wnt5a in adipose tissue of mice decreased SREBP-2 activity and cholesterol biosynthesis.¹²

Absence of Wnt5a Is Associated With Cholesterol Accumulation in Late Endosomes

We next analyzed in which compartment cholesterol accumulated in the absence of Wnt5a. Electron microscopic analysis of treated human Wnt5a^{-/-} VSMCs

Figure 1 Continued. **C**, Quantitative analysis of atherosclerotic lesion size in whole aortas from cholesterol-fed mice. The plot shows individual values with mean \pm SEM (n=4 mice/group). **D**, Relative protein levels of Gal-3 in mutant and control mice. The plot shows individual scatter with mean \pm SEM (n=4 mice/group). **E**, Representative electron micrographs show enlarged late endosomes/lysosomes (LEs) in VSMCs from thoracic aortas of mutants vs controls. Endothelial cells, nuclei, and lipid droplets are indicated (EC, N, LD). Scale bar **top**: 5 μ m, **bottom**: 2 μ m (n=6 mice/group). **F**, Relative transcript levels of indicated genes in human VSMCs Wnt5a^{-/-} and Wnt5a^{+/+} untreated (D0) and upon cholesterol loading (D10) taking GAPDH as internal control. The plot shows individual scatter with mean \pm SEM (myocardin n=5 and Klf4 n=5). Statistical significance was assessed by Shapiro-wilk to test normality and Mann-Whitney test for **C**, **D**, and **F**. **C**, $P=0.015$; **D**, $P=0.028$ indicate significance relative to smWnt5a+; **F**, $P=0.015$ and $P=0.0079$ indicate significance relative to Wnt5a^{+/+}. ns indicates not significant.




Figure 2. Cholesterol accumulation and altered lysosomal function in human vascular smooth muscle cells (VSMCs) $Wnt5a^{-/-}$. Human VSMCs $Wnt5a^{-/-}$ and $Wnt5a^{+/+}$ were analyzed untreated (D0) or upon treatment with a cholesterol accumulation protocol during 10 days (D10). **A**, Quantification of total cholesterol (n=5 D0 $Wnt5a^{+/+}$, n=7 D0 $Wnt5a^{-/-}$, n=8 D10 $Wnt5a^{+/+}$, n=10 D10 $Wnt5a^{-/-}$). **B**, Quantification of cholesteryl esters (n=5 D0 $Wnt5a^{+/+}$, n=7 D0 $Wnt5a^{-/-}$, n=8 D10 $Wnt5a^{+/+}$, n=10 D10 $Wnt5a^{-/-}$). **C**, Relative transcript levels of SREBP-2 (sterol regulatory element-binding protein 2; n=4 D0 and D10) and HMG CoA reductase (n=9 D0 $Wnt5a^{+/+}$, n=8 D10 $Wnt5a^{-/-}$, n=4 D10 $Wnt5a^{+/+}$ and n=4 D10 $Wnt5a^{-/-}$) taking GAPDH 18S as internal control. **D**, Oil Red O staining upon treatment for cholesterol accumulation during 10 days in presence of human recombinant Wnt5a (+rWnt5a) or BSA (+BSA) in human $Wnt5a^{-/-}$ VSMCs (n=3). **E**, Representative electron micrographs show presence of large lipofuscin containing vesicles (Lipo), lipid droplets (LD), and large late endosomes/lysosomes (LELs) in human $Wnt5a^{-/-}$ VSMCs vs controls. Scale bar: 1 μ m (n=3). **A**, **B**, and **C** show individual value along with mean \pm SEM. Statistical significance was assessed by Shapiro-wilk to test normality, unpaired *t* test for **A** and **B**, and Mann-Whitney test for **C**. **A**, $P=0.034$; **B**, $P=0.038$; **C**, $P=0.028$ and $P=0.028$ for SREBP-2 and $P=0.034$ and $P=0.028$ for HMG CoA reductase indicate significance relative to $Wnt5a^{+/+}$. ns indicates not significant.

revealed accumulation of lipid droplets, accumulation of numerous very LELs containing cholesterol crystals, and accumulation of very large electron-dense cytoplasmic lipofuscin containing vesicles (Figure 2E, Figure S4B) that were comparable in shape and size to those seen in the heart and in the aortas of *smWnt5a*⁻ mice (Figure S2B). To determinate whether lipofuscin accumulated in endosomes, we used confocal imaging and the late endosomes marker Lamp1 (lysosome-associated membrane protein 1). We found that the lipofuscin-enriched large cytosolic vesicles that accumulate in *Wnt5a*⁻ VSMCs are Lamp1 positive LELs (Figure S4C).

The increased in cholesterol and CE levels upon treatment with the cholesterol accumulation cocktail, together with the alteration of the endosomal compartments suggested that in the absence of Wnt5a, cholesterol does not reach the ER. To test this, we quantified cholesterol inside cells with a specific fluorescent probe: the D4 fragment of perfringolysin O fused to GFP (GFP-D4). This probe can be efficiently used to label sterols in the PM or in subcellular structures such as endosomes.²⁰ We found more cholesterol-enriched cytosolic vesicles in *Wnt5a*⁻ VSMCs as shown by immunofluorescence (Figure 3A) and by quantification of GFP-D4 positive vesicles (Figure 3B). Moreover, co-labelling between Lamp1 and the GFP-D4 probe showed an overlap of both signals indicating that cholesterol accumulates in endosomal compartments (Figure 3A). Quantification of colocalization between GFP-D4 and Lamp1 show that there is three times more Lamp1+/GFP-D4+ overlapping vesicles in VSMCs lacking Wnt5a compared with controls upon cholesterol accumulation (Figure 3C). In addition, immunofluorescence analysis of the LELs marker Lamp1 indicated that Lamp1 protein levels are substantially higher in cholesterol-fed mutant cells compared with controls (Figure 3A). A time course experiment revealed that, whereas in control cells Lamp1 protein expression increased modestly during the course of cholesterol accumulation protocol, its expression reached higher levels in VSMCs lacking Wnt5a (Figure 3D, Figure S4D). Analysis of acidic compartments using LysoTracker reagent showed numerous and bigger acidic compartments in the absence of Wnt5a (Figure 3E, F). Similarly, upon treatments with the ox-LDL protocol immunohistochemistry experiments revealed numerous and bigger Lamp1 positive compartments in the absence of Wnt5a (Figure S4A). Thus, in the absence of Wnt5a cholesterol accumulated in multi-vesicular inclusions and enlarged LELs.

Absence of Wnt5a Blocks Cholesterol Egress From LELs

Because lysosome-derived cholesterol correlates closely with ER cholesterol, we indirectly assessed ER cholesterol levels by measuring ACAT1 (acyl-CoA

cholesterolacyltransferase 1) expression and activity. ACAT1 is an ER resident enzyme that converts free cholesterol to cholesteryl ester. Its activity is controlled by cholesterol availability in the ER membrane.²¹ In untreated cells, ACAT1 protein levels were markedly decreased in human *Wnt5a*⁻ VSMCs compared with controls (Figure 4A and 4B), indicative of low ER cholesterol. In addition, ACAT1 activity (Figure 4C) was almost undetectable in human *Wnt5a*⁻ VSMCs untreated or treated for cholesterol accumulation (Figure 4C). LAL is the enzyme that hydrolyzes CE in lysosomes. Its activity in human VSMCs is markedly lower when compared with macrophages and this represent a mechanism for VSMCs foam cell formation.²² To rule out the possibility that the CE accumulation observed in *Wnt5a*⁻ VSMCs is due to decreased lysosomal hydrolysis of CE, we measured its expression. We found the LAL protein levels almost undetectable in the absence of Wnt5a (Figure 4D). Thus, the loss of Wnt5a may reduce its capacity to catabolize atherogenic lipoproteins and CE. This results in the accumulation of cholesterol and CE in Lamp1 positive compartments at the expense of the ER.

Localization of Wnt5a in LEL Particles

To determine how Wnt5a enter the endocytic compartment we postulate that, as previously described in the cerebrospinal fluid,²³ Wnt5a associates with LDL particles. For this, recombinant Wnt5a was incubated with either 0.6% CHAPS or human LDL lipoprotein fraction for 4 hours at 37°C and filtered through 100 kDa cut-off protein concentrator. When recombinant Wnt5a was incubated with LDL, Western blot confirmed presence of Wnt5a only in >100 kDa fraction but not in the <100 kDa fraction indicating that recombinant Wnt5a localized on LDL particles and forms a high-molecular weight complex (Figure 4E).

Next, we hypothesize that Wnt5a might be internalized together with LDLs in LELs. For this, we used a subcellular fractionation approach to isolate LELs from human VSMCs as outlined in Figure 4F. The cell homogenate (fraction A) was subjected to 2 centrifugation steps to separate LELs membranes from other organelles (Figure 4F, left panel). Centrifugation of the homogenate at 3000g separates unbroken cells and the heaviest membranes. As assessed by immunoblot analysis of the fractions (Figure 4F, right panel), a plasma membrane protein marker (caveolin-1) and an ER membrane protein marker (ER-72) are mostly retained in the 3000g pellet (fraction B), whereas significant amounts of all other membrane markers, including the LEL protein marker Lamp1 were present in the 3000g supernatant (fraction C). Fraction C was then subjected to centrifugation through a discontinuous sucrose gradient and yielded 2 distinct bands of membranes (fractions D and E). Fraction E contained ER membrane protein marker, whereas the LEL membrane




Figure 3. Cholesterol accumulated in Lamp1 positive cytosolic vesicles in human *Wnt5a*^{-/-} vascular smooth muscle cells (VSMCs).

A, Cholesterol accumulation in human VSMCs *Wnt5a*^{-/-} and controls labeled with anti-Lamp1 antibodies (red) a LELs marker, and with the fluorescent cholesterol probe GFP (D4 fragment of perfringolysin O fused to GFP)-D4 (green). Nuclei were stained in blue with Dapi. Higher magnification (2.5 \times) images of the areas outlined in white are shown in the **right**. Scale bar 5 μ m (n=4). **B**, Quantification of GFP-D4 positive vesicles/cells in human VSMCs *Wnt5a*^{-/-} and controls (n=8 *Wnt5a*^{+/+}, n=4 *Wnt5a*^{-/-}). **C**, Quantification of confocal analysis shows increased colocalization between GFP-D4 (cholesterol) and Lamp1 (LELs) in human VSMCs *Wnt5a*^{-/-} compared with controls (n=9 *Wnt5a*^{+/+}, n=6 *Wnt5a*^{-/-}). **D**, A representative immunoblot shows the protein expression of Lamp1 during the course of cholesterol accumulation (n=3). **E**, Human VSMCs *Wnt5a*^{-/-} and controls were treated with 50 nmol/L lysosome-tracker red. Higher magnification (2.5 \times) images of the area outlined in white are shown on the right. Scale bar: 6 μ m (n=4). **F**, Quantification of lysosome-tracker red positive vesicles (n=6 *Wnt5a*^{+/+} and *Wnt5a*^{-/-}). **B**, **C**, and **F** show individual value along with mean \pm SEM. For **C**, statistical significance was assessed by Shapiro-wilk to test normality followed by an unpaired *t* test. For **B** and **F**, statistical significance was assessed by Shapiro-wilk to test normality followed by a Mann-Whitney test. **B**, $P=0.004$; **C**, $P=0.035$; and **F**, $P=0.008$ indicate significance relative to *Wnt5a*^{+/+}.




Figure 4. Decreased concentrations of endoplasmic reticulum (ER) membrane cholesterol in human vascular smooth muscle cells (VSMCs) Wnt5a^{-/-}.

A, A representative immunoblot shows the protein expression of ACAT1 (acyl-CoA cholesterolacyltransferase 1) and **(B)** its quantification in human Wnt5a^{-/-} VSMCs vs controls upon cholesterol accumulation protocol calculated as relative to GAPDH expression (n=3). **C**, ACAT1 activity in human Wnt5a^{-/-} VSMCs vs controls before (D0) and upon (D10) cholesterol accumulation treatments during 10 days (n=4). **D**, A representative immunoblot shows the protein expression of the LAL (lysosomal acid lipase; n=4). **E**, Wnt5a was incubated with human LDL (low-density lipoprotein) particles as described in the method section. A representative immunoblot shows the presence of (Continued)

protein Lamp1 was essentially present in fraction D (Figure 4F, right panel). Interestingly, Wnt5a was present in fractions C and D, indicative of its localization in LELs. In addition, cholesterol in the fraction enriched in ER membranes (the heavy fraction E) is decreased in mutant cells compared with controls, whereas cholesterol in the fraction enriched in LEL membranes (the light fraction D) is increased (Figure 4G). Thus, Wnt5a is directly associated with the LEL compartments, and in the absence of Wnt5a cholesterol is retained in LEL fractions.

Wnt5a Down Regulates the mTORC1 Pathway

LELs have emerged as the cellular site where the master growth regulator and nutrient sensor, mTORC1 is activated. Given that high mTORC1 activity coincides with low ER cholesterol, high ACAT-1 expression, and activation of SREBP-2,⁹ we assessed mTORC1 activation in Wnt5a^{-/-} VSMCs. We found a robust increase of p-mTORC1 and its downstream targets, p-P70S6 and p-4EBP-1, and of p-Akt (phospho Akt) in the absence of Wnt5a in human VSMCs (Figure 5A). During the course of cholesterol accumulation, p-mTORC1 remains highly activated in the absence of Wnt5a (Figure S5A). Interestingly, rescue experiments show that treatment of human Wnt5a^{-/-} VSMCs with conditioned media enriched in Wnt5a (Figure S5B) decreased the p-mTORC1 target gene p-P70S6 to control levels, whereas mock media had no effect (Figure 5B). Similarly, treatment with recombinant human Wnt5a decreased p-mTORC1 in human Wnt5a^{-/-} VSMCs back to control levels (Figure 5D). Treatment with Everolimus, an mTORC1 inhibitor, decreased p-mTORC1 and p-Akt in human Wnt5a^{-/-} VSMCs to control levels (Figure S5C). Everolimus treatment decreased cholesterol accumulation in the CD63 positive compartments in human Wnt5a^{-/-} VSMCs (Figure 5E). Similarly, Oil-red-o staining showed that neural lipids and cholesterol are reduced upon Everolimus treatments (Figure S5D), and GC-MS quantification showed that Everolimus treatments decreased total cholesterol and cholesteryl esters in Wnt5a^{-/-} VSMCs, however, without reaching significance (Figure S5E). Thus, loss of Wnt5a coincides with high mTORC1 activity, and cholesterol accumulation.

Changes in cholesterol levels can alter the activity of signaling pathways originating at the plasma membrane involving PI3K that can activate mTORC1.²⁴ To test whether Wnt5a acts upstream of mTORC1 or directly

on mTORC1 we measured PI3K activation in human VSMCs. We found the p85 subunit of PI3K to be highly tyrosine-phosphorylated in human Wnt5a^{-/-} VSMCs compared with controls (Figure 5C) and treatment with the PI3K inhibitor Wortmannin inhibited mTORC1 activation in human Wnt5a^{-/-} VSMCs (Figure S5F). Marked activation of the Akt/mTORC1 pathway in the absence of Wnt5a was also observed in vivo with increased protein levels of p-Akt, p-P70 S6, and p-4EBP1 (phospho-eukaryotic translation initiation factor 4E-binding protein 1) in aortas from smWnt5a- mice compared with controls (Figure 5F and 5G). Overall, these data suggest that Wnt5a promotes LEL cholesterol egress by down regulating the PI3K/Akt/mTORC1 pathway in human VSMCs.

Absence of Wnt5a Activated the Calcineurin-Dependent TFEB Nuclear Relocalization

The MiT/TFE transcription factor family member, TFEB (transcription factor EB) drives the expansion of the lysosomal compartment and lysosomal biogenesis.²⁵ To assess whether Wnt5a modulates TFEB activity, we measured mRNA levels of its target genes in human Wnt5a^{-/-} VSMCs and controls. Results show an increase in SGSH (N-sulphoglucosamine sulphohydrolase), CTSE (cathepsin F), and TPP1 (tripeptidyl peptidase 1) mRNA levels in untreated mutant cells, and an increase in ATP6V (V-type proton ATPase) mRNA levels in cholesterol-fed mutant cells (Figure S4E and S4F) suggesting that TFEB is activated in the absence of Wnt5a. In agreement, we found that upon cholesterol accumulation TFEB translocated to the nucleus in the absence of Wnt5a (Figure 6A and 6B). Conditions that promote TFEB nuclear translocation, such as starvation and lysosomal stress, dephosphorylated TFEB, and one phosphatase involved in TFEB dephosphorylation has been identified as calcineurin.²⁶ Conversely, mTORC1-mediated phosphorylation of TFEB at serine residues Ser142 and Ser211 promotes its cytoplasmic localization.²⁷ To test whether in the absence of Wnt5a, TFEB dephosphorylation is driven by calcineurin, human Wnt5a^{-/-} VSMCs were treated with the calcineurin inhibitor FK506 upon cholesterol accumulation. FK506 treatments relocalized TFEB to the cytosol (Figure 6A and 6B). In addition, using the Fura-8 fluorescent calcium marker, we found that basal intracellular calcium levels are increased in human Wnt5a^{-/-} VSMCs compared with controls (Figure 6C), an

Figure 4 Continued. Wnt5a in the <100 kDa fraction when incubated with CHAPS, or in >100 kDa apoB positive fraction when incubated with human LDL (n=3). **F**, Purification of ER and LELs, purification of mitochondria, plasma and early endosomal membranes. **Left** is showing the diagram of LELs membrane fractionation scheme. **A–E** denote major fractions recovered and analyzed by Western blot. **Right**, VSMCs were treated according to the fractionation scheme and as described in the method section. Aliquots representing equal volumes of each fraction (**A–E**) were subjected to immunoblot analysis for the indicated organelles markers and Wnt5a. **G**, Quantification of cholesterol concentrations in heavy (**E**, ER) and light (**D**, lysosomes) membrane fractions from human Wnt5a^{-/-} VSMCs and Wnt5a^{+/+} (n=8 and n=7 Wnt5a^{-/-} heavy fractions). **B**, **C**, and **G** show individual data along with mean±SEM. For **B**, **C**, and **G**, data were analyzed using a Shapiro-Wilk test to assess normality followed by Mann-Whitney test. For **B**, $P=0.028$; **C**, $P=0.028$; and **G**, $P=0.049$ and $P=0.040$ indicate significance relative to Wnt5a^{+/+}.




Figure 5. Wnt5a decreases mTORC1 (mechanistic target of rapamycin complex 1) activity.

A, Representative immunoblot of the indicated genes in human Wnt5a^{-/-} and control vascular smooth muscle cells (VSMCs) upon cholesterol accumulation (n=3). **B**, Western blot analysis showing p-P70 S6 (Thr389) expressions in Wnt5a^{-/-} VSMCs untreated (UT) or treated with conditioned medium enriched in Wnt5a (CM Wnt5a⁺) or mock medium (CM Wnt5a⁻) as described in the method section (n=3). **C**, Immunoprecipitation in Wnt5a^{-/-} VSMCs and controls of the PI3K (phosphatidylinositol 3-kinase)-p85 subunit followed with immunoblotting with anti-phosphotyrosine (4G10) antibodies show the tyrosine-phosphorylated form of p85 (p-p85) in the absence of Wnt5a. NI indicates nonimmune antibodies; and beads, empty beads. Input is from whole cell lysate (n=3). **D**, A representative immunoblot shows p-mTORC1, total mTORC1, p-GSK3, and Wnt5a expressions in Wnt5a^{-/-} VSMCs (-/-) and control cells (+/+) untreated (UT) or treated (rWnt5a) with human recombinant Wnt5a (n=3). **E**, Confocal analysis showing colocalization (yellow) between GFP-D4 (D4 fragment of perfringolysin O fused to GFP) positive vesicle (cholesterol, green) and CD63 (LELs, red) in untreated (UT) and Everolimus (30 nmol/L) treated Wnt5a^{-/-} VSMCs. Scale bars are 5 μ m (n=3). **F**, Western blot analysis and **G** quantification of relative protein levels of the indicated genes in aortas from control (smWnt5a⁺) and mutant (smWnt5a⁻) mice (n=4 mice/group). **G**, Individual data along with mean \pm SEM. Data were analyzed using Shapiro-wilk followed by Mann-Whitney test. **G**, P=0.028 indicate significance relative to smWnt5a⁺ mice.




Figure 6. Calcineurin-dependent nuclear relocalization of TFEB (transcription factor EB) in *Wnt5a*^{-/-} vascular smooth muscle cells (VSMCs).

A, Nuclear localization of TFEB-GFP and its quantification **B**, in human VSMC *Wnt5a*^{-/-} and controls untreated (UT) or treated with the calcineurin inhibitor FK506 (+FK506). Scale bars are 5 μ m ($n=17$ *Wnt5a*^{+/+} UT, $n=14$ *Wnt5a*^{-/-} UT, $n=12$ *Wnt5a*^{+/+} +FK506 and $n=23$ *Wnt5a*^{+/+} +FK506). **C**, Intracellular calcium quantification in human VSMC *Wnt5a*^{-/-} and controls ($n=184$ *Wnt5a*^{+/+} and $n=187$ *Wnt5a*^{-/-}). **B** and **C**, Individual data with mean \pm SEM. Statistical analysis was done using Shapiro-Wilk test followed by Mann-Whitney test. **B** and **C**, $P < 0.0001$ indicate significance relative to *Wnt5a*^{+/+}.

event known to activate calcineurin.²⁶ Collectively, these data suggest that in the absence of *Wnt5a*, activation of calcineurin by Ca^{2+} release from intracellular stores overrides the mTORC1 signal and causes TFEB to go to the nucleus.

Wnt5a Physically Interacts With NPC Proteins

On the basis of our results indicating that absence of *Wnt5a* interferes with cholesterol trafficking in LELs, we examined whether the *Wnt5a* sequence contains cholesterol-binding motifs. We identified 5 putative cholesterol recognition amino acid consensus motifs that we named D1-5 (Figure 7A). The consensus cholesterol recognition amino acid consensus motifs sequence (L/V)-X₁₋₅-Y-X₁₋₅-(K/R) has been described in transmembrane domains of cholesterol-regulated proteins such as caveolins, the

Ca^{2+} channel Orai 1, and SLC38A9 within the mTORC1 scaffolding complex.^{28,29} However, cholesterol-binding domains have also been identified outside the membrane-spanning regions of soluble proteins such as α -synuclein.³⁰ Using the 3-dimensional structure of *Wnt8* another *Wnt* ligand that is 33 % homologous to *Wnt5a* as a template,³¹ we modeled in silico the 3-dimensional structure of *Wnt5a* and the subsequent localization of cholesterol recognition amino acid consensus motifs (Figure 7A). Cholesterol recognition amino acid consensus motifs D2, D3, and D4 were found at the periphery of the 3-dimensional structure of *Wnt5a*, accessible to cholesterol, and thus in a biologically relevant position (Figure 7A). We next investigated the ability of *Wnt5a* to bind cholesterol in liposomes reconstituted with phosphatidylcholine, enriched with cholesterol (PC+Chol) or not (PC). Co-sedimentation assays revealed that *Wnt5a*




Figure 7. Wnt5a interacts with NPC (Niemann–Pick C) proteins and cholesterol-enriched liposomes.

A, Predictive 3D structure of Wnt5a and localization of the cholesterol recognition amino acid consensus (CARC) motifs. The arrow is showing localization for the palmitoleic acid lipid group. The essential Arg/Lys within the CARC motifs are boxed in dark. **B**, Conditioned medium enriched in Wnt5a (CM Wnt5a+) or mock medium (CM Wnt5a-) were incubated with liposomes containing 80%–85% phosphatidylcholine (PC) or 80%–85% phosphatidylcholine+5% cholesterol (PC+Chol). Liposomes were sedimented, washed, and analyzed by immunoblotting for the presence of Wnt5a. Supernatant (S) and pellets (P) were loaded (n=3). **C**, Relative transcript levels of NPC1 in human VSMC Wnt5a^{-/-} and controls untreated (D0) or treated for cholesterol accumulation during 10 days (D10). GAPDH is taken as internal control. The plot shows individual values with mean±SEM (n=4). **D**, HEK293 cells were co-transfected with either pCMV-Wnt5a and pCMV-NPC1-EGFP or pCMV-Wnt5a and pCMV-NPC2-Tag expression plasmids. Immunoprecipitations of NPC1 (lane 3) or with NPC2 (lane 4), followed with immunoblotting with Wnt5a show that Wnt5a interacts with both NPC proteins. NI indicates nonimmune antibodies; and beads, empty beads (n=3). **E**, In vitro direct binding between His-Wnt5a and Flag-NPC2. Recombinant Flag-NPC2 was incubated with Flag-NTA agarose beads allowing NPC2 to bind to the beads. (Continued)

did not pellet with PC-containing liposomes (Figure 7B, lane 4). However, the inclusion of cholesterol in PC-containing liposomes enhanced their association with Wnt5a (Figure 7B, lane 2) indicating that Wnt5a binds to cholesterol-enriched membranes.

We next tested whether Wnt5a interacts with NPC1 and NPC2 and measured their mRNA levels in the absence of Wnt5a. We found that mRNA levels of NPC1 increased by 2-3X in Wnt5a^{-/-} VSMCs compared with controls (Figure 7C), whereas NPC2 levels remain unchanged (not shown). Moreover, when co-expressed in HEK293T cells, we found that Wnt5a co-immunoprecipitated with NPC1 and NPC2 (Figure 7D). The immunoprecipitate does not contain the late endosome marker, Lamp1 suggesting that the interaction between Wnt5a and NPC proteins (Figure 7D) is not a mere co-precipitation of proteins present in the same membrane complex. We also found that Wnt5a co-immunoprecipitated with NPC1 and NPC2 using mass spectrometry analysis of the interactome (Table S4). We further tested the specific interaction between Wnt5a and NPC2 using in vitro binding experiments and recombinant proteins. We found that purified recombinant Wnt5a physically interacts with recombinant NPC2 (Figure 7E). This in vitro interaction demonstrates a direct binding between these 2 proteins and indicates that Wnt5a interacts with the lysosomal cholesterol export machinery.

The U18666A compound is a cholesterol transport-inhibiting agent, used to mimic NPC1 deficiency, which causes the accumulation of cholesterol in cells. Interestingly, we found that in human Wnt5a^{-/-} VSMCs, Wnt5a-enriched conditioned medium decreased mTORC1 activity induced by U18666A treatments, whereas mock medium had no significant effect (Figure 7F). Thus, loss of Wnt5a or NPC1 inhibition have similar effects on mTORC1 activation and cholesterol trafficking, and Wnt5a counteracts the activation of mTORC1 mediated by U18666A-induced NPC1 inhibition.

DISCUSSION

How cholesterol moves between cellular membranes in mammalian cells is not understood? Our study identifies Wnt5a as essential for lysosomal cholesterol egress.

Wnt5a act as intercellular signaling molecules but is lipid modified. Therefore, its intercellular transport requires extracellular carriers.¹ Here, we report that Wnt5a enter the endocytic compartment through binding

to the LDL lipoparticles. In human or murine VSMCs Wnt5a^{-/-} the LAL expression is very low and the cholesterol hydrolysis processes are overwhelmed. Cholesterol and CE accumulated in enlarged LELs and this coincide with high mTORC1 activity. Within LELs, Wnt5a interacts with NPC proteins, down regulates mTORC1, and promotes cholesterol egress to the ER and lysosomal function. Thus, Wnt5a appears to sense sterol abundance and translates cholesterol levels directly into mTORC1 activity, in parallel with the NPC proteins (Figure 7G).

We previously reported that preadipocytes and mouse embryonic fibroblasts LRP1^{-/-}, that expressed low endogenous Wnt5a, did not form adipocytes when submitted to an adipogenic cocktail. Instead, they accumulated cholesterol and CE.^{11,12} Fatty acid synthesis is impaired, and triglycerides levels remained unchanged.^{11,12} Here, we found essentially the same changes in human Wnt5a^{-/-} VSMCs.

In agreement with our data, previous reports showed that siRNA-mediated silencing of Wnt5a in both macrophages and VSMC significantly increased cholesterol content upon exposure to ox-LDL.¹⁵ In addition, Wnt signaling enhanced the macrophage response to inflammation resolving factors.⁴ However, it was also reported that siRNA-mediated silencing of Wnt5a attenuated atherosclerosis potentially through inhibiting the proinflammatory NF- κ B and MAPK (mitogen-activated protein kinase) signaling pathways³² suggesting that immune function and endothelial cell function are affected by the loss of VSMC Wnt5a. Here, we did not find any significant difference in the expression of the endothelial cell markers CD31 in the absence of Wnt5a in VSMC. In addition, an alteration of the immune or endothelial cell function would be against the atheroprotective effects of LRP6 observed in human. These apparent discrepancies may be related to the receptor availability of Wnt5a that can both activate canonical and noncanonical pathways. For instance, the noncanonical Wnt5a/CaMKII (Ca²⁺/calmodulin dependent protein kinases II) pathway contributes to the inflammatory response of human macrophages,¹⁶ whereas the Wnt5a/ β -catenin or the LRP6 pathway prevent against cholesterol accumulation and atherosclerosis.^{3,11}

As cholesterol accumulated, Wnt5a protein levels increased in human Wnt5a^{+/+} VSMCs (Figure S3C) and, as described by others and we, also increased in the aortic wall during atherosclerosis.¹³⁻¹⁵ However, the correlation between its expression and the accumulation of cholesterol in the arterial wall does not mean there is

Figure 7 Continued. Then 10 μ g of human recombinant Wnt5a were added and incubated at room temperature. Complex of proteins was then eluted with a glycine buffer and analyzed by Western blotting (n=4). **F**, Effects of the U18666A compound on the phosphorylation of mTORC1 (mechanistic target of rapamycin complex 1) in presence of control medium (CM Wnt5a-) or Wnt5a enriched condition medium (CM Wnt5a+) in human VSMC Wnt5a^{-/-} and controls (n=3). **G**, Wnt5a is required for proper lysosomal functions. It promotes cholesterol egress from late endosomes to ER through inhibition of p-mTORC1. In late endosomes/lysosomes (LELs), upon binding to NPC2 and cholesterol, Wnt5a might facilitate cholesterol transfer to NPC1 and to the ER membrane. This suppresses SREBP-2 (sterol regulatory element-binding protein-2) activity, limits cholesterol accumulation in VSMCs, and protect against atherosclerosis. Statistical analysis was done using a Shapiro-Wilk test followed by 1-way ANOVA and Tukey post hoc analysis for **C**, **C**, $P=0.023$ and $P=0.088$ indicate significance relative to Wnt5a^{+/+}.

a causal effect. Here, by deleting *Wnt5a* in the VSMCs of mice we directly addressed its protective role against cholesterol-induced atherosclerosis. In addition, the absence of *Wnt5a* activated the PI3K/mTORC1 pathway. PI3K integrates signal from cellular growth and functions as a checkpoint for slowing or preventing degenerative disorders of the vascular wall. For instance, LRP1 mutant mice containing in VSMCs a mutated PDGF receptor- β kinase domain that disrupted the PI3K activation are protected against atherosclerosis.³³

Under nutrient-rich conditions, through mTORC1-mediated phosphorylation, TFEB is retained inactive in the cytoplasm. Upon starvation or under conditions of lysosomal dysfunction, TFEB is dephosphorylated by calcineurin and rapidly translocates to the nucleus to activate the transcription of its target genes. In human *Wnt5a*^{-/-} VSMCs, we found TFEB in the nucleus and increased intracellular Ca²⁺ levels, an event known to activate calcineurin.²⁶ In the absence of *Wnt5a*, activated calcineurin overrides the mTORC1 signal and causes TFEB to go to the nucleus where it drives the expansion of the lysosomal compartment allowing cholesterol to accumulate.

In conclusion, the work presented here identifies a previously unrecognized crucial role for *Wnt5a* in controlling cholesterol homeostasis. The absence of *Wnt5a* down regulates LAL and this leads to the accumulation of CE in enlarged LELs. Accumulation of cholesterol activates the nutrient-dependent mTORC1 axis at the surface of LELs, whereas *Wnt5a* interacts with NPC proteins to communicate lysosomal sterol abundance to the mTORC1 complex. This is required for cholesterol transport across the lysosomal membrane and its trafficking to the ER for subsequent redistribution and export via ATP binding cassette (ABC) transporters. Through this mechanism, *Wnt5a* limits cholesterol accumulation in VSMCs and protects against atherosclerosis in mice.

ARTICLE INFORMATION

Received January 14, 2021; revision received December 8, 2021; accepted December 9, 2021.

Affiliations

UMR-S INSERM 1109 (S.A., M.L., A.I., R.L.M., J.T., P.B.), Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC) (F.A., C.T., M.O.-A., R.S.-R., J.B.), Institut Pluridisciplinaire Hubert Curien (C.S., C.M.), and CNRS, UMR 7178 (D.J.-D.), University of Strasbourg, France. Institute of Metabolic and Cardiovascular Diseases, I2MC, INSERM, UMR, 1048, 1 avenue du Professeur Jean Poulhès, France (S.N., L.O.M., X.C.). Department of Molecular Genetics, University of Texas Southwestern Medical Center at Dallas (J.H.).

Acknowledgments

We are grateful to Toshihide Kobayashi, Nario Tomishige, and Motohide Murate (UMR CNRS 7021, University of Strasbourg) for helpful suggestions and EM analysis, Russell Debose-Boyd (UT Southwestern Medical Center at Dallas) for antibodies against HMG CoA reductase, and the Mouse Clinical Institute of Strasbourg (UMR CNRS 7021) for technical assistance.

Sources of Funding

This work was supported by grants from Fédération Française de Cardiologie, and the Fondation de France (No. 00107044). R. Silva-Rojas was funded by

Fondation Recherche Médicale doctoral fellowship (FRM, PLP20170939073). J. Herz was supported by grants from the National Heart, Lung, and Blood Institute (R37 HL063762), NIA (RF AG053391), the National Institute of Neurological Disorders and Stroke and NIA (R01 NS093382), BrightFocus A2016396S, the Bluefield Project to Cure FTD, and a Harrington Scholar Innovator Award (2019).

Disclosures

None.

Supplemental Materials

Expanded Materials and Methods

Figures S1–S5

Table S4

REFERENCES

- Logan CY, Nusse R. The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol*. 2004;20:781–810. doi: 10.1146/annurev.cellbio.20.010403.113126
- Go GW, Srivastava R, Hernandez-Ono A, Gang G, Smith SB, Booth CJ, Ginsberg HN, Mani A. The combined hyperlipidemia caused by impaired Wnt-LRP6 signaling is reversed by Wnt3a rescue. *Cell Metab*. 2014;19:209–220. doi: 10.1016/j.cmet.2013.11.023
- Mani A, Radhakrishnan J, Wang H, Mani A, Mani MA, Nelson-Williams C, Carew KS, Mane S, Najmabadi H, Wu D, et al. LRP6 mutation in a family with early coronary disease and metabolic risk factors. *Science*. 2007;315:1278–1282. doi: 10.1126/science.1136370
- Weinstock A, Rahman K, Yaacov O, Nishi H, Menon P, Nikain CA, Garabedian ML, Pena S, Akbar N, Sansbury BE, et al. Wnt signaling enhances macrophage responses to IL-4 and promotes resolution of atherosclerosis. *Elife*. 2021;10:e67932. doi: 10.7554/eLife.67932
- Maxfield FR, Tabas I. Role of cholesterol and lipid organization in disease. *Nature*. 2005;438:612–621. doi: 10.1038/nature04399
- Zhang JR, Coleman T, Langmade SJ, Scherrer DE, Lane L, Lanier MH, Feng C, Sands MS, Schaffer JE, Semenkovich CF, et al. Niemann-Pick C1 protects against atherosclerosis in mice via regulation of macrophage intracellular cholesterol trafficking. *J Clin Invest*. 2008;118:2281–2290. doi: 10.1172/JCI32561
- Infante RE, Wang ML, Radhakrishnan A, Kwon HJ, Brown MS, Goldstein JL. NPC2 facilitates bidirectional transfer of cholesterol between NPC1 and lipid bilayers, a step in cholesterol egress from lysosomes. *Proc Natl Acad Sci USA*. 2008;105:15287–15292. doi: 10.1073/pnas.0807328105
- Settembre C, Zoncu R, Medina DL, Vetrini F, Erdin S, Erdin S, Huynh T, Ferron M, Karsenty G, Vellard MC, et al. A lysosome-to-nucleus signalling mechanism senses and regulates the lysosome via mTOR and TFEB. *EMBO J*. 2012;31:1095–1108. doi: 10.1038/emboj.2012.32
- Eid W, Dauner K, Courtney KC, Gagnon A, Parks RJ, Sorisky A, Zha X. mTORC1 activates SREBP-2 by suppressing cholesterol trafficking to lysosomes in mammalian cells. *Proc Natl Acad Sci USA*. 2017;114:7999–8004. doi: 10.1073/pnas.1705304114
- Boucher P, Gotthardt M, Li WP, Anderson RG, Herz J. LRP: role in vascular wall integrity and protection from atherosclerosis. *Science*. 2003;300:329–332. doi: 10.1126/science.1082095
- Terrand J, Bruban V, Zhou L, Gong W, El Asmar Z, May P, Zurhove K, Haffner P, Philippe C, Woldt E, et al. LRP1 controls intracellular cholesterol storage and fatty acid synthesis through modulation of Wnt signaling. *J Biol Chem*. 2009;284:381–388. doi: 10.1074/jbc.M806538200
- El Asmar Z, Terrand J, Jenty M, Host L, Mlih M, Zerr A, Justiniano H, Matz RL, Boudier C, Scholler E, et al. Convergent signaling pathways controlled by LRP1 (Receptor-related Protein 1) cytoplasmic and extracellular domains limit cellular cholesterol accumulation. *J Biol Chem*. 2016;291:5116–5127. doi: 10.1074/jbc.M116.714485
- Woldt E, Terrand J, Mlih M, Matz RL, Bruban V, Coudane F, Foppolo S, El Asmar Z, Chollet ME, Ninio E, et al. The nuclear hormone receptor PPAR γ counteracts vascular calcification by inhibiting Wnt5a signaling in vascular smooth muscle cells. *Nat Commun*. 2012;3:1077. doi: 10.1038/ncomms2087
- Christman MA 2nd, Goetz DJ, Dickerson E, McCall KD, Lewis CJ, Benencia F, Silver MJ, Kohn LD, Malgor R. Wnt5a is expressed in murine and human atherosclerotic lesions. *Am J Physiol Heart Circ Physiol*. 2008;294:H2864–H2870. doi: 10.1152/ajpheart.00982.2007
- Qin L, Hu R, Zhu N, Yao HL, Lei XY, Li SX, Liao DF, Zheng XL. The novel role and underlying mechanism of Wnt5a in regulating cellular

- cholesterol accumulation. *Clin Exp Pharmacol Physiol*. 2014;41:671–678. doi: 10.1111/1440-1681.12258
16. Bhatt PM, Malgor R. Wnt5a: a player in the pathogenesis of atherosclerosis and other inflammatory disorders. *Atherosclerosis*. 2014;237:155–162. doi: 10.1016/j.atherosclerosis.2014.08.027
 17. Salmon M, Gomez D, Greene E, Shankman L, Owens GK. Cooperative binding of KLF4, pELK-1, and HDAC2 to a G/C repressor element in the SM22 α promoter mediates transcriptional silencing during SMC phenotypic switching in vivo. *Circ Res*. 2012;111:685–696. doi: 10.1161/CIRCRESAHA.112.269811
 18. Bauer M, Bénard J, Gaasterland T, Willert K, Cappellen D. WNT5A encodes two isoforms with distinct functions in cancers. *PLoS One*. 2013;8:e80526. doi: 10.1371/journal.pone.0080526
 19. Harris CA, Haas JT, Streeper RS, Stone SJ, Kumari M, Yang K, Han X, Brownell N, Gross RW, Zechner R, et al. DGAT enzymes are required for triacylglycerol synthesis and lipid droplets in adipocytes. *J Lipid Res*. 2011;52:657–667. doi: 10.1194/jlr.M013003
 20. Ohno-Iwashita Y, Shimada Y, Waheed AA, Hayashi M, Inomata M, Nakamura M, Maruya M, Iwashita S, Perfringolysin O, a cholesterol-binding cytolysin, as a probe for lipid rafts. *Anaerobe*. 2004;10:125–134. doi: 10.1016/j.anaerobe.2003.09.003
 21. Suckling KE, Stange EF. Role of acyl-CoA: cholesterol acyltransferase in cellular cholesterol metabolism. *J Lipid Res*. 1985;26:647–671.
 22. Dubland JA, Allahverdian S, Besler KJ, Ortega C, Wang Y, Pryma CS, Boukais K, Chan T, Seidman MA, Francis GA. Low LAL (Lysosomal Acid Lipase) expression by smooth muscle cells relative to macrophages as a mechanism for arterial foam cell formation. *Arterioscler Thromb Vasc Biol*. 2021;41:e354–e368. doi: 10.1161/ATVBAHA.120.316063
 23. Kaiser K, Gyllborg D, Procházka J, Salašová A, Kompaniková P, Molina FL, Laguna-Goya R, Radaszkiewicz T, Harnoš J, Procházková M, et al. WNT5A is transported via lipoprotein particles in the cerebrospinal fluid to regulate hindbrain morphogenesis. *Nat Commun*. 2019;10:1498. doi: 10.1038/s41467-019-09298-4
 24. Perera RM, Zoncu R. The lysosome as a regulatory hub. *Annu Rev Cell Dev Biol*. 2016;32:223–253. doi: 10.1146/annurev-cellbio-111315-125125
 25. Sardiello M, Palmieri M, di Ronza A, Medina DL, Valenza M, Gennarino VA, Di Malta C, Donaudy F, Embrione V, Polishchuk RS, et al. A gene network regulating lysosomal biogenesis and function. *Science*. 2009;325:473–477. doi: 10.1126/science.1174447
 26. Medina DL, Di Paola S, Peluso I, Armani A, De Stefani D, Venditti R, Montefusco S, Scotto-Rosato A, Prezioso C, Forrester A, et al. Lysosomal calcium signalling regulates autophagy through calcineurin and TFEB. *Nat Cell Biol*. 2015;17:288–299. doi: 10.1038/ncb3114
 27. Settembre C, Zoncu R, Medina DL, Vetrini F, Erdin S, Erdin S, Huynh T, Ferron M, Karsenty G, Vellard MC, et al. A lysosome-to-nucleus signalling mechanism senses and regulates the lysosome via mTOR and TFEB. *EMBO J*. 2012;31:1095–1108. doi: 10.1038/emboj.2012.32
 28. Castellano BM, Thelen AM, Moldavski O, Feltes M, van der Welle RE, Mydock-McGrane L, Jiang X, van Eijkeren RJ, Davis OB, Louie SM, et al. Lysosomal cholesterol activates mTORC1 via an SLC38A9-Niemann-Pick C1 signaling complex. *Science*. 2017;355:1306–1311. doi: 10.1126/science.aag1417
 29. Fantini J, Barrantes FJ. How cholesterol interacts with membrane proteins: an exploration of cholesterol-binding sites including CRAC, CARC, and tilted domains. *Front Physiol*. 2013;4:31. doi: 10.3389/fphys.2013.00031
 30. Fantini J, Carlus D, Yahi N. The fusogenic tilted peptide (67–78) of α -synuclein is a cholesterol binding domain. *Biochim Biophys Acta*. 2011;1808:2343–2351. doi: 10.1016/j.bbame.2011.06.017
 31. Janda CY, Waghray D, Levin AM, Thomas C, Garcia KC. Structural basis of Wnt recognition by Frizzled. *Science*. 2012;337:59–64. doi: 10.1126/science.1222879
 32. Yang L, Chu Y, Wang Y, Zhao X, Xu W, Zhang P, Liu X, Dong S, He W, Gao C. siRNA-mediated silencing of Wnt5a regulates inflammatory responses in atherosclerosis through the MAPK/NF- κ B pathways. *Int J Mol Med*. 2014;34:1147–1152. doi: 10.3892/ijmm.2014.1860
 33. Zhou L, Takayama Y, Boucher P, Tallquist MD, Herz J. LRP1 regulates architecture of the vascular wall by controlling PDGFR β -dependent phosphatidylinositol 3-kinase activation. *PLoS One*. 2009;4:e6922. doi: 10.1371/journal.pone.0006922
 34. Rohlmann A, Gotthardt M, Willnow TE, Hammer RE, Herz J. Sustained somatic gene inactivation by viral transfer of Cre recombinase. *Nat Biotechnol*. 1996;14:1562–1565. doi: 10.1038/nbt1196-1562
 35. Student AK, Hsu RY, Lane MD. Induction of fatty acid synthetase synthesis in differentiating 3T3-L1 preadipocytes. *J Biol Chem*. 1980;255:4745–4750.
 36. Garratt AN, Voiculescu O, Topilko P, Charnay P, Birchmeier C. A dual role of erbB2 in myelination and in expansion of the schwann cell precursor pool. *J Cell Biol*. 2000;148:1035–1046. doi: 10.1083/jcb.148.5.1035
 37. Wilhelm LP, Wendling C, Védie B, Kobayashi T, Chenard MP, Tomasetto C, Drin G, Alpy F. STARD3 mediates endoplasmic reticulum-to-endosome cholesterol transport at membrane contact sites. *EMBO J*. 2017;36:1412–1433. doi: 10.15252/embj.201695917
 38. BLIGH EG, DYER WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol*. 1959;37:911–917. doi: 10.1139/o59-099
 39. Zhang X, Julien-David D, Miesch M, Geoffroy P, Raul F, Roussi S, Aoudé-Werner D, Marchioni E. Identification and quantitative analysis of beta-sitosterol oxides in vegetable oils by capillary gas chromatography-mass spectrometry. *Steroids*. 2005;70:896–906. doi: 10.1016/j.steroids.2005.06.004
 40. Lada AT, Davis M, Kent C, Chapman J, Tomoda H, Omura S, Rudel LL. Identification of ACAT1- and ACAT2-specific inhibitors using a novel, cell-based fluorescence assay: individual ACAT uniqueness. *J Lipid Res*. 2004;45:378–386. doi: 10.1194/jlr.D300037-JLR200
 41. Ortiz-Miranda S, Ji R, Jurczyk A, Aryee KE, Mo S, Fletcher T, Shaffer SA, Greiner DL, Bortell R, Gregg RG, et al. A novel transgenic mouse model of lysosomal storage disorder. *Am J Physiol Gastrointest Liver Physiol*. 2016;311:G903–G919. doi: 10.1152/ajpgi.00313.2015