

Update on preclinical models of cancer therapy-related cardiac dysfunction: Challenges and perspectives. A scientific statement of the Heart Failure Association (HFA) of the ESC, the ESC Council of Cardio-Oncology, and the ESC Working Group on Cellular Biology of the Heart

Alessandra Ghigo^{1*}, Pietro Ameri^{2,3}, Aarti Asnani⁴, Edoardo Bertero^{2,3}, Rudolf A. de Boer⁵, Dimitrios Farmakis⁶, Arantxa González^{7,8}, Stephane Heymans^{9,10}, Borja Ibáñez^{8,11,12}, Teresa López-Fernández^{13,14,15}, Alexander R. Lyon¹⁶, Piero Pollesello¹⁷, Amina Rakisheva^{18,19}, Konstantinos Stellos^{20,21,22,23}, Katrin Streckfuss-Bömeke^{24,25,26,27}, Carlo Gabriele Tocchetti²⁸, Thomas Thum²⁹, Peter van der Meer³⁰, Eva Van Rooij^{31,32}, Piotr Ponikowski³³, Marco Metra³⁴, Giuseppe Rosano^{35,36}, and Sophie Van Linthout^{37,38*}

¹Molecular Biotechnology Center Guido Tarone, Department of Molecular Biotechnology and Health Sciences, University of Torino, Turin, Italy; ²Cardiac, Vascular, and Thoracic Department, IRCCS Ospedale Policlinico San Martino, Genoa, Italy; ³Department of Internal Medicine, University of Genova, Genoa, Italy; ⁴Division of Cardiovascular Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA, USA; ⁵Cardiovascular Institute, Thorax Center, Department of Cardiology, Erasmus Medical Center, Rotterdam, The Netherlands; ⁶Department of Cardiology, Athens University Hospital Attikon, National and Kapodistrian University of Athens Medical School, Athens, Greece; ⁷Program of Cardiovascular Diseases, Foundation for Applied Medical Research (CIMA), Department of Cardiology, Clínica Universidad de Navarra, Instituto de Investigación Sanitaria de Navarra (IdiSNA), Universidad de Navarra, Pamplona, Spain; ⁸Centro de Investigación Biomédica en Red Cardiovascular (CIBERCV), Instituto de Salud Carlos III, Madrid, Spain; ⁹Department of Cardiology, CARIM Cardiovascular Research Institute Maastricht, Maastricht University Medical Centre, Maastricht, The Netherlands; ¹⁰Centre of Cardiovascular Research, University of Leuven, Leuven, Belgium; ¹¹Centro Nacional de Investigaciones Cardiovasculares, Madrid, Spain; ¹²Cardiology Department, IIS-Fundación Jiménez Díaz Hospital, Madrid, Spain; ¹³Cardiology Department, La Paz University Hospital, Madrid, Spain; ¹⁴IdiPAZ research Institute, Madrid, Spain; ¹⁵Cardiology Department, Quironsalud Madrid University Hospital, Madrid, Spain; ¹⁶Cardio-Oncology Centre of Excellence, Royal Brompton Hospital, London, UK; ¹⁷Content and Communication, Branded Products, Orion Pharma, Espoo, Finland; ¹⁸Department of Cardiology, City Cardiology Center, Almaty, Kazakhstan; ¹⁹Department of Cardiology, Qonaev City Hospital, Almaty, Kazakhstan; ²⁰Department of Cardiovascular Research, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany; ²¹German Centre for Cardiovascular Research (DZHK), Partner Site Heidelberg/Mannheim, Mannheim, Germany; ²²Biosciences Institute, Vascular Biology and Medicine Theme, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK; ²³Medizinische Klinik I, Department of Cardiology, Angiology, Haemostaseology and Intensive Care Medicine, University Hospital Mannheim of the Heidelberg University, Mannheim, Germany; ²⁴Clinic for Cardiology and Pneumology, Georg-August University Göttingen, Göttingen, Germany; ²⁵Medical Clinic I, Cardiology and Angiology, Justus-Liebig-University, Giessen, Germany; ²⁶Institute of Pharmacology and Toxicology, University of Würzburg, Würzburg, Germany; ²⁷Comprehensive Heart Failure Center Würzburg, Würzburg, Germany; ²⁸Department of Translational Medical Sciences (DISMET), Center for Basic and Clinical Immunology Research (CISI), Interdepartmental Center of Clinical and Translational Sciences (CIRCET), Interdepartmental Hypertension Research Center (CIRIAPA), Federico II University, Naples, Italy; ²⁹Institute of Molecular and Translational Therapeutic Strategies (IMTTS), Hannover Medical School, Hannover, Germany;

*Corresponding authors. Alessandra Ghigo, Molecular Biotechnology Center Guido Tarone, Department of Molecular Biotechnology and Health Sciences, University of Torino, Via Nizza 52, 10126 Turin, Italy. Tel: +39 011 6706335, Fax: +39 011 6706432, Email: alessandra.ghigo@unito.it
 Sophie Van Linthout, Translational Immunocardiology, Berlin Institute of Health at Charité - Universitätsmedizin Berlin, BIH Center for Regenerative Therapies (BCRT), Föhrer Strasse 15, 13353 Berlin, Germany. Tel: +49 30 450539486, Fax: +49 30 450539409, Email: sophie.van-linthout@bih-charite.de

³⁰Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; ³¹Hubrecht Institute, Royal Netherlands Academy of Arts and Sciences (KNAW) and University Medical Center Utrecht, Utrecht, The Netherlands; ³²Department of Cardiology, University Medical Center Utrecht, Utrecht, The Netherlands; ³³Center for Heart Diseases, University Hospital, Wrocław Medical University, Wrocław, Poland; ³⁴Institute of Cardiology, ASST Spedali Civili, Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, Brescia, Italy; ³⁵Department of Human Sciences and Promotion of Quality of Life, San Raffaele Open University of Rome, Rome, Italy; ³⁶Cardiology, San Raffaele Cassino Hospital, Cassino, Italy; ³⁷Berlin Institute of Health at Charité-Universitätsmedizin Berlin, BIH Center for Regenerative Therapies (BCRT), Berlin, Germany; and ³⁸German Center for Cardiovascular Research (DZHK), Partner site Berlin, Berlin, Germany

Received 29 October 2024; revised 7 February 2025; accepted 14 February 2025; online publish-ahead-of-print 11 March 2025

New anticancer therapies with potential cardiovascular side effects are continuously being introduced into clinical practice, with new and often unexpected toxicities becoming apparent only after clinical introduction. These unknown toxicities should be identified and understood beforehand to better prepare patients and physicians, enabling the implementation of effective treatments. Therefore, there is a crucial need for appropriate preclinical models to understand the biological basis of their cardiotoxicity. This scientific statement summarizes the preclinical models hitherto used, from *in vitro* two- and three-dimensional human systems to small and large animals, to pinpoint the molecular mechanisms behind the cardiotoxicity of new-generation anticancer therapies, particularly immunotherapies, and to develop potential cardioprotective strategies. Furthermore, it discusses how preclinical models have contributed to the provocative concept of heart failure being potentially tumorigenic and how the discovery of drugs with both anticancer and cardioprotective actions has revealed a common mechanistic basis for heart failure and cancer. Finally, it discusses the existing gaps between preclinical models and clinical observations in patients, how these discrepancies affect regulatory pathways and the drug development process in cardio-oncology and provides recommendations for closing these gaps.

Keywords

Cardiotoxicity • Cancer • Anticancer therapy • Immunotherapy • Preclinical models

Introduction and scope

Cancer therapy-related cardiovascular toxicity is an important clinical problem. Medications with established cardiotoxicity, but also high efficacy, like anthracycline chemotherapy and human epidermal growth factor receptor 2 (HER2)-targeted therapies, are widely used in clinical practice. In addition, many new anticancer therapies with potential cardiac side effects are continuously introduced into clinical practice. Cancer therapy-related cardiovascular toxicity can manifest as a wide range of potentially life-threatening complications, including cancer therapy-related cardiac dysfunction (CTRCD), arrhythmias, and myocarditis.¹ Preclinical models, including two- (2D) and three-dimensional (3D) cellular systems as well as small and large animals, have proven to be indispensable tools for elucidating the mechanistic pathways responsible for the cardiac side effects of traditional anticancer therapies.² This scientific statement summarizes studies from the past 5 years (2019–2024) that have utilized these models to advance the mechanistic understanding of the cardiotoxicity of new generation oncological treatments, and of the complex interaction between cancer and heart failure (HF). It also discusses the current gaps between preclinical modelling and clinical observations in patients (Figure 1), how these gaps affect regulatory pathways and the overall feasibility of the drug development process, and provides recommendations for closing these gaps. While anticancer therapies can have various adverse effects on the cardiovascular system, such as arrhythmias, hypertension, and thrombosis, and others, this review will specifically concentrate on the mechanisms leading to cardiac dysfunction and HF, aiming to provide a detailed analysis while maintaining a clear and targeted scope.

Mechanisms and clinical manifestation of cardiac dysfunction of conventional and emerging cancer therapies

Conventional therapies

Conventional cancer treatments with known cardiotoxic effects include anthracyclines, anti-HER2 therapies, tyrosine kinase inhibitors and radiotherapy. Anthracycline-induced cardiotoxicity (AIC), which irreversibly progresses from asymptomatic left ventricular dysfunction to HF,^{1,3,4} is the most extensively studied form of cardiotoxicity due to the availability of established preclinical models across various species. Studies using zebrafish, rodents, pigs and human cells have identified increased reactive oxygen species, activation of the DNA damage response, impaired autophagy, and mitochondrial dysfunction as key drivers of toxicity in cardiomyocytes⁵ and other cardiac cell types.^{6,7}

Another example is trastuzumab, a humanized monoclonal antibody targeting the ERBB2/HER2 receptor tyrosine kinase. Unlike anthracycline regimens, trastuzumab-induced cardiotoxicity is highly reversible, but can amplify and exacerbate AIC when co-administered.¹ Initial insights into the cardiotoxicity of anti-HER2 therapies came from conditional ErbB2 mouse mutants that develop severe dilated cardiomyopathy⁸ and from mice with cardiac-specific ErbB2 overexpression that are protected against doxorubicin-induced oxidative stress.⁹ Human-induced pluripotent stem cells (iPSCs) subsequently extended our mechanistic understanding of trastuzumab cardiotoxicity beyond inhibition of the pro-survival NRG1/ErbB pathway in cardiomyocytes¹⁰ and revealed alterations in metabolic pathways¹¹




Figure 1 Current preclinical models of cancer therapy-related cardiac dysfunction (CTRCD) and recommended future model systems addressing gaps between current preclinical models and clinical observations. Left panel: Overview of preclinical model systems across different species utilized in studies aimed at identifying new approaches for diagnosing, preventing, or curing CTRCD. The pros and cons of each model are highlighted. Right panel: Recommended future model systems addressing current gaps in preclinical modelling of CTRCD. 2D, two-dimensional; 3D, three-dimensional; AIC, anthracycline-induced cardiotoxicity; CM, cardiomyocyte; EHT, engineered heart tissue; iPSC, induced pluripotent stem cell; MRI, magnetic resonance imaging. (The figure has been created with Biorender.com).

and in the interaction between cardiomyocytes and vascular endothelial cells.¹²

Tyrosine kinase inhibitors (TKIs) also cause cardiac side effects.¹ Although some are specific, the majority of clinically licensed TKIs inhibit multiple kinases. Targeting BCR-ABL with ponatinib causes cardiomyopathy, HF, and vascular occlusion,¹³ with mitochondrial toxicity,¹⁴ integrated stress response activation¹⁴ and S100A8/A9-NLRP3-interleukin (IL)-1 β -mediated inflammation¹⁵ identified as key mechanisms in human iPSC and murine models.^{14,15} Sunitinib, a multi-targeted TKI in the class of vascular endothelial growth factor (VEGF) TKIs that causes hypertension and left ventricular dysfunction,¹ similarly induces mitochondrial dysfunction in engineered cardiac microtissues.¹⁶ Bruton's tyrosine kinase inhibitors (BTKi) have revolutionized the treatment of B-cell malignancies including chronic lymphocytic leukaemia and mantle cell lymphoma. Patients treated with BTKi have an increased risk of arterial hypertension, atrial fibrillation and CTRCD. Insights from studies in mice lacking an active Bruton's tyrosine kinase suggest that part of the arrhythmia risk associated with the BTKi ibrutinib is related to an off-target blockade of C-terminal Src kinase.¹⁷ However, more specific second-generation BTKi also have an increased risk of atrial fibrillation and hypertension,¹⁸ suggesting that direct Bruton's tyrosine kinase inhibition may be relevant in the pathogenesis of atrial fibrillation and hypertension in humans.

Radiation-induced heart disease, which typically manifests several years to decades after therapy, includes coronary artery disease, valvular heart disease, pericardial disease, conduction abnormalities, and HF.^{1,19} Studies in myocytes and mice exposed to radiation that mimics the dose, fractionation, and beam delivery of clinical radiotherapy protocols have identified cardiac fibrosis²⁰ and cardiomyocyte Ca²⁺ mishandling²¹ as key pathogenic events in radiation-induced heart disease.

Two decades of preclinical research have revealed the molecular underpinnings of the cardiotoxicity of most conventional oncological treatments, as extensively reviewed in Morelli *et al.*²² However, the potential interactions between cancer therapies in multimodal regimens, as well as the effects of cardiotoxic treatments against the backdrop of pre-existing cardiovascular risk factors or disease, have been insufficiently addressed in preclinical modelling and should be the focus of future studies.

Immune checkpoint inhibitors

The therapeutic landscape in cancer has been profoundly transformed by the advent of immune checkpoint inhibitors (ICIs), a class of monoclonal antibodies that boost the anticancer immune response by targeting immune-cell surface receptors.²³ ICIs target various co-stimulatory signalling molecules on T lymphocytes and antigen-presenting cells, such as cytotoxic T-lymphocyte antigen 4 (CTLA-4), programmed cell death protein 1 (PD-1), and its ligand (PD-L1), thereby breaking the immune tolerance of T cells against cancer cells.²⁴

Despite their proven efficacy,^{25,26} the use of ICIs has led to the occurrence of several immune-related adverse events (irAEs) caused by the hyperactivation of the immune system against non-cancerous tissues.²⁷ Although irAEs depend on the type of cancer, the ICI used, the duration of exposure and the dose administered, ICI immunotherapy is associated with an increased risk of cardiac events, including myocarditis, arrhythmias, HF, and cardiovascular death.^{28–31}

Animal models have been instrumental in deciphering the exact pathophysiological mechanisms underlying the cardiovascular toxicities of ICIs (extensively reviewed in Tocchetti *et al.*³¹ and Gergely *et al.*³²). Studies in mice with genetic deletion of immune checkpoints like *Pdcd1* and *Ctla4*, which are the murine equivalents

of the genes encoding human PD-1 and CTLA-4, have shown that ICI cardiovascular toxicity is associated with increased inflammation. This is secondary to infiltration of macrophages³³ and cytotoxic CD8⁺ T cells specific for α -myosin³⁴ into the heart. In mice with monoallelic loss of *Ctla4* combined with a complete genetic absence of *Pdcd1*, myocardial infiltration was associated with severe electrocardiographic abnormalities and left ventricular dysfunction observed in some ICI-treated patients.³⁵ Interestingly, in this murine model, which closely recapitulates the clinical and pathological features of ICI-associated myocarditis, intervention with CTLA4-Ig (abatacept) ameliorated disease progression.³⁵ This was supported by a case series of patients receiving abatacept, which successfully attenuated the fulminant course of ICI myocarditis, although the optimal dosing for this indication and pharmacokinetics in humans are complex and have not been definitively established.³⁵

Preclinical models will be essential to validate new strategies, like fasting mimicking diet cycles,³⁶ to protect the heart from the adverse effects of immunotherapy. At the same time, these models should be used to address current knowledge gaps. The observation that only 1–1.5% of all patients receiving ICI develop severe cardiovascular toxicity,^{27,31} along with the variable progression of ICI-mediated myocarditis among individuals, suggests that additional genetic and environmental factors must contribute to the manifestation of this condition. Indeed, traditional cardiovascular risk factors such as diabetes, hypertension, and dyslipidaemia do not predict which patients are at risk. In contrast, patients undergoing combination ICI therapy, those with pre-existing autoimmune diseases, and those with specific cancers, such as thymic cancers, are at increased risk.^{31,37} This indicates that the immune phenotype plays a critical role, presenting a challenge for translational models due to significant species-related differences in immunobiology. Therefore, preclinical models which recapitulate the human immune response are critical to decipher the patient-specific response towards ICI.

CAR-T and TCR-T cells

Chimeric antigen receptor T cell (CAR-T) therapy is a form of immunotherapy in which patient's own T cells are genetically engineered to recognize and eliminate cancer cells. CAR-T cells are generated by inducing the expression of an artificial protein, the chimeric antigen receptor, consisting of a T cell signalling domain and an antigen recognition domain that selectively targets a cancer-specific antigen. CAR-T therapy has drastically improved the prognosis of previously incurable haematologic malignancies, particularly those derived from B cells.³⁸ The potential of CAR-T cells to treat a range of conditions beyond cancer, including autoimmune diseases, chronic infections, cardiac fibrosis, senescence-associated diseases and other conditions, is also being explored in preclinical models.³⁹

The therapeutic efficacy of CAR-T therapy is compromised by serious clinical complications, the most common being cytokine release syndrome (CRS), a systemic inflammatory syndrome characterized by immune cell activation and high levels of circulating cytokines.⁴⁰ CRS develops on average 5–7 days after CAR-T

infusion and its manifestation varies in severity from fever, with or without constitutional symptoms, to a potentially life-threatening condition.⁴¹ Evidence from preclinical mouse models⁴² has demonstrated the importance of IL-1 β ⁴³ and IL-6⁴⁴ in the pathophysiology of CRS, and the benefit of IL-1 receptor and IL-6 blockade in ameliorating CRS. To date, several clinical trials are underway testing the efficacy of CAR-Ts combined with IL-1 receptor antagonism,⁴⁵ while IL-6 receptor blockade with tocilizumab has shown benefit in patients with moderate or severe CRS.⁴⁶

Cardiovascular toxicity is a frequently reported complication of CAR-T therapy. CAR-T-related cardiovascular toxicity often occurs in the context of CRS,^{47,48} and the risk of cardiovascular events correlates with CRS severity.⁴⁹ The most common manifestations are new-onset HF and supraventricular tachyarrhythmias, but CAR-T-related severe CRS and cardiac dysfunction can also lead to refractory cardiogenic shock and death (Table 1).^{47,49–56} However, it is important to emphasize that these clinical data are derived from retrospective analyses, which are inherently susceptible to bias and may not fully reflect real-world clinical scenarios. The pathophysiology of CAR-T-related cardiovascular toxicity has not been investigated so far in animal models of CRS,⁵⁷ but myocardial depressing effects of cytokines like IL-6 have been described in mice.⁵⁸

Cardiac safety concerns have also arisen with T cell receptor (TCR) T cell (TCR-T) therapies, immunotherapies which use engineered TCR-Ts expressing cancer antigen-specific TCRs to redirect T cells against tumour cells.⁵⁹ Unlike CAR-T, which target cell surface antigens, TCR-T recognize epitopes presented by the major histocompatibility complex (MHC). Although TCR-T is less likely to induce CRS compared with CAR-T therapy,⁵⁹ two cases of fatal myocarditis due to TCR-T targeting MHC class I-restricted epitopes derived from the melanoma-associated antigen-3 (MAGE-A3) have been reported.^{60,61} Mechanistically, these effects were linked to cross-reactivity against titin, as demonstrated in iPSC-derived cardiomyocytes (iPSC-CMs) exposed to MAGE-A3-transduced T cells.^{60,61}

In conclusion, CAR-T and TCR-T therapies are associated with cardiac events stemming from cardiotoxic effects of interleukins in the context of CRS, and off-target cytotoxic effects of engineered T cells, respectively. A deeper understanding of these pathomechanisms requires further investigation in appropriate preclinical models.⁵⁷

Preclinical models to study cancer therapy-related cardiac dysfunction

In vitro models

Induced pluripotent stem cell-derived cardiomyocyte and non-cardiomyocyte cells in 2D and 3D tissue models

For a long time, there was a lack of human cardiomyocyte culture models that could be used to understand the molecular and cellular physiology of CTRCD; as a result, preclinical cardio-oncology research relied primarily on rodent cardiomyocytes as an *in vitro*

Table 1 Frequency of cytokine release syndrome and cardiovascular events in adult patients receiving CAR-T therapy

Study	No. of patients and type of malignancy	CRS	LVSD/AHF	Arrhythmias	CV death
Alvi et al. ⁴⁷	n = 137 DLBCL 83 (60.6), follicular lymphoma 36 (26.3), MM 11 (8.0), other 7 (5.1)	81 (59%)	6 (4.4) AHF, of which 4 new-onset	5 (3.6), of which 3 AF and 2 SVT	6 (4.4)
Lefebvre et al. ⁴⁹	n = 145 ALL 36 (24.8), CLL 66 (45.5), DLBCL 43 (29.7)	176 events in 104 pts (71.7)	22 AHF events in 21 pts (14.5)	12 episodes of AF in 11 pts (7.6), 1 SVT, 1 NSVT	2 (13.8)
Ganatra et al. ⁵⁰	n = 187 DLBCL 137 (73.3), follicular lymphoma 35 (18.7), other B-cell malignancies 15 (8.0)	155 (82.9)	12 (10.3% of 116 pts with echo) LVSD, of which 6 AHF events	10 (5.3), unspecified	3 (1.6)
Goldman et al. ⁵¹	n = 2657 NHL 1777 (83.7), ALL 306 (14.4), other 39 (1.8)	1457 (54.8)	Cardiomyopathy 69 (2.6)	74 (2.8), of which 55 AF, 10 VT, 4 VF	NA
Qi et al. ⁵²	n = 126 MM 78, NHL 25, ALL 23	103 (81.7)	15 (11.9) AHF, 1 (0.8) LVSD	7 (5.6), unspecified	2 (1.6)
Lee et al. ⁵³	n = 78 RRMM 78 (100)	16 (20.5) grade 2 or above	4 (5.1) LVSD, of which 1 (1.3) AHF	11 (14.1), of which 8 AF/flutter and 3 NSVT	1 (1.3)
Lefebvre et al. ⁵⁴	n = 44 Lymphoma 43 (97.7), ALL 1 (2.3)	24 events in 23 (52.3) pts	1 (2.3) AHF with preserved LVEF	1 (2.3) AF	None
Mahmood et al. ⁵⁵	n = 202 B-cell lymphoma 143 (70.8), ALL 59 (29.2)	139 (68.8)	26 (12.8) AHF, of which 2 (1.0) cardiogenic shock	37 (18.3) atrial arrhythmias	1 (0.5)
Korell et al. ⁵⁶	n = 137; B-cell lymphoma 104 (75.9), B-ALL 19 (13.8), MM 14 (10.2)	75 (60.1)	4 (2.9) LVSD	5 (3.6) AF	None

AF, atrial fibrillation; AHF, acute heart failure; ALL, acute lymphoblastic leukaemia; CRS, cytokine release syndrome; CV, cardiovascular; DLBCL, diffuse large B-cell lymphoma; LVEF, left ventricular ejection fraction; LVSD, left ventricular systolic dysfunction; MM, multiple myeloma; NA, not available; NHL, non-Hodgkin's lymphoma; NSVT, non-sustained ventricular tachycardia; pts, patients; RRMM, relapsed and refractory multiple myeloma; SVT, supraventricular tachycardia; VF, ventricular fibrillation; VT, ventricular tachycardia.

model. More recently, the development of 2D and 3D culture systems has enabled the use of iPSC-CMs as a screening platform for cardiotoxic effects of cancer therapies, to predict cardiotoxicity risk,^{62,63} or to analyse the underlying pathophysiology.^{10,11} By generating human iPSC-CM and non-cardiomyocyte cell types, it is possible to establish patient-specific disease models to define cardiotoxicities at the individual and population levels. In addition, patient-specific iPSC models can be used to identify and validate single nucleotide polymorphisms associated with CTRCD, ultimately guiding the design of clinical trials.⁶⁴

Healthy control iPSC-CMs have initially been used to demonstrate general cardiotoxic effects of anthracyclines, such as apoptosis, reactive oxygen species production, mitochondrial dysfunction, and electrophysiological and Ca²⁺ changes,^{65,66} and led to the identification of telomerase overexpression as a therapeutic option in CTRCD.⁶⁷ On the other hand, patient-specific iPSC models have been shown to recapitulate the predisposition of

individual breast cancer patients to low doxorubicin-induced cardiotoxicity⁶² or trastuzumab-induced cardiac dysfunction.¹¹ Another patient-specific stem cell model of CTRCD demonstrated that patients with aggressive B-cell lymphoma are persistently more susceptible to detrimental effects of high doxorubicin treatment with hyperactivation of Ca²⁺/calmodulin-dependent protein kinase II δ (CaMKII δ), resulting in arrhythmogenic triggers and cellular dysfunction.⁶⁸ Furthermore, iPSC-CMs generated from paediatric oncology patients with acute anthracycline cardiotoxicity identified cardioprotective micro⁶⁹ and circular⁷⁰ RNAs. Finally, a stem cell model of paediatric CTRCD patients allowed the identification of genomic variants of drug transporters associated with doxorubicin-induced cardiotoxicity, thereby confirming the evidence for a genetic predisposition to CTRCD,⁷¹ including several somatic and systemic mutations that trigger the development of CTRCD.^{68,72}

Although iPSC-CMs hold great promise as an *in vitro* research tool for understanding the cardiotoxicity of oncological drugs,

they also differ significantly from adult cardiomyocytes in that they are structurally and functionally immature, resembling foetal cardiomyocytes.⁷³ This immaturity can be partially corrected by embedding the iPSC-CMs in a 3D environment, such as the so-called dynamic engineered heart tissue model⁷⁴ or engineered heart myocardium.⁷⁵ In addition to promoting maturation, as shown by a faster conduction velocity, better alignment, and increased contractile power, this also provided a method to analyse iPSC-CM behaviour in a more physiological environment and the possibility to include other cell types, including endothelial cells and cardiac fibroblasts. Exposing dynamic engineered heart tissues or engineered heart myocardium to anthracyclines resulted in tissue dilatation and reduced contractile force, both key features of the clinical phenotype of AIC.^{68,76} Furthermore, these 3D dynamic-engineered cardiac tissues could recapitulate the induction of senescence observed in the hearts of patients with severe doxorubicin-induced cardiotoxicity. However, pharmacological inhibition of senescence did not show any functional improvement in these models, calling into question the use of senolytic drugs as a potential cardioprotective therapy.⁷⁶ Other maturation strategies for iPSC-CMs currently being explored include long-term culture, mechanical and/or electrical stimulation, 3D cardiac organoids of different cell types, the addition of anti-inflammatory macrophages⁷⁷ and combinatorial approaches.^{78,79} While the majority of the studies with CTRCD stem cell models have focused specifically on autonomic iPSC-CM processes, the interaction with other relevant (cardiovascular) cell types, including immune cells, and their integration into 3D engineered tissue systems, should also be considered. The inclusion of multicellular models into preclinical studies of CTRCD is expected to uncover previously unexplored aspects of the disease. Beyond immaturity and limited ability to reproduce multicellular communications, iPSC-CMs face additional challenges, including subtype specification, scarce interlaboratory reproducibility, labour-intensive protocols, and high costs.⁸⁰ Moreover, the reprogramming, culture, and differentiation of human iPSCs demand significant time and resources, limiting the feasibility of large-scale research for non-established iPSC laboratories. While substantial progress has been made,^{81,82} further efforts are needed to optimize cardiomyocyte differentiation protocols, prioritizing simplicity, reproducibility, and cost-effectiveness, to enable broader application.

Cardiac slices

Although iPSC models have revolutionized our understanding of CTRCD, they still do not fully emulate the complex cellular composition of human heart tissue. Myocardial slices – thin, living sections of heart tissue – offer a unique advantage for studying cardiac function and drug effects in a context that closely resembles the native tissue environment. Using a biomimetic culture system, pig and human heart slices can remain completely viable and functionally and structurally intact for 6 days in culture, also enabling the assessment of contraction and relaxation kinetics on isolated single myofibrils after culture.^{83,84} To date, canine, porcine, and human myocardial slices have been used to study the cardiotoxicity of doxorubicin, trastuzumab, and sunitinib. Heart slices effectively

recapitulate the expected toxicity of doxorubicin^{85,86} and trastuzumab,⁸⁶ similar to human iPSC-CMs, but are superior in detecting sunitinib cardiovascular toxicity at clinically relevant concentrations.⁸⁶ Therefore, these systems should be considered for validating translational findings from other preclinical models in future studies.

In vivo models

Zebrafish

The zebrafish is an *in vivo* whole organism model that has been used for decades to study the mechanisms of cardiovascular disease (CVD), as well as cancer growth and metastasis. In cardio-oncology, zebrafish have been used to identify new small molecule therapies that protect against doxorubicin-induced cardiomyopathy without compromising anti-tumour activity.⁸⁷ Zebrafish models have also been developed to study cardiovascular toxicities associated with cyclophosphamide⁸⁸ and VEGF inhibitors.^{89,90} Zebrafish have the following strengths when compared to *in vitro*, rodent, or large animal models: (1) small size and optical transparency in the embryonic stage, facilitating high throughput screening with direct cardiac imaging; (2) high degree of genetic conservation between zebrafish and humans, particularly for metabolic and cell death-related pathways; (3) advances in genome editing technology, allowing rapid and efficient generation of CRISPR/Cas9 mutants; (4) studies at scale with large numbers and significantly lower cost compared to rodent models. One disadvantage is the early developmental stage required for most high-throughput applications. As a result, validation of new molecular mechanisms discovered in zebrafish embryos typically requires the use of complementary adult model systems.

Rodents

Rodents are the most widely used preclinical model in cardio-oncology research. CTRCD can be easily recapitulated by treating rats and mice with the anticancer drug according to protocols that closely mimic human therapeutic regimens in terms of route of administration, dosage and pharmacokinetic profile.² Over the past decades, these models have been instrumental in identifying the mechanisms behind the cardiotoxicity of conventional chemotherapies, primarily anthracyclines, and in discovering potential cardioprotective strategies.⁹¹ However, direct treatment of rodents with anticancer therapies does not always effectively reproduce the associated cardiotoxicity, as seen with trastuzumab and other anti-HER2 antibodies, because these drugs are unable to recognize the murine counterpart of the HER2 receptor.⁹² Homozygous human HER2 transgenic mice do not represent a valuable alternative as they are embryonic lethal.⁹³ Consequently, most of the knowledge on the cardiotoxicity of anti-HER2 therapies has been obtained from mice with cardiomyocyte-specific ErbB2 deletion or overexpression.^{8,9} These studies, along with later investigations involving the genetic deletion of immune checkpoints, exemplify the strengths of mouse genetics in advancing our understanding of the mechanistic basis of cardiotoxicity. Furthermore, this highlights the added value of murine models,

particularly compared to large animals, where genetic manipulation is more challenging.

Another significant advantage of rodent models over human *in vitro* systems is the ability to assess the impact of novel cardioprotective candidates on the anti-tumour activity of cancer therapies. For example, xenotransplantation of cells derived from cutaneous melanoma, a prototypical cancer type treated with immunotherapy, originally led to the identification of tumour necrosis factor (TNF)- α blockade, but not CD8⁺ T cell depletion, as a means of preventing ICI-mediated myocarditis without compromising the anticancer efficacy of the anti-PD1 therapy.⁹⁴ Nevertheless, despite this positive preclinical evidence, caution is advised against the use of TNF- α antibodies for ICI myocarditis and HF,¹ as clinical studies in HF patients have shown neutral or even negative outcomes.⁹⁵ In addition to xenotransplantation of solid tumours into syngeneic mice, the use of patient-derived xenografts (PDX) can further strengthen the translational value of the study. PDX generated using tissue from a woman with triple-negative breast cancer, and implanted in severe combined immunodeficiency mice, demonstrated that a small molecule allosteric inhibitor of the pro-apoptotic molecule BAX can provide cardioprotection, without compromising the efficacy of doxorubicin in reducing the burden of breast cancer *in vivo*.⁹⁶ However, a major disadvantage of xenografts in immunodeficient mice is the inability to model interactions of the immune system with the tumour and the cardiovascular system. In this respect, genetic models with spontaneous tumour growth represent a valuable alternative. For example, using *ErbB2/Neu* transgenic mice, it has been shown that inhibition of PI3K γ effectively protects against AIC, while inducing anti-tumour immune responses, leading to reduced mammary cancer growth.⁹⁷

In summary, although novel human *in vitro* models of CTRCD are emerging, rodents remain invaluable for studying complex cell–cell and organ–organ interactions, including tumour–heart interactions as well as the involvement of the immune system.

Large animals

Despite their scientific value, murine models differ significantly from human anatomy and physiology, limiting the straightforward translation to the clinic. Because the hearts of large mammals have increased similarity to humans, diagnostic and therapeutic strategies that prove beneficial in large animal studies are more likely to succeed in clinical trials than those tested only in small animals.⁹⁸ Several large animal models of CTRCD (e.g. pigs, primates, sheep, cows, dogs) have been reported,^{99–105} most of which test for cardiac side effects of anthracyclines. Large animal models offer the opportunity to test diagnostic protocols that can be replicated in humans using exactly the same equipment. One example is the discovery of T2 relaxation time prolongation on magnetic resonance imaging as a very early marker of anthracycline cardiotoxicity in pigs and monkeys,^{102,104} which was later validated in clinical studies.^{106,107}

Large animal models are also useful when testing potential cardioprotective therapies prior to clinical trials. Through the use of these models, there are examples of some therapies that have been withdrawn from the translational portfolio because they did not

show any benefit in a large animal model.¹⁰⁰ A successful example of this strategy is the testing of remote ischaemic conditioning as an intervention to prevent AIC. This non-invasive intervention was tested in a porcine model of AIC and showed that its use prior to each doxorubicin injection resulted in a significant improvement in left ventricular ejection fraction through a mechanism involving mitochondrial protection.¹⁰⁸ These results in the pig model have served as basis for the execution of the ongoing RESILIENCE trial (NCT05223413), which is investigating the benefits of weekly remote ischaemic conditioning in a population of cancer patients at high risk of AIC.

Despite their undeniable advantages, large animal models present significant challenges, including high costs, specialized infrastructure and handling requirements, and longer lifespans and gestation periods. Additionally, genetic manipulation in these animals is complex and may raise ethical concerns.⁹⁸ Consequently, small animals, like murine models, remain the preferred choice in preclinical research, including in the field of cardio-oncology.

Mechanisms and preclinical models of reverse cardio-oncology

Recent studies suggest that the relationship between CVD and cancer extends well beyond CTRCD. Community-based cohort studies have shown that individuals with HF are at increased risk of cancer incidence and mortality compared with the general population, a phenomenon referred to as ‘reverse cardio-oncology’.^{109–111} Murine models, particularly tumour xenografts, have been instrumental in demonstrating that the relationship between cancer and HF is not merely a clinical association, but rather a causal one. Landmark studies first demonstrated that HF, induced by permanent ligation of the left anterior descending coronary artery or by pressure overload-induced left ventricular remodelling via transverse aortic constriction, leads to increased tumour burden in a mouse model of spontaneous intestinal adenomatosis,^{112,113} independent of haemodynamic changes.¹¹³ Subsequent studies have confirmed these findings, showing that cardiac remodelling from left anterior descending coronary artery ligation, transverse aortic constriction, ATF3 overexpression, or prolonged low-dose phenylephrine infusion promotes proliferation, and in some cases metastatic dissemination, of MMTV-PyMT or 4T1 mammary cancer and Lewis lung carcinoma cells (Table 2).^{114–121}

Preclinical models have further provided evidence for the role of secreted factors¹¹³ and the immune system¹¹⁸ as mediators between HF and cancer. Remodelling factors that have been suggested as mediators of the tumour-supporting effect of HF include serpinA3, periostin, nerve growth factor, connective tissue growth factor, fibronectin, and inflammatory cytokines.^{112–115,117,120,121} Myocardial infarction (MI)-induced epigenetic reprogramming of Ly6C^{hi} monocytes in the bone marrow reservoir to an immunosuppressive phenotype and their recruitment to tumours has also been shown to underlie MI-induced tumour growth.¹¹⁸ However, HF induced by transverse aortic constriction surgery also promoted tumorigenesis in immunodeficient mice, indicating that immune

Table 2 Preclinical models in reverse cardio-oncology studies

Cardiac intervention	Cancer model	Strain, sex	Effects of intervention	Main identified pathway	Effects on tumour cells (<i>In vitro</i>)	Ref.
TAC (vs. sham surgery)	ApcMin/J mouse (intestinal adenoma)	Male	On the heart LVH, fibrosis	Not provided (suggestion of fibrotic factors)	Not studied	112
TAC (vs. sham surgery)	MMTV-PyMT (mammary cancer) i.v. MMTV-PyMT cells (cancer metastases)	C57Bl/6, female	On the heart LVH, LVSD, no fibrosis	–	↑ proliferation of MMTV-PyMT and LCC cells upon incubation with serum from TAC-operated mice	114
TAC (vs. sham surgery)	LCC cells into the s.c. tissue of the flank (lung cancer) i.v. LCC cells (cancer metastases)	C57Bl/6, male	On the heart LVH, LVSD, no fibrosis	Cardiac remodelling → tumorigenic factors (CTGF, POSTN)	↑ proliferation of MMTV-PyMT and LCC cells upon incubation with POSTN	
TAC (vs. sham surgery)	MMTV-PyMT cells into the mammary fat (mammary cancer)	NOD/SCID, female	On the heart LVH, LVSD, no fibrosis	–	–	
TAC (vs. sham surgery)	MMTV-PyMT cells into the mammary fat (mammary cancer)	JDP2xATF3 female	On the heart No cardiac remodelling	= CTGF and periostin levels	–	
TAC (vs. sham surgery)	LCC cells into the sc tissue of the flank (lung cancer)	Postn ^{-/-} C57Bl/6, male	On the heart LVH, LVSD	Inflammatory and tumorigenic cytokines, FN = serum POSTN	↑ proliferation of MMTV-PyMT and LCC cells upon incubation with serum from Postn ^{-/-} mice	115
Permanent LAD ligation (vs. sham surgery)	Renca cells into the kidney (renal cancer)	BALB/c, NS	On the heart LVH, LVSD	–	Low expression of ITGB1 in Renca cells	
TAC (vs. sham surgery)	MMTV-PyMT cells into the mammary fat (mammary cancer)	C57Bl/6, female	On the heart LVH, LVSD	–	= proliferation of ITGB1 ^{-/-} and Renca cells upon incubation with serum from TAC-operated mice	
Permanent LAD ligation (vs. sham surgery)	ITGB1 ^{-/-} vs WT CRC tumour model (germ-free mice, AZM/DDDS)	C57Bl/6, male	On the heart LAD: LVSD, fibrosis TAC: LVH, fibrosis	Foecal transplantation: ↑ tumour growth in AZM/DDDS model	↑ proliferation in AZM/DDDS colonic tumors	116
Permanent LAD ligation (vs. sham surgery)	4 T1 cells into the mammary fat pad (mammary cancer)	BALB/c, female	On the heart LVSD, fibrosis, congestion	Cardiac remodelling → NGF	↑ proliferation of 4 T1 cells upon incubation with NGF, abrogated by silencing or GW441756	117
Permanent LAD ligation (vs. sham surgery)	4 T1 cells into the mammary fat pad (mammary cancer) + GW441756	BALB/c, female	On the heart LVSD, fibrosis, congestion	Reduced tumour growth and proliferation		

Table 2 (Continued)

Cardiac intervention	Cancer model	Strain, sex	Effects of intervention		Main identified pathway	Effects on tumour cells (<i>in vitro</i>)	Ref.
			On the heart	On the tumour			
Permanent LAD ligation (vs. sham surgery)	E0771 cells into the mammary fat pad (mammary cancer) MMTV-PyMT (mammary cancer)	C57BL/6j, female	No cardiac remodelling	↑ tumour growth and proliferation	↑ Ly6C ^{hi} monocytes, which halt anti-tumour immune responses	–	118
Permanent LAD ligation (vs. sham surgery) Transplantation of infarcted heart (vs. transplantation of sham-operated heart)	ApcMin/J mouse (intestinal adenoma)	C57BL/6j, male	LVSD, fibrosis, ↓ SBP, congestion	↑ tumour growth and proliferation ↑ dissemination to the lung	Cardiac remodelling → tumorigenic factors (Serpina3)	↑ proliferation of HT29 colorectal cancer cells upon incubation with SerpinA3	113
Permanent LAD ligation (vs. sham surgery)	Renca cells into the kidney (renal cancer)	BALB/c, male	LVSD, fibrosis, no congestion	↑ tumour growth and proliferation	–	= proliferation of Renca cells upon incubation with of SerpinA3 and other remodelling factors	119
αMHC-τTA × ATF3	MMTV-PyMT cells into the mammary fat (mammary cancer) LCC cells into the s.c. tissue of the flank (lung cancer) i.v. MMTV-PyMT cells (cancer metastases)	C57Bl6, female C57Bl6, male C57Bl6, female	LVH, LVSD, fibrosis LVH, LVSD –	↑ tumour growth, and proliferation ↑ tumour fibrosis ↑ tumour growth ↑ lung metastases	Cardiac remodelling → tumorigenic factors (POSTN, CTGF, FN, SerpinE1, SerpinA3)	↑ proliferation of MMTV-PyMT and LLC cells upon incubation with serum from ATF3-transgenics	120
PE (10 mg/kg/day) for 4 weeks	MMTV-PyMT cells into the mammary fat (mammary cancer)	C57Bl/6, female	LVH and fibrosis, no dysfunction	↑ tumour growth and proliferation	Cardiac remodelling → tumorigenic factors (POSTN, CTGF, FN)	↑ proliferation of MMTV-PyMT cells upon incubation with serum from PE-infused mice	121

αMHC-τTA, ^α myosin heavy chain-tetracycline transactivator; ApcMin/J, adenomatous polyposis coli mutant; ATF3, activating transcription factor 3; AZM/DDDS, azithromycin/dexamethasone; CTGF, connective tissue growth factor; CRC, colorectal cancer; FN, fibronectin; ITGB1^{-/-}, integrin beta knockout; i.v., intravenous; JDP2, Jun dimerization protein 2; LAD, left anterior descending coronary artery; LCC, large cell carcinoma; LLC, Lewis lung carcinoma; LVH, left ventricular hypertrophy; LVSD, left ventricular systolic dysfunction; Ly6C^{hi}, lymphocyte antigen 6 complex, locus c high; MMTV-PyMT, mouse mammary tumour virus-polyoma middle T antigen; NGF, nerve growth factor; NOD/SCID, non-obese diabetic/severe combined immunodeficiency; NS, not specified; PE, phenylephrine; POSTN, periostin; Postn^{-/-}, periostin knockout; SBP, systolic blood pressure; s.c., subcutaneous; SerpinA3, serine protease inhibitor A3; SerpinE1, serine protease inhibitor E1; TAC, transverse aortic constriction; WT, wild-type.

dysregulation may not be essential.¹¹⁴ In addition, microbial dysbiosis, which often is observed in HF, is present in a mouse model of MI-induced HF and transplantation of the 'HF-related microbiome' into recipient mice accelerated colon tumour growth.¹¹⁶

In summary, preclinical models suggest that HF may promote cancer growth through secreted factors and inflammatory mediators, indicating the potential of available preclinical models to evaluate the anticancer effect of HF medication.¹²²

Novel insights from drugs with both cardioprotective and anti-tumour activity

In addition to the intriguing concept that HF may be tumorigenic, the fact that patients with CVD also have a higher risk of cancer than that in the general population suggests potential common mechanisms and therapeutic targets.^{123–125} This notion has been supported by some recently developed anti-tumour therapies, particularly those that target the NLRP3 inflammasome, such as NLRP3 and IL-1 inhibitors, which have also been shown to protect the cardiovascular system.

Consistent with preclinical results obtained on the role of IL-1 in tumour progression, IL-1 receptor antagonists and anti-IL-1 monoclonal antibodies can inhibit primary tumour growth and metastasis in animal studies.¹²⁶ Furthermore, genetic associations in cancer patients indicate that combining anti-IL-1 strategies with checkpoint blockade immunotherapy may benefit patients with haematological, pancreatic or breast cancer, and clinical trials are ongoing.¹²⁶ As IL-1 can stimulate CRS-related toxicity of CAR-T therapy, this combination might not only be effective against cancer but may also help to reduce the cardiovascular toxicity associated with cell therapies.⁵⁷ The Canakinumab Anti-Inflammatory Thrombosis Outcome Study (CANTOS), a large, randomized, double-blind, placebo-controlled phase 3 trial of canakinumab in patients with a history of MI with atherosclerosis,¹²⁷ showed that canakinumab was associated with a dramatic (>60%) reduction in lung cancer incidence and mortality,¹²⁸ specifically in molecular subtypes of lung cancer.¹²⁹ Based on these results, single-agent canakinumab has been prospectively investigated in a randomized, double-blind, placebo-controlled phase 3 trial in patients with surgically resected stage II–III non-small cell lung cancer (NCT03447769). However, this trial (CANOPY-A) did not show a benefit in disease-free survival,¹³⁰ and further results from other trials will follow.

NLRP3 inhibition is another therapy with potential combined anti-tumour and cardioprotective properties. The NLRP3 inflammasome is a source of IL-1 β and IL-18, and regulates pyroptosis, autophagy, and oxidative stress,¹³¹ as evidenced by the known anti-inflammatory and cardiovascular protective properties of colchicine,^{132,133} the oldest non-specific inflammasome inhibitor. Colchicine is an effective treatment for pericarditis and may also reduce the risk of ischaemic cardiovascular events in patients with a recent MI, although at the expense of increased non-CVD mortality.¹³⁴ Beside preclinical evidence of its potential cardioprotective properties, and the clinical evidence of IL-1 and IL-6 inhibition against cardiovascular risk,¹³⁵ direct clinical evidence of NLRP3

inhibition in protecting against HF or CVD is still lacking. Various clinical trials studying colchicine or IL-1 inhibition in stable or acutely decompensated HF with reduced ejection fraction (reviewed in Olsen *et al.*¹³⁶) have also failed to convincingly demonstrate a strong protective potential. On the other hand, targeting the NLRP3 inflammasome to further improve immunotherapy may be a promising therapeutic approach for prostate, breast, colorectal and lung cancer, as well as melanoma.¹³⁷ However, NLRP3 may act as a double-edged sword in carcinogenesis, exerting both positive and negative effects.¹³⁸ More preclinical and clinical research is therefore needed to identify the best subgroup of cancer patients who will benefit from NLRP3 inhibition with common cardiovascular and cancer risk. This also applies to the danger associated with molecular pattern and activator of the NLRP3 inflammasome, S100A9,^{139,140} which has an ambivalent role in cancer¹⁴¹ and CVD.¹⁴²

In conclusion, specific inflammatory molecular pathways, particularly those related to the NLRP3 inflammasome, may serve as common therapeutic targets for both cancer and HF. However, further mechanistic validation in preclinical models is essential, especially given the unclear role of the NLRP3 inflammasome (and its drivers) in both tumorigenesis and CVD.

Limitations of current models

Gaps between preclinical modelling and clinical observations in human patients

The interpretation of data from preclinical modelling is crucial for avoiding translational failures.¹⁴³ The aim is not only to avoid loss of time and the misallocation of funds and resources, but more importantly to avoid patients entering clinical phases that are doomed to fail or, conversely, to reduce the risk of promising new therapies being overlooked. Information obtained in the preclinical phase of a new chemical entity, or a new method, provides valuable insights during the clinical development phases. Ignoring or misinterpreting such signs has proven to be costly and even dangerous when candidate therapies enter clinical evaluation, leading to time and resources lost, and patients enrolled in fruitless trials. This underscores the importance of effective translational preclinical models and the need for precise preclinical model requirements to guide translational scientists.

Despite efforts to replicate the key features of human CTRCD, current preclinical models have significant limitations that need to be addressed in future research (Table 3, Figure 1). To accurately model CTRCD, it is essential to recognize that many anticancer treatments cause late-onset cardiotoxicity, requiring the development of *in vivo* and *in vitro* models that reflect the progressive dysfunction observed in patients.^{76,144,145} This can be accomplished by mimicking clinical regimens with repeated exposures to low to medium doses of anticancer drugs or radiotherapy. Yet, a significant number of preclinical studies still utilize *in vivo* or *in vitro* systems exposed to a single high dose of anticancer treatment, which raises concerns about the translational relevance of the

study conclusions. Another important aspect to consider in pre-clinical modelling of CTRCD is species selection. For instance, the use of inappropriate animal species, such as rodents, has led to confusion in studies investigating the cardiotoxicity of the humanized monoclonal antibody trastuzumab, which does not bind to the equivalent of the HER2 receptor (Neu) in rats or mice.⁹² Therefore, future research on the cardiac side effects of the emerging class of drugs known as antibody-drug conjugates¹⁵⁶ should involve careful species selection or the use of humanized models,¹⁵⁷ particularly for those based on trastuzumab.

Efforts should also focus on modelling inter-organ interactions, which are increasingly recognized as crucial in the pathogenesis of CTRCD. In this respect, despite increasing political pressure to reduce animal research in the European Union, animal models remain indispensable in basic cardio-oncology studies. They play a crucial role in the translational pathway, providing the only means of gaining essential pathophysiological insights into disease mechanisms at both (inter)-organ and systemic level.⁹⁸ Specifically, interactions between the myocardium and the primary target of anticancer therapies, the tumour, are essential to understand. While most basic cardio-oncology research is conducted using tumour-free models, growing evidence suggests that CTRCD can arise not only from the direct toxic effects of anticancer drugs on cardiac cells, but also from effects mediated through the tumour. Consequently, the effects of cancer therapy can vary depending on whether it is administered to tumour-free or tumour-bearing animals. For example, ICI therapy has been shown to induce left ventricular dysfunction in mice transplanted with syngeneic PD1-sensitive melanomas but not in tumour-free animals.⁹⁴ Therefore, employing tumour-bearing models is advised to more accurately replicate all factors involved in CTRCD pathogenesis (Table 3). In large animals, this approach is rarely feasible due to the lack of syngeneic tumour cell lines and immunocompromised strains. This limitation requires the use of immunosuppressive therapy, which can lead to severe and difficult-to-manage complications,¹⁵⁸ while also altering the effects of drugs or interventions. This limitation is effectively addressed by murine xenograft models, which utilize either syngeneic tumour cell implantation or human tumour cell inoculation in immunocompromised mice, as well as genetically engineered mice with spontaneous tumour growth (Table 2). Additionally, *in vitro* heart-tumour chips offer a promising alternative for studying these tumour–heart–drug interactions.

The immune system also plays a critical role in interacting with the myocardium and contributing to the development of CTRCD. However, immune cells are still rarely incorporated into *in vitro* models of CTRCD, highlighting the need for a more comprehensive use of cardiac slices and 2D and 3D co-culture systems including inflammatory cells.⁷⁷

Finally, as previously pointed out, recognizing early warning signs in the translational phase is essential for drug development in cardiovascular medicine. Ignoring these signs could become costly and dangerous when candidate therapies move into clinical evaluation.¹⁴³

In conclusion, growing evidence suggests that genetic,¹⁵⁹ gender,¹⁵¹ and ethnic¹⁵⁵ factors influence predisposition to CTRCD. While groundbreaking research using genomic editing of human

iPSC have started validating significant variants identified in candidate-gene and genome-wide association studies,^{160–162} gender and ethnic considerations have been largely overlooked in preclinical modelling of CTRCD. Therefore, future research should address these neglected aspects by increasing the inclusion of both male and female animals as well as utilizing *in vitro* model systems with cells from patients of diverse ethnic backgrounds.

Impact of limitations on regulatory paths and full development process

Pathways for drug approval are becoming increasingly complex, with both the European Medicines Agency and US Food and Drug Administration (FDA) focusing on accelerated approval pathways. Regulatory agencies face the challenge of balancing the demand for rapid access to new and effective drugs with the need for comprehensive safety and efficacy data prior to patient use. Unfortunately, drug development regulations in different therapeutic areas are far from harmonized, and this affects the number of new chemical entities reaching the market.¹⁶³ As an example, under the FDA's new 'breakthrough therapy', designation of the approvals granted from 1987 to 2017, approximately 50% were for oncology, 20% for infectious diseases, and none for CVDs.¹⁶⁴ There is no doubt that many CVDs, including the acute or chronic consequences of cancer treatment, should be considered severely-debilitating or life-threatening diseases, and it stands to reason that the approval of severely-debilitating or life-threatening disease therapeutics should parallel that of advanced oncology drugs. However, although the FDA provides regulatory guidelines for advanced cancer therapies,¹⁶⁵ there is minimal FDA guidance for the early development of drugs for severely-debilitating or life-threatening diseases for which there are few available therapeutic options. Consequently, from an industry business perspective, drug development projects in CVDs could become less attractive, which may contribute to the diversion of funds to other areas deemed more lucrative. A negative trend can be already seen in the percentages of new CVD treatment over the total approval by FDA in the latest decades: between 2000 and 2017, in fact, the approved CVD therapies over the total of the new approvals by FDA declined from 9.09% to 5.29%.¹⁶⁶ Industry-funded development research will just focus elsewhere if the trend continues.

From an academic perspective, several factors can hinder or slow down the development process, including limited awareness and infrastructure for patenting, spin-offs, and industry partnerships. Researchers often prioritize publishing over commercial applications, missing opportunities to translate innovations into practical solutions, particularly in fields like life sciences where intellectual property protection is crucial. Many lack knowledge of the patenting process, including optimal timing and procedures. While universities have technology transfer offices, their effectiveness varies, and structured pathways for spin-offs are often lacking. Early industry collaborations, which provide funding and expertise, are underutilized due to limited connections and concerns over research independence. Strengthening academia–industry networks, improving intellectual property training, and fostering

Table 3 Gaps in preclinical modelling and recommendations for future research

Topic	Gap in modelling	Recommendation for future research	Examples/reference
Late-onset manifestation of CTRCD (especially anthracyclines and radiation-induced cardiotoxicity)	<i>In vivo</i> : study of CTRCD in animals treated with a single, high dose of anticancer drug/radiation, usually leading to early and high mortality rates due to non-cardiac toxicity.	Implement treatment protocols that closely replicate human therapeutic regimens in terms of administration route, number of cycles, dosage and pharmacokinetic profile.	Establishment of a murine model that provokes modest and progressive doxorubicin cardiotoxicity without constitutional symptoms, reminiscent of the effects seen in patients ¹⁴⁴
	<i>In vitro</i> : study of CTRCD in cells/slices/EHT following one dose of anticancer drug/radiation.	Implement repeated treatment schemes.	4-hit model ⁷⁶ 2-hit model ¹⁴⁵
Crosstalk between tumour cells and the myocardium in CTRCD	Study of CTRCD in non-tumour-bearing animals.	Perform treatment with anticancer drug/radiation in animals with tumour xenografts (syngeneic models or PDX in SCID mice) or in genetically engineered mice with spontaneous tumour growth.	AML cells killed by doxorubicin release IL-1 α leading to cardiac metabolic rewiring ¹⁴⁶ ICI-induced LV dysfunction detected in tumour-bearing mice only. ⁹⁴
	Study of CTRCD in non-tumour comprising <i>in vitro</i> model systems.	Development and use of heart-tumour chips.	A heart-breast cancer-on-a-chip platform for disease modelling and monitoring of cardiotoxicity induced by cancer chemotherapy. ¹⁴⁷
Interaction of the immune system with the myocardium (and/or the tumour) in CTRCD	<i>In vivo</i> : study of CTRCD in immunodeficient animals.	<i>In vivo</i> : use of immunocompetent animals.	Targeting early stages of cardiotoxicity from anti-PD-1 ICI therapy. ⁹⁴
	<i>In vitro</i> : study of CTRCD in cells/slices/EHT in the absence of immune cells.	<i>In vitro</i> : implement 2D and 3D co-culture systems, including immune cells in addition to cardiac cells. Use of cardiac slices.	Effects of co-cultured macrophages on iPSC proliferation, cardiac differentiation, and maturation. ⁷⁷
Intra- and inter-species phenotypic variability of anticancer therapy cardiotoxicity	<i>Intraspecies variability</i> : the severity of cardiotoxicity phenotypes may vary based on the genetic background of the animal strain used, with potential differences even between the N and J substrains of C57BL/6 mice due to genetic variations.	Select sensitive strain backgrounds and/or optimizing treatment doses/regimens based on the sensitivity of the selected strain.	Strain background differences in the severity of cardiovascular phenotypes. ^{148,149}
	<i>Interspecies variability</i> : rodents have been used to study trastuzumab cardiotoxicity although the drug does not bind rat or mouse <i>ErbB2/Neu</i> , generating results hard to interpret (off-target rather than on-target cardiotoxicity).	Select relevant binding species (e.g. human pluripotent stem cells or humanized models like transgenic animals expressing human HER-2) to study HER-2 inhibitors cardiotoxicity.	Trastuzumab, and by extension trastuzumab drug conjugates, do not bind rat or mouse <i>Neu</i> . ⁹²

Table 3 (Continued)

Topic	Gap in modelling	Recommendation for future research	Examples/reference
Sex-specific cardiovascular risks of cancer and its therapies	Sex-based differences of cancer and cancer therapy cardiotoxicity are underexplored.	<i>In vivo</i> : evaluate anticancer therapies in both male and female animals of the same background, or in the sex specific to the cancer. Report the sex of the animals (ARRIVE guidelines). ¹⁵⁰ <i>In vitro</i> : evaluate anticancer therapies in both male and female <i>in vitro</i> model systems, or in the sex specific to the cancer. Report the sex of the cells/ <i>in vitro</i> model used. Use of isogenic male vs. female iPSC-derived CMs.	Sex-specific cardiovascular risks of cancer and its therapies. ^{151–153}
Genetic predisposition to CTRCD	Need for model systems reflecting genetic predisposition to CTRCD.	Use of genomic editing of human iPSC for validation of significant variants identified in CGAS and GWAS studies.	Human <i>in vitro</i> models for assessing the genomic basis of CTRCD. ¹⁵⁴
Racial/ethnic disparity in CTRCD	Racial/ethnic disparity in the use of cells/ <i>in vitro</i> systems for CTRCD.	Use of <i>in vitro</i> model systems with cells from patients with different ethnic background.	Racial disparity in anthracycline-induced CTRCD. ¹⁵⁵

2D, two-dimensional; 3D, three-dimensional; AML, acute myeloid leukaemia; ARRIVE, animal research, reporting of *in vivo* experiment; CGAS, candidate gene association study; CM, cardiomyocyte; CTRCD, cancer therapy-related cardiac dysfunction; EHT, engineered heart tissue; GWAS, genome-wide association study; HER-2, human epidermal growth factor receptor 2; ICI, immune checkpoint inhibitor; IL-1 α , interleukin-1 α ; iPSC, induced pluripotent stem cell; LV, left ventricular; PD-1, programmed cell death protein 1; PDX, patient-derived xenograft; SCID, severe combined immunodeficiency.

entrepreneurial support would help streamline the development process.^{167,168}

For these reasons, we cannot emphasize enough the need for continuous and open discussions between translational researchers, trialists, pharmaceutical industry representatives and regulatory authorities to ensure that cardio-oncology is properly positioned in the regulatory compass and, overall, in the world health-economic compass.

Conclusions and future directions

Over the past decade, cancer therapy has experienced a paradigm shift from non-specific cytotoxic chemotherapies, such as anthracyclines, to more targeted treatments including monoclonal antibodies, TKIs, ICIs, and cell therapies. While these advancements have led to better clinical outcomes for patients with both solid and haematological malignancies, they also come with a broad spectrum of side effects, including cardiac toxicity. Consequently, the development of new anticancer drugs presents the challenge of uncovering the mechanisms underlying the relative CTRCD, which necessitates the creation of effective preclinical disease models. Although a combination of preclinical *in vitro* and *in vivo* models across various species has provided a comprehensive understanding of the cardiotoxicity associated with conventional anticancer therapies, the cardiac effects of some newer treatments are only

beginning to be explored, while others remain largely unexplored and warrant future research. For instance, further investigation is required into the cardiac impact of cyclin-dependent kinase 4/6 inhibitors¹⁶⁹ or the antibody-drug conjugates – an innovative and evolving class of antineoplastic agents that combine monoclonal antibodies with biologically active drugs.¹⁵⁶ Furthermore, the potential interactions among cancer therapies in multimodal regimens have been insufficiently addressed in preclinical modelling, and thus warrant the attention of future studies.

The development of new immunotherapies, coupled with the growing recognition of the pivotal role of the immune system in CTRCD, and as a common denominator in cancer and HF,¹⁷⁰ underscores the importance of using immunocompetent animals and multicellular 2D and 3D model systems, encompassing, in addition to cardiomyocytes, fibroblasts (which possess inflammatory properties)¹⁷¹ and immune cells. The mutual interaction between the tumour and the heart further supports the use of tumour-bearing animals⁹⁴ and of heart-tumour-on-a-chip platforms.¹⁴⁷ Genetic predisposition to¹⁵⁴ and sex-based differences in CTRCD¹⁵¹ also highlight the importance of genomic editing of human iPSC for validation of significant genetic variants associated with CTRCD and the testing of anticancer therapies in both male and female cells/model systems.¹⁵² Racial/ethnic disparity in CTRCD emphasizes the need for *in vitro* model systems with cells from patients with different ethnical background.¹⁵⁵ Finally, in

view of personalized medicine, patient-derived model systems may serve as predictive models of CTRCD.¹⁷² Beyond the recognized use of patient-specific iPSC-CM and multicellular model systems, the use of cardiac slices derived from endomyocardial biopsies holds promise.

In conclusion, the emergence of novel anticancer drugs, the recognition of the role of the immune system in CTRCD, our increased understanding of the reciprocal relationship between the heart and cancer, and the growing appreciation for sex-based and ethnic disparities of CTRCD are driving the development of novel, improved preclinical model systems and approaches to close the gap between preclinical models and clinical observations in patients.

Acknowledgement

Open access publishing facilitated by Università degli Studi di Torino, as part of the Wiley - CRUI-CARE agreement.

Funding

A.G. was supported by grants from the Italian Ministry of Health (GR-2021-12371950) and the Italian Ministry of Education, Universities and Research (PRIN 2022 20223YPL49; PRIN PNRR 2022 P2022ZB72T). P.A. was supported by PNRR-MAD-2022-12376632 (CUP C63C22001360006) funded by the European Union - Next Generation EU - NRRP M6C2 - Investment 2.1 Enhancement and strengthening of biomedical research in the NHS. A.A. is supported by NIH R01HL163172, R01HL166541, and R01HL157530; a sponsored research agreement with Genentech; and a HESI Thrive Award. R.A.d.B. is supported by the Netherlands Heart Foundation (grants 2020B005 and 01-003-2022-0358) and by the European Research Council (ERC CoG 818715). A.Go. was supported by the Spanish Ministry of Science, Innovation and Universities (Instituto de Salud Carlos III: CB16/11/00483 and PI21/00946 co-financed with ERDF funds and CIBERCV InCare project). B.I. is supported by grants from the European Commission (ERC-CoG 819775 and H2020-HEALTH 945118), the Spanish Ministry of Science and Innovation (PID2022-140176OB-I00), La Caixa Foundation (HR22-00533), and the Red Madrileña de Nanomedicina en Imagen Molecular -Comunidad de Madrid (P2022/BMD-7403 RENIM-CM). A.R.L. is supported by the Fondation Leducq Network of Excellence in Cardio-Oncology, the Royal Brompton Cardio-Oncology Centre of Excellence and The Big Heart Foundation. K.S. is supported by grants from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (MODVASC, grant agreement No 759248), the German Research Foundation DFG (CRC1366 C07, project number 394046768), the Health+Life Science Alliance Heidelberg Mannheim GmbH and the Biotechnology and Biological Sciences Research Council (BBSRC) of the UK Research and Innovation (UKRI). K.S.B. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, Project 471241922, CRC1213-B07 and the research training group RTG2824). C.G.T. is supported by the Italian Ministry of Health (PNRR-MAD-2022-12376632 and RF-2016-02362988). T.T. was supported by the ERC Advanced Grant REVERSE and the Deutsche Forschungsgemeinschaft (CRC1470). P.v.d.M. is supported by a grant from the European Research Council (ERC CoG 101045236, DISSECT-HF). S.V.L. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, CRC-1470-A07 and Project 536819681) and the Deutsche Krebshilfe (Project 70115119).

Conflict of interest: A.G. is co-founder and shareholder of Kither Biotech, a pharmaceutical company focused on the development of PI3K inhibitors for airway diseases not in conflict with the content of this manuscript. P.A. received speaker and/or advisory fees from Boehringer Ingelheim, AstraZeneca, Bayer, Daiichi-Sankyo, MSD, Janssen, and Gossamer Bio, outside the scope of this work. A.A. serves as the PI for a sponsored research agreement with Genentech, receives royalties from patent #US20210163495A1 (Mass General Brigham), and sits on the Board of Directors for Corventum, Inc. The institution of R.A.d.B. has received research grants and/or fees from Alnylam, AstraZeneca, Abbott, Bristol-Myers Squibb, Cardior Pharmaceuticals GmbH, Novo Nordisk, and Roche; R.A.d.B. has had speaker engagements with and/or received fees from and/or served on an advisory board for Abbott, AstraZeneca, Bristol Myers Squibb, Cardior Pharmaceuticals GmbH, Novo Nordisk, and Roche; travel support from Abbott, Cardior Pharmaceuticals GmbH, and Novo Nordisk. D.F. reports speaker honoraria or consultation fees from Abbott Laboratories, AstraZeneca, Bayer, Boehringer Ingelheim, Leo, Roche Diagnostics and Viatrix, outside the submitted work. S.H. receives personal fees for independent scientific advice on early development in the field of heart failure for AstraZeneca, Ribocure, and CSL Behring, and research support from AstraZeneca and CSL Behring. T.L.F. has received speaker, advisory board or consultancy fees from AstraZeneca, Bristol Myers Squibb, Roche, Janssens-Cilag Ltd, Eisai Ltd, Myocardial Solutions, Philips, Bayer, Daiichi-Sankyo and Beigene. A.R.L. has received speaker, advisory board or consultancy fees and/or research grants from Pfizer, Novartis, Servier, AstraZeneca, Bristol Myers Squibb, GSK, Astellas, Amgen, Takeda, Roche, Janssens-Cilag Ltd, Clinigen Group, Eli Lilly, Eisai Ltd, Ferring Pharmaceuticals, Boehringer Ingelheim, Akcea Therapeutics, Myocardial Solutions, iOWNA Health and Heartfelt Technologies Ltd. K.S.B. received research support from Novartis, BionTECH, and speaker honoraria from Novartis. C.G.T. has received honoraria or consultancy fees from VivaLyfe, Univers Formazione, Solaris, Summeet, AstraZeneca, Medtronic, and Myocardial Solutions, and funding from Amgen and MSD, outside the submitted work, and is listed as an inventor of two patents related to heart failure. T.T. is founder and CSO/CMO of Cardior Pharmaceuticals GmbH, a wholly-owned subsidiary of Novo Nordisk Europe A/S. The UMCG, which employs P.v.d.M., received consultancy fees and/or grants from Novartis, Pharmacosmos, Vifor Pharma, AstraZeneca, Pfizer, Pharma Nord, BridgeBio, Novo Nordisk, Daiichi Sankyo, Boehringer Ingelheim and Ionis. M.M. has received consulting honoraria from Abbott Structural Heart, AstraZeneca, Bayer, Boehringer Ingelheim, Edwards Lifesciences, Novo Nordisk, Pharmacosmos, Roche Diagnostics in the last 3 years. All other authors have nothing to disclose.

References

1. Lyon AR, Lopez-Fernandez T, Couch LS, Asteggiano R, Aznar MC, Bergler-Klein J, et al. 2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J Cardiovasc Imaging* 2022;23:e333-e465. <https://doi.org/10.1093/ehjci/jeac106>
2. Asnani A, Moslehi JJ, Adhikari BB, Baik AH, Beyer AM, de Boer RA, et al.; American Heart Association Council on Basic Cardiovascular Sciences; Cardio-Oncology Science Subcommittee of Council on Genomic and Precision Medicine and Council on Clinical Cardiology; Council on Peripheral Vascular Disease; and Council on Arteriosclerosis, Thrombosis and Vascular Biology. Preclinical models of cancer therapy-associated cardiovascular toxicity: A scientific statement from the American Heart Association. *Circ Res* 2021;129:e21-e34. <https://doi.org/10.1161/RES.0000000000000473>
3. Larsen CM, Garcia Arango M, Dasari H, Arciniegas Calle M, Adjei E, Rico Mesa J, et al. Association of anthracycline with heart failure in patients treated for breast cancer or lymphoma, 1985-2010. *JAMA Netw Open* 2023;6:e2254669. <https://doi.org/10.1001/jamanetworkopen.2022.54669>

4. Herrmann J, Lenihan D, Armenian S, Barac A, Blaes A, Cardinale D, et al. Defining cardiovascular toxicities of cancer therapies: An International Cardio-Oncology Society (IC-OS) consensus statement. *Eur Heart J* 2021;**43**:280–299. <https://doi.org/10.1093/eurheartj/ehab674>
5. Salloum FN, Tocchetti CG, Ameri P, Ardehali H, Asnani A, de Boer RA, et al. International Cardio-Oncology Society. Priorities in cardio-oncology basic and translational science: GCOS 2023 Symposium Proceedings: JACC: CardioOncology state-of-the-art review. *JACC CardioOncol* 2023;**5**:715–731. <https://doi.org/10.1016/j.jacc.2023.08.003>
6. Farhad, H, Staziaki, PV, Addison D, Coelho-Filho OR, Shah RV, Mitchell RN, et al. Characterization of the changes in cardiac structure and function in mice treated with anthracyclines using serial cardiac magnetic resonance imaging. *Circ Cardiovasc Imaging* 2016;**9**: e003584. <https://doi.org/10.1161/CIRCIMAGING.115.003584>
7. Tocchetti CG, Ameri P, de Boer RA, D'Alessandra Y, Russo M, Sorriento D, et al. Cardiac dysfunction in cancer patients: Beyond direct cardiomyocyte damage of anticancer drugs: Novel cardio-oncology insights from the joint 2019 meeting of the ESC Working Groups of Myocardial Function and Cellular Biology of the Heart. *Cardiovasc Res* 2020;**116**:1820–1834. <https://doi.org/10.1093/cvr/cvaa222>
8. Ozcelik C, Erdmann B, Pilz B, Wetschreck N, Britsch S, Hübner N, et al. Conditional mutation of the ErbB2 (HER2) receptor in cardiomyocytes leads to dilated cardiomyopathy. *Proc Natl Acad Sci USA* 2002;**99**:8880–8885. <https://doi.org/10.1073/pnas.122249299>
9. Belmonte F, Das S, Saha-Shah P, Sivakumaran V, Stanley B, Guo X, et al. ErbB2 overexpression upregulates antioxidant enzymes, reduces basal levels of reactive oxygen species, and protects against doxorubicin cardiotoxicity. *Am J Physiol Heart Circ Physiol* 2015;**309**:H1271–1280. <https://doi.org/10.1152/ajpheart.00517.2014>
10. Lyra-Leite DM, Burridge PV. Pluripotent stem cell modeling of anticancer therapy-induced cardiotoxicity. *Curr Cardiol Rep* 2020;**22**:56. <https://doi.org/10.1007/s11886-020-01325-x>
11. Kitani T, Ong SG, Lam CK, Rhee JW, Zhang JZ, Oikonomopoulos A, et al. Human-induced pluripotent stem cell model of trastuzumab-induced cardiac dysfunction in patients with breast cancer. *Circulation* 2019;**139**:2451–2465. <https://doi.org/10.1161/CIRCULATIONAHA.118.037357>
12. Xu Z, Gao Z, Fu H, Zeng Y, Jin Y, Xu B, et al. PTX3 from vascular endothelial cells contributes to trastuzumab-induced cardiac complications. *Cardiovasc Res* 2023;**119**:1250–1264. <https://doi.org/10.1093/cvr/cvad012>
13. Medeiros BC, Possick J, Fradley M. Cardiovascular, pulmonary, and metabolic toxicities complicating tyrosine kinase inhibitor therapy in chronic myeloid leukemia: Strategies for monitoring, detecting, and managing. *Blood Rev* 2018;**32**:289–299. <https://doi.org/10.1016/j.blre.2018.01.004>
14. Yan G, Han Z, Kwon Y, Jousma J, Nukala SB, Prosser BL, et al. Integrated stress response potentiates ponatinib-induced cardiotoxicity. *Circ Res* 2024;**134**:482–501. <https://doi.org/10.1161/CIRCRESAHA.123.323683>
15. Tousif S, Singh AP, Umbarkar P, Galindo C, Wheeler N, Toro Cora A, et al. Ponatinib drives cardiotoxicity by S100A8/A9-NLRP3-IL-1 β mediated inflammation. *Circ Res* 2023;**132**:267–289. <https://doi.org/10.1161/CIRCRESAHA.122.321504>
16. Truitt R, Mu A, Corbin EA, Vite A, Brandimarto J, Ky B, et al. Increased afterload augments sunitinib-induced cardiotoxicity in an engineered cardiac microtissue model. *JACC Basic Transl Sci* 2018;**3**:265–276. <https://doi.org/10.1016/j.jacbs.2017.12.007>
17. Xiao L, Salem JE, Clauss S, Hanley A, Bapat A, Hulsmans M, et al. Ibrutinib-mediated atrial fibrillation attributable to inhibition of C-terminal Src kinase. *Circulation* 2020;**142**:2443–2455. <https://doi.org/10.1161/CIRCULATIONAHA.120.049210>
18. Quartermaine C, Ghazi SM, Yasin A, Awan FT, Fradley M, Wiczer T, et al. Cardiovascular toxicities of BTK inhibitors in chronic lymphocytic leukemia: JACC: CardioOncology state-of-the-art review. *JACC CardioOncol* 2023;**5**:570–590. <https://doi.org/10.1016/j.jacc.2023.09.002>
19. Mitchell JD, Cehic DA, Morgia M, Bergom C, Toohey J, Guerrero PA, et al. Cardiovascular manifestations from therapeutic radiation: A multidisciplinary expert consensus statement from the International Cardio-Oncology Society. *JACC CardioOncol* 2021;**3**:360–380. <https://doi.org/10.1016/j.jacc.2021.06.003>
20. Lee CL, Lee JW, Daniel AR, Holbrook M, Hasapis S, Wright AO, et al. Characterization of cardiovascular injury in mice following partial-heart irradiation with clinically relevant dose and fractionation. *Radiother Oncol* 2021;**157**:155–162. <https://doi.org/10.1016/j.radonc.2021.01.023>
21. Sag CM, Wolff HA, Neumann K, Opiela MK, Zhang J, Steuer F, et al. Ionizing radiation regulates cardiac Ca handling via increased ROS and activated CaMKII. *Basic Res Cardiol* 2013;**108**:385. <https://doi.org/10.1007/s00395-013-0385-6>
22. Morelli MB, Bongiovanni C, Da Pra S, Miano C, Sacchi F, Lauriola M, et al. Cardiotoxicity of anticancer drugs: Molecular mechanisms and strategies for cardioprotection. *Front Cardiovasc Med* 2022;**9**:847012. <https://doi.org/10.3389/fcvm.2022.847012>
23. Thuny F, Naidoo J, Neilan TG. Cardiovascular complications of immune checkpoint inhibitors for cancer. *Eur Heart J* 2022;**43**:4458–4468. <https://doi.org/10.1093/eurheartj/ehac456>
24. Ball S, Ghosh RK, Wongsangsak S, Bandyopadhyay D, Ghosh GC, Aronow WS, et al. Cardiovascular toxicities of immune checkpoint inhibitors. *J Am Coll Cardiol* 2019;**74**:1714–1727. <https://doi.org/10.1016/j.jacc.2019.07.079>
25. Robert C, Schachter J, Long GV, Arance A, Grob JJ, Mortier L, et al.; KEYNOTE-006 Investigators. Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med* 2015;**372**:2521–2532. <https://doi.org/10.1056/NEJMoa1503093>
26. Wolchok JD, Chiarion-Sileni V, Gonzalez R, Rutkowski P, Grob J-J, Cowey CL, et al. Overall survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2017;**377**:1345–1356. <https://doi.org/10.1056/NEJMoa1709684>
27. Martins F, Sofiya L, Sykiotis GP, Lamine F, Maillard M, Fraga M, et al. Adverse effects of immune-checkpoint inhibitors: Epidemiology, management and surveillance. *Nat Rev Clin Oncol* 2019;**16**:563–580. <https://doi.org/10.1038/s41571-019-0218-0>
28. D'Souza M, Nielsen D, Svane IM, D'Souza M, Iversen K, Rasmussen PV, et al. The risk of cardiac events in patients receiving immune checkpoint inhibitors: A nationwide Danish study. *Eur Heart J* 2020;**42**:1621–1631. <https://doi.org/10.1093/eurheartj/ehaa884>
29. Moslehi JJ, Salem JE, Sosman JA, Lebrun-Vignes B, Johnson DB. Increased reporting of fatal immune checkpoint inhibitor-associated myocarditis. *Lancet* 2018;**391**:933. [https://doi.org/10.1016/S0140-6736\(18\)30533-6](https://doi.org/10.1016/S0140-6736(18)30533-6)
30. Rubio-Infante N, Ramirez-Flores YA, Castillo EC, Lozano O, Garcia-Rivas G, Torre-Amione G. Cardiotoxicity associated with immune checkpoint inhibitor therapy: A meta-analysis. *Eur J Heart Fail* 2021;**23**:1739–1747. <https://doi.org/10.1002/ejhf.2289>
31. Tocchetti CG, Farmakis D, Koop Y, Andres MS, Couch LS, Formisano L, et al. Cardiovascular toxicities of immune therapies for cancer – a scientific statement of the Heart Failure Association (HFA) of the ESC and the ESC Council of Cardio-Oncology. *Eur J Heart Fail* 2024;**26**:2055–2076. <https://doi.org/10.1002/ejhf.3340>
32. Gergely TG, Drobni ZD, Kallikourdis M, Zhu H, Meijers WC, Neilan TG, et al. Immune checkpoints in cardiac physiology and pathology: Therapeutic targets for heart failure. *Nat Rev Cardiol* 2024;**21**:443–462. <https://doi.org/10.1038/s41569-023-00986-9>
33. Grabie N, Lichtman AH, Padera R. T cell checkpoint regulators in the heart. *Cardiovasc Res* 2019;**115**:869–877. <https://doi.org/10.1093/cvr/cvz025>
34. Axelrod ML, Meijers WC, Screever EM, Qin J, Carroll MG, Sun X, et al. T cells specific for α -myosin drive immunotherapy-related myocarditis. *Nature* 2022;**611**:818–826. <https://doi.org/10.1038/s41586-022-05432-3>
35. Wei SC, Meijers WC, Axelrod ML, Anang NAS, Screever EM, Wescott EC, et al. A genetic mouse model recapitulates immune checkpoint inhibitor-associated myocarditis and supports a mechanism-based therapeutic intervention. *Cancer Discov* 2021;**11**:614–625. <https://doi.org/10.1158/2159-8290.CD-20-0856>
36. Cortellino S, Quagliarriello V, Delfanti G, Blažević O, Chiodoni C, Maurea N, et al. Fasting mimicking diet in mice delays cancer growth and reduces immunotherapy-associated cardiovascular and systemic side effects. *Nat Commun* 2023;**14**:5529. <https://doi.org/10.1038/s41467-023-41066-3>
37. Fenioux C, Abbar B, Boussouar S, Bretagne M, Power JR, Moslehi JJ, et al. Thymus alterations and susceptibility to immune checkpoint inhibitor myocarditis. *Nat Med* 2023;**29**:3100–3110. <https://doi.org/10.1038/s41591-023-02591-2>
38. Irvine DJ, Maus MV, Mooney DJ, Wong WW. The future of engineered immune cell therapies. *Science* 2022;**378**:853–858. <https://doi.org/10.1126/science.abq6990>
39. Baker DJ, Arany Z, Baur JA, Epstein JA, June CH. CAR T therapy beyond cancer: The evolution of a living drug. *Nature* 2023;**619**:707–715. <https://doi.org/10.1038/s41586-023-06243-w>
40. Fajgenbaum DC, June CH. Cytokine storm. *N Engl J Med* 2020;**383**:2255–2273. <https://doi.org/10.1056/NEJMr2026131>
41. Lee DW, Santomasso BD, Locke FL, Ghobadi A, Turtle CJ, Brudno JN, et al. ASTCT consensus grading for cytokine release syndrome and neurologic toxicity associated with immune effector cells. *Biol Blood Marrow Transplant* 2019;**25**:625–638. <https://doi.org/10.1016/j.bbmt.2018.12.758>
42. Jamal A, Ho N, Braun A, Adabi E, Thalheimer FB, Buchholz CJ. Early induction of cytokine release syndrome by rapidly generated CAR T cells in preclinical models. *EMBO Mol Med* 2024;**16**:784–804. <https://doi.org/10.1038/s44321-024-00055-9>
43. Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M. CAR T cell-induced cytokine release syndrome is mediated by macrophages and

- abated by IL-1 blockade. *Nat Med* 2018;**24**:731–738. <https://doi.org/10.1038/s41591-018-0041-7>
44. Li X, Gong N, Tian F, Zhang S, Zhang Y, Wang Y, et al. Suppression of cytokine release syndrome during CAR-T-cell therapy via a subcutaneously injected interleukin-6-adsorbing hydrogel. *Nat Biomed Eng* 2023;**7**:1129–1141. <https://doi.org/10.1038/s41551-023-01084-4>
 45. Gelfo V, Romaniello D, Mazzeschi M, Sgarzi M, Grilli G, Morselli A, et al. Roles of IL-1 in cancer: From tumor progression to resistance to targeted therapies. *Int J Mol Sci* 2020;**21**:6009. <https://doi.org/10.3390/ijms21176009>
 46. Lee DW, Gardner R, Porter DL, Louis CU, Ahmed N, Jensen M, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood* 2014;**124**:188–195. <https://doi.org/10.1182/blood-2014-05-552729>
 47. Alvi RM, Frigalet MJ, Fradley MG, Jain MD, Mahmood SS, Awadalla M, et al. Cardiovascular events among adults treated with chimeric antigen receptor T-cells (CAR-T). *J Am Coll Cardiol* 2019;**74**:3099–3108. <https://doi.org/10.1016/j.jacc.2019.10.038>
 48. Salem JE, Ederhy S, Lebrun-Vignes B, Moslehi JJ. Cardiac events associated with chimeric antigen receptor T-cells (CAR-T): A VigiBase perspective. *J Am Coll Cardiol* 2020;**75**:2521–2523. <https://doi.org/10.1016/j.jacc.2020.02.070>
 49. Lefebvre B, Kang Y, Smith AM, Frey NV, Carver JR, Scherrer-Crosbie M. Cardiovascular effects of CAR T cell therapy. *JACC CardioOncol* 2020;**2**:193–203. <https://doi.org/10.1016/j.jacc.2020.04.012>
 50. Ganatra S, Redd R, Hayek SS, Parikh R, Azam T, Yanik GA, et al. Chimeric antigen receptor T-cell therapy-associated cardiomyopathy in patients with refractory or relapsed non-Hodgkin lymphoma. *Circulation* 2020;**142**:1687–1690. <https://doi.org/10.1161/CIRCULATIONAHA.120.048100>
 51. Goldman A, Maor E, Bomze D, Liu JE, Herrmann J, Fein J, et al. Adverse cardiovascular and pulmonary events associated with chimeric antigen receptor T-cell therapy. *J Am Coll Cardiol* 2021;**78**:1800–1813. <https://doi.org/10.1016/j.jacc.2021.08.044>
 52. Qi K, Yan Z, Cheng H, Chen W, Wang Y, Wang X, et al. An analysis of cardiac disorders associated with chimeric antigen receptor T cell therapy in 126 patients: A single-centre retrospective study. *Front Oncol* 2021;**11**:691064. <https://doi.org/10.3389/fonc.2021.691064>
 53. Lee DH, Kumar A, Mohammed T, Peres LC, Alsina M, Bachmeier C, et al. Cardiac events after standard of care idexcabtagene vicleucel for relapsed and refractory multiple myeloma. *Blood Adv* 2023;**7**:4247–4257. <https://doi.org/10.1182/bloodadvances.2023009766>
 54. Lefebvre B, Kang Y, Vakilpour A, Onoue T, Frey NV, Brahmabhatt P, et al. Incidence of MACE in patients treated with CAR-T cell therapy: A prospective study. *JACC CardioOncol* 2023;**5**:747–754. <https://doi.org/10.1016/j.jacc.2023.07.009>
 55. Mahmood SS, Riedell PA, Feldman S, George G, Sansoterra SA, Althaus T, et al. Biomarkers and cardiovascular outcomes in chimeric antigen receptor T-cell therapy recipients. *Eur Heart J* 2023;**44**:2029–2042. <https://doi.org/10.1093/eurheartj/ehad117>
 56. Korell F, Entenmann L, Romann S, Giannitsis E, Schmitt A, Müller-Tidow C, et al. Evaluation of all-cause mortality and cardiovascular safety in patients receiving chimeric antigen receptor T cell therapy: A prospective cohort study. *EClinicalMedicine* 2024;**69**:102504. <https://doi.org/10.1016/j.eclinm.2024.102504>
 57. Norelli M, Camisa B, Barbiera G, Falcone L, Purevdorj A, Genua M, et al. Monocyte-derived IL-1 and IL-6 are differentially required for cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nat Med* 2018;**24**:739–748. <https://doi.org/10.1038/s41591-018-0036-4>
 58. Pathan N, Hemingway CA, Alizadeh AA, Stephens AC, Boldrick JC, Oragui EE, et al. Role of interleukin 6 in myocardial dysfunction of meningococcal septic shock. *Lancet* 2004;**363**:203–209. [https://doi.org/10.1016/S0140-6736\(03\)15326-3](https://doi.org/10.1016/S0140-6736(03)15326-3)
 59. Shafer P, Kelly LM, Hoyos V. Cancer therapy with TCR-engineered T cells: Current strategies, challenges, and prospects. *Front Immunol* 2022;**13**:835762. <https://doi.org/10.3389/fimmu.2022.835762>
 60. Cameron BJ, Gerry AB, Dukes J, Harper JV, Kannan V, Bianchi FC, et al. Identification of a titin-derived HLA-A1-presented peptide as a cross-reactive target for engineered MAGE A3-directed T cells. *Sci Transl Med* 2013;**5**:197ra103. <https://doi.org/10.1126/scitranslmed.3006034>
 61. Linette GP, Stadtmayer EA, Maus MV, Rapoport AP, Levine BL, Emery L, et al. Cardiovascular toxicity and titin cross-reactivity of affinity-enhanced T cells in myeloma and melanoma. *Blood* 2013;**122**:863–871. <https://doi.org/10.1182/blood-2013-03-490565>
 62. Burridge PW, Li YF, Matsa E, Wu H, Ong SG, Sharma A, et al. Human induced pluripotent stem cell-derived cardiomyocytes recapitulate the predilection of breast cancer patients to doxorubicin-induced cardiotoxicity. *Nature Med* 2016;**22**:547–556. <https://doi.org/10.1038/nm.4087>
 63. Sang L, Zhou Z, Luo S, Zhang Y, Qian H, Zhou Y, et al. An in silico platform to predict cardiotoxicity risk of anti-tumor drug combination with hiPSC-CMs based in vitro study. *Pharm Res* 2024;**41**:247–262. <https://doi.org/10.1007/s11095-023-03644-4>
 64. Pinheiro EA, Fetterman KA, Burridge PW. hiPSCs in cardio-oncology: Deciphering the genomics. *Cardiovasc Res* 2019;**115**:935–948. <https://doi.org/10.1093/cvr/cvz018>
 65. Karhu ST, Kinnunen SM, Tolli M, Valimäki MJ, Szabo Z, Talman V, et al. GATA4-targeted compound exhibits cardioprotective actions against doxorubicin-induced toxicity in vitro and in vivo: Establishment of a chronic cardiotoxicity model using human iPSC-derived cardiomyocytes. *Arch Toxicol* 2020;**94**:2113–2130. <https://doi.org/10.1007/s00204-020-02711-8>
 66. Maillet A, Tan K, Chai X, Sadananda SN, Mehta A, Ooi J, et al. Modeling doxorubicin-induced cardiotoxicity in human pluripotent stem cell derived-cardiomyocytes. *Sci Rep* 2016;**6**:25333. <https://doi.org/10.1038/srep25333>
 67. Chatterjee S, Hofer T, Costa A, Lu D, Batkai S, Gupta SK, et al. Telomerase therapy attenuates cardiotoxic effects of doxorubicin. *Mol Ther* 2021;**29**:1395–1410. <https://doi.org/10.1016/j.yjth.2020.12.035>
 68. Haupt LP, Rebs S, Maurer W, Hübscher D, Tiburcy M, Pabel S, et al. Doxorubicin induces cardiotoxicity in a pluripotent stem cell model of aggressive B cell lymphoma cancer patients. *Basic Res Cardiol* 2022;**117**:13. <https://doi.org/10.1007/s00395-022-00918-7>
 69. Reinal I, Ontoria-Oviedo I, Selva M, Casini M, Peiró-Molina E, Fambuena-Santos C, et al. Modeling cardiotoxicity in pediatric oncology patients using patient-specific iPSC-derived cardiomyocytes reveals down-regulation of cardioprotective microRNAs. *Antioxidants (Basel)* 2023;**12**:1378. <https://doi.org/10.3390/antiox12071378>
 70. Lu D, Chatterjee S, Xiao K, Riedel I, Huang CK, Costa A, et al. A circular RNA derived from the insulin receptor locus protects against doxorubicin-induced cardiotoxicity. *Eur Heart J* 2022;**43**:4496–4511. <https://doi.org/10.1093/eurheartj/ehac337>
 71. Rassaf T, Totzeck M, Backs J, Bokemeyer C, Hallek M, Hilfiker-Kleiner D, et al. Onco-cardiology: Consensus paper of the German Cardiac Society, the German Society for Pediatric Cardiology and Congenital Heart Defects and the German Society for Hematology and Medical Oncology. *Clin Res Cardiol* 2020;**109**:1197–1222. <https://doi.org/10.1007/s00392-020-01636-7>
 72. Wojnowski L, Kulle B, Schirmer M, Schlüter I, Kramer A, Rosenberger A, et al. NAD(P)H oxidase and multidrug resistance protein genetic polymorphisms are associated with doxorubicin-induced cardiotoxicity. *Circulation* 2005;**112**:3754–3762. <https://doi.org/10.1161/CIRCULATIONAHA.105.576850>
 73. Kane C, Couch L, Terracciano CM. Excitation-contraction coupling of human induced pluripotent stem cell-derived cardiomyocytes. *Front Cell Dev Biol* 2015;**3**:59. <https://doi.org/10.3389/fcell.2015.00059>
 74. Biley JM, Vermeer M, Duffy, RM, Batalov I, Kramer D, Tashman JW, et al. Dynamic loading of human engineered heart tissue enhances contractile function and drives a desmosome-linked disease phenotype. *Sci Transl Med* 2021;**13**:1–15. <https://doi.org/10.1126/scitranslmed.abd1817>
 75. Tiburcy M, Hudson JE, Balfanz P, Schlick S, Meyer T, Chang Liao ML, et al. Defined engineered human myocardium with advanced maturation for applications in heart failure modeling and repair. *Circulation* 2017;**135**:1832–1847. <https://doi.org/10.1161/CIRCULATIONAHA.116.024145>
 76. Linders AN, Dias IB, Ovchinnikova ES, Vermeer MCSC, Hoes MF, Markousis Mavrogenis G, et al. Evaluation of senescence and its prevention in doxorubicin-induced cardiotoxicity using dynamic engineered heart tissues. *JACC CardioOncol* 2023;**5**:298–315. <https://doi.org/10.1016/j.jacc.2023.03.012>
 77. Long C, Guo R, Han R, Li K, Wan Y, Xu J, et al. Effects of macrophages on the proliferation and cardiac differentiation of human induced pluripotent stem cells. *Cell Commun Signal* 2022;**20**:108. <https://doi.org/10.1186/s12964-022-00916-1>
 78. Emanuelli G, Zoccarato A, Reumiller CM, Papadopoulos A, Chong M, Rebs S, et al. A roadmap for the characterization of energy metabolism in human cardiomyocytes derived from induced pluripotent stem cells. *J Mol Cell Cardiol* 2022;**164**:136–147. <https://doi.org/10.1016/j.yjmcc.2021.12.001>
 79. Ronaldson-Bouchard K, Ma SP, Yeager K, Chen T, Song LJ, Sirabella D, et al. Advanced maturation of human cardiac tissue grown from pluripotent stem cells. *Nature* 2018;**556**:239–243. <https://doi.org/10.1038/s41586-018-0016-3>
 80. Gintant G, Burridge P, Gepstein L, Harding S, Herron T, Hong C, et al. Use of human induced pluripotent stem cell-derived cardiomyocytes in preclinical cancer drug cardiotoxicity testing: A scientific statement from the American Heart Association. *Circ Res* 2019;**125**:e75–e92. <https://doi.org/10.1161/RES.0000000000000291>
 81. Lyra-Leite DM, Copley RR, Freeman PP, Pongpamorn P, Shah D, McKenna DE, et al. Nutritional requirements of human induced pluripotent stem cells. *Stem Cell Reports* 2023;**18**:1371–1387. <https://doi.org/10.1016/j.stemcr.2023.05.004>

82. Lyra-Leite DM, Fonoudi H, Gharib M, Burridge PW. An updated protocol for the cost-effective and weekend-free culture of human induced pluripotent stem cells. *STAR Protoc* 2021;**2**:100213. <https://doi.org/10.1016/j.xpro.2020.100213>
83. Ou Q, Jacobson Z, Abouleisa RRE, Tang XL, Hindi SM, Kumar A, et al. Physiological biomimetic culture system for pig and human heart slices. *Circ Res* 2019;**125**:628–642. <https://doi.org/10.1161/CIRCRESAHA.119.314996>
84. Shi R, Reichardt M, Fiegle DJ, K upfer LK, Czajka T, Sun Z, et al. Contractility measurements for cardiotoxicity screening with ventricular myocardial slices of pigs. *Cardiovasc Res* 2023;**119**:2469–2481. <https://doi.org/10.1093/cvr/cvad141>
85. Boukhalifa A, Robinson SR, Meola DM, Robinson NA, Ling LA, LaMastro JN, et al. Using cultured canine cardiac slices to model the autophagic flux with doxorubicin. *PLoS One* 2023;**18**:e0282859. <https://doi.org/10.1371/journal.pone.0282859>
86. Miller JM, Meki MH, Ou Q, George SA, Gams A, Abouleisa RRE, et al. Heart slice culture system reliably demonstrates clinical drug-related cardiotoxicity. *Toxicol Appl Pharmacol* 2020;**406**:115213. <https://doi.org/10.1016/j.taap.2020.115213>
87. Liu Y, Asnani A, Zou L, Bentley VL, Yu M, Wang Y, et al. Visnagin protects against doxorubicin-induced cardiomyopathy through modulation of mitochondrial malate dehydrogenase. *Sci Transl Med* 2014;**6**:266ra170. <https://doi.org/10.1126/scitranslmed.3010189>
88. Zhu JJ, Xu YQ, He JH, Yu HP, Huang CJ, Gao JM, et al. Human cardiotoxic drugs delivered by soaking and microinjection induce cardiovascular toxicity in zebrafish. *J Appl Toxicol* 2014;**34**:139–148. <https://doi.org/10.1002/jat.2843>
89. Cheng H, Kari G, Dicker AP, Rodeck U, Koch WJ, Force T. A novel preclinical strategy for identifying cardiotoxic kinase inhibitors and mechanisms of cardiotoxicity. *Circ Res* 2011;**109**:1401–1409. <https://doi.org/10.1161/CIRCRESAHA.111.255695>
90. Singh AP, Glennon MS, Umbarkar P, Gupta M, Galindo CL, Zhang Q, et al. Ponatinib-induced cardiotoxicity: Delineating the signalling mechanisms and potential rescue strategies. *Cardiovasc Res* 2019;**115**:966–977. <https://doi.org/10.1093/cvr/cvz006>
91. Sawicki KT, Sala V, Prever L, Hirsch E, Ardehali H, Ghigo A. Preventing and treating anthracycline cardiotoxicity: New insights. *Annu Rev Pharmacol Toxicol* 2021;**61**:309–332. <https://doi.org/10.1146/annurev-pharmtox-030620-104842>
92. Lewis Phillips G, Guo J, Kiefer JR, Proctor W, Bumbaca Yadav D, Dybdal N, et al. Trastuzumab does not bind rat or mouse ErbB2/neu: Implications for selection of non-clinical safety models for trastuzumab-based therapeutics. *Breast Cancer Res Treat* 2022;**191**:303–317. <https://doi.org/10.1007/s10549-021-06427-w>
93. Yong CS, Sharkey J, Duscio B, Yong CSM, Venville B, Wei WZ, et al. Embryonic lethality in homozygous human Her-2 transgenic mice due to disruption of the Pds5b gene. *PLoS One* 2015;**10**:e0136817. <https://doi.org/10.1371/journal.pone.0136817>
94. Michel L, Helfrich I, Hendgen-Cotta UB, Mincu RI, Korste S, Mroczek SM, et al. Targeting early stages of cardiotoxicity from anti-PD1 immune checkpoint inhibitor therapy. *Eur Heart J* 2021;**43**:316–329. <https://doi.org/10.1093/eurheartj/ehab430>
95. Liu X, Wu W, Fang L, Liu Y, Chen W. TNF-alpha inhibitors and other biologic agents for the treatment of immune checkpoint inhibitor-induced myocarditis. *Front Immunol* 2022;**13**:922782. <https://doi.org/10.3389/fimmu.2022.922782>
96. Amgalan D, Garner TP, Pekson R, Jia XF, Yanamandala M, Paulino V, et al. A small-molecule allosteric inhibitor of BAX protects against doxorubicin-induced cardiomyopathy. *Nat Cancer* 2020;**1**:315–328. <https://doi.org/10.1038/s43018-020-0039-1>
97. Li M, Sala V, De Santis MC, Cimino J, Cappello P, Pianca N, et al. Phosphoinositide 3-kinase gamma inhibition protects from anthracycline cardiotoxicity and reduces tumor growth. *Circulation* 2018;**138**:696–711. <https://doi.org/10.1161/CIRCULATIONAHA.117.030352>
98. van der Velden J, Asselbergs FW, Bakkers J, Batkai S, Bertrand L, Bezzina CR, et al. Animal models and animal-free innovations for cardiovascular research: Current status and routes to be explored. Consensus document of the ESC Working Group on Myocardial Function and the ESC Working Group on Cellular Biology of the Heart. *Cardiovasc Res* 2022;**118**:3016–3051. <https://doi.org/10.1093/cvr/cvab370>
99. Bartoli CR, Brittan KR, Giridharan GA, Koenig SC, Hamid T, Prabhu SD. Bovine model of doxorubicin-induced cardiomyopathy. *J Biomed Biotechnol* 2011;**2011**:758736. <https://doi.org/10.1155/2011/758736>
100. Carbone A, Psaltis PJ, Nelson AJ, Metcalf R, Richardson JD, Weightman M, et al. Dietary omega-3 supplementation exacerbates left ventricular dysfunction in an ovine model of anthracycline-induced cardiotoxicity. *J Card Fail* 2012;**18**:502–511. <https://doi.org/10.1016/j.cardfail.2012.03.005>
101. Engwall MJ, Evers N, Turk JR, Vargas HM. The effects of repeat-dose doxorubicin on cardiovascular functional endpoints and biomarkers in the telemetry-equipped Cynomolgus monkey. *Front Cardiovasc Med* 2021;**8**:587149. <https://doi.org/10.3389/fcvm.2021.587149>
102. Galan-Arriola C, Lobo M, Vilchez-Tschischke JP, L opez GJ, de Molina-Iracheta A, P erez-Martinez C, et al. Serial magnetic resonance imaging to identify early stages of anthracycline-induced cardiotoxicity. *J Am Coll Cardiol* 2019;**73**:779–791. <https://doi.org/10.1016/j.jacc.2018.11.046>
103. Galan-Arriola C, Vilchez-Tschischke JP, Lobo M, L opez GJ, de Molina-Iracheta A, P erez-Martinez C, et al. Coronary microcirculation damage in anthracycline cardiotoxicity. *Cardiovasc Res* 2022;**118**:531–541. <https://doi.org/10.1093/cvr/cvab053>
104. Melendez GC, Vasu S, Lesnfsky EJ, Kaplan JR, Appt S, D'Agostino RB Jr, et al. Myocardial extracellular and cardiomyocyte volume expand after doxorubicin treatment similar to adjuvant breast cancer therapy. *JACC Cardiovasc Imaging* 2020;**13**:1084–1085. <https://doi.org/10.1016/j.jcmg.2019.10.020>
105. Muehlberg F, Funk S, Zange L, von Knobelsdorff-Brenkenhoff F, Blaszczyk E, Schulz A, et al. Native myocardial T1 time can predict development of subsequent anthracycline-induced cardiomyopathy. *ESC Heart Fail* 2018;**5**:620–629. <https://doi.org/10.1002/ehf2.12277>
106. Giusca S, Korosoglou G, Montenbruck M, Ger ak B, Schwarz AK, Esch S, et al. Multiparametric early detection and prediction of cardiotoxicity using myocardial strain, T1 and T2 mapping, and biochemical markers: A longitudinal cardiac resonance imaging study during 2 years of follow-up. *Circ Cardiovasc Imaging* 2021;**14**:e012459. <https://doi.org/10.1161/CIRCIMAGING.121.012459>
107. Martin-Garcia A, Diaz-Pelaez E, Lopez-Corral L, Sanchez-Pablo C, Macias de Plasencia G, Galan-Arriola C, et al. T2 mapping identifies early anthracycline-induced cardiotoxicity in elderly patients with cancer. *JACC Cardiovasc Imaging* 2020;**13**:1630–1632. <https://doi.org/10.1016/j.jcmg.2020.01.017>
108. Galan-Arriola C, Villena-Gutierrez R, Higuero-Verdejo MI, Diaz-Rengifo IA, Pizarro G, L opez GJ, et al. Remote ischaemic preconditioning ameliorates anthracycline-induced cardiotoxicity and preserves mitochondrial integrity. *Cardiovasc Res* 2021;**117**:1132–1143. <https://doi.org/10.1093/cvr/cvaa181>
109. Aboumsallem JP, Moslehi J, de Boer RA. Reverse cardio-oncology: Cancer development in patients with cardiovascular disease. *J Am Heart Assoc* 2020;**9**:e013754. <https://doi.org/10.1161/JAHA.119.013754>
110. Bertero E, Robusto F, Rulli E, D'Etorre A, Bisceglia L, Staszewsky L, et al. Cancer incidence and mortality according to pre-existing heart failure in a community-based cohort. *JACC CardioOncol* 2022;**4**:98–109. <https://doi.org/10.1016/j.jacc.2021.11.007>
111. Jaiswal V, Ang SP, Agrawal V, Hameed M, Saleeb MRA, Jaiswal A, et al. Association between heart failure and the incidence of cancer: A systematic review and meta-analysis. *Eur Heart J Open* 2023;**3**:oead073. <https://doi.org/10.1093/ehjopen/oead073>
112. de Wit S, Aboumsallem JP, Shi C, Schouten EM, Bracon V, Meijers WC, et al. Pressure overload-induced cardiac hypertrophy stimulates tumor growth in tumor-prone Apc^{Min} mice. *Circ Heart Fail* 2023;**16**:e010740. <https://doi.org/10.1161/CIRCHEARTFAILURE.123.010740>
113. Meijers WC, Maglione M, Bakker SJL, Oberhuber R, Kieneker LM, de Jong S, et al. Heart failure stimulates tumor growth by circulating factors. *Circulation* 2018;**138**:678–691. <https://doi.org/10.1161/CIRCULATIONAHA.117.030816>
114. Avraham S, Abu-Sharki S, Shofti R, Haas T, Korin B, Kalfon R, et al. Early cardiac remodeling promotes tumor growth and metastasis. *Circulation* 2020;**142**:670–683. <https://doi.org/10.1161/CIRCULATIONAHA.120.046471>
115. Langier Goncalves I, Awwad L, Aviram S, Izraeli T, Achlaug L, Aronheim A. Heart failure promotes cancer progression in an integrin β 1-dependent manner. *Int J Mol Sci* 2023;**24**:17367. <https://doi.org/10.3390/ijms242417367>
116. de Wit W, Geerlings L, Shi C, Dronkers J, Schouten EM, Blanche G, et al. Heart failure-induced microbial dysbiosis contributes to colonic tumour formation in mice. *Cardiovasc Res* 2024;**120**:612–622. <https://doi.org/10.1093/cvr/cvae038>
117. Tani T, Oikawa M, Misaka T, Ishida T, Takeishi Y. Heart failure post-myocardial infarction promotes mammary tumor growth through the NGF-TRKA pathway. *JACC CardioOncol* 2024;**6**:55–66. <https://doi.org/10.1016/j.jacc.2023.10.002>
118. Koelwyn GJ, Newman AAC, van Afonso MS, Solingen C, Corr EM, Brown EJ, et al. Myocardial infarction accelerates breast cancer via innate immune reprogramming. *Nat Med* 2020;**26**:1452–1458. <https://doi.org/10.1038/s41591-020-0964-7>
119. Shi C, Aboumsallem JP, de Wit S, Schouten EM, Bracon V, Meijers WC, et al. Evaluation of renal cancer progression in a mouse model of heart failure. *Cancer Commun (Lond)* 2021;**41**:796–799. <https://doi.org/10.1002/cac2.12185>
120. Awwad L, Aronheim A. Cardiac dysfunction promotes cancer progression via multiple secreted factors. *Cancer Res* 2022;**82**:1753–1761. <https://doi.org/10.1158/0008-5472.CAN-21-2463>
121. Awwad L, Goldenberg T, Langier-Goncalves I, Aronheim A. Cardiac remodeling in the absence of cardiac contractile dysfunction is sufficient to promote cancer progression. *Cells* 2022;**11**:1108. <https://doi.org/10.3390/cells11071108>

122. Koelwyn GJ, Aboumsallem JP, Moore KJ, de Boer RA. Reverse cardio-oncology: Exploring the effects of cardiovascular disease on cancer pathogenesis. *J Mol Cell Cardiol* 2022;163:1–8. <https://doi.org/10.1016/j.yjmcc.2021.09.008>
123. Bertero E, Carmisciano L, Jonasson C, Butler J, Maack C, Ameri P. Association of inflammatory markers with incident heart failure or cancer in the HUNT3 and Health ABC population studies. *Eur J Prev Cardiol* 2024;31:1400–1407. <https://doi.org/10.1093/eurjpc/zwae089>
124. Jovani M, Liu EE, Paniagua SM, Lau ES, Li SX, Takvorian KS, et al. Cardiovascular disease related circulating biomarkers and cancer incidence and mortality: Is there an association? *Cardiovasc Res* 2022;118:2317–2328. <https://doi.org/10.1093/cvr/cvab282>
125. Meijers WC, de Boer RA. Common risk factors for heart failure and cancer. *Cardiovasc Res* 2019;115:844–853. <https://doi.org/10.1093/cvr/cvz035>
126. Garlanda C, Mantovani A. Interleukin-1 in tumor progression, therapy, and prevention. *Cancer Cell* 2021;39:1023–1027. <https://doi.org/10.1016/j.ccell.2021.04.011>
127. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, et al.; CANTOS Trial Group. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 2017;377:1119–1131. <https://doi.org/10.1056/NEJMoa1707914>
128. Ridker PM, MacFadyen JG, Thuren T, B. M. Everett P, Libby R, J. Glynn, et al.; CANTOS Trial Group. Effect of interleukin-1 β inhibition with canakinumab on incident lung cancer in patients with atherosclerosis: Exploratory results from a randomised, double-blind, placebo-controlled trial *Lancet* 2017 **390** 1833–1842. [https://doi.org/10.1016/S0140-6736\(17\)32247-X](https://doi.org/10.1016/S0140-6736(17)32247-X) 10105
129. Wong CC, Baum J, Silvestro A, Beste MT, Bharani-Dharan B, Xu S, et al. Inhibition of IL1 β by canakinumab may be effective against diverse molecular subtypes of lung cancer: An exploratory analysis of the CANTOS trial. *Cancer Res* 2020;80:5597–5605. <https://doi.org/10.1158/0008-5472.CAN-19-3176>
130. Garon EB, Lu S, Goto Y, De Marchi P, Paz-Ares L, Spigel DR, et al. Canakinumab as adjuvant therapy in patients with completely resected non-small-cell lung cancer: Results from the CANOPY-A double-blind, randomized clinical trial. *J Clin Oncol* 2024;42:180–191. <https://doi.org/10.1200/JCO.23.00910>
131. Paik S, Kim JK, Silwal P, Sasakawa C, Jo EK. An update on the regulatory mechanisms of NLRP3 inflammasome activation. *Cell Mol Immunol* 2021;18:1141–1160. <https://doi.org/10.1038/s41423-021-00670-3>
132. Nidorf SM, Fiolet ATL, Mosterd A, Eikelboom JW, Schut A, Opstal TSJ, et al.; LoDoCo2 Trial Investigators. Colchicine in patients with chronic coronary disease. *N Engl J Med* 2020;383:1838–1847. <https://doi.org/10.1056/NEJMoa2021372>
133. Pappritz K, Lin J, El-Shafeey M, Fechner H, Kühl U, Alogna A, et al. Colchicine prevents disease progression in viral myocarditis via modulating the NLRP3 inflammasome in the cardiopleic axis. *ESC Heart Fail* 2022;9:925–941. <https://doi.org/10.1002/ehf2.13845>
134. Tardif JC, Kouz S, Waters DD, Bertrand OF, Diaz R, Maggioni AP, et al. Efficacy and safety of low-dose colchicine after myocardial infarction. *N Engl J Med* 2019;381:2497–2505. <https://doi.org/10.1056/NEJMoa1912388>
135. He F, Xie T, Ni D, Tang T, Cheng X. Efficacy and safety of inhibiting the NLRP3/IL-1beta/IL-6 pathway in patients with ST-elevation myocardial infarction: A meta-analysis. *Eur J Clin Invest* 2023;53:e14062. <https://doi.org/10.1111/eci.14062>
136. Olsen MB, Gregersen I, Sandanger O, Yang K, Sokolova M, Halvorsen BE, et al. Targeting the inflammasome in cardiovascular disease. *JACC Basic Transl Sci* 2022;7:84–98. <https://doi.org/10.1016/j.jacbts.2021.08.006>
137. Shadab A, Mahjoor M, Abbasi-Kolli M, Afkhami H, Moenian P, Safdarian AR. Divergent functions of NLRP3 inflammasomes in cancer: A review. *Cell Commun Signal* 2023;21:232. <https://doi.org/10.1186/s12964-023-01235-9>
138. Van Linthout S, Volk HD. Immuno-cardio-oncology: Killing two birds with one stone? *Front Immunol* 2022;13:1018772. <https://doi.org/10.3389/fimmu.2022.1018772>
139. Muller I, Vogl T, Kuhl U, Krannich A, Banks A, Trippel T, et al. Serum alarmin S100A8/S100A9 levels and its potential role as biomarker in myocarditis. *ESC Heart Fail* 2020;7:1442–1451. <https://doi.org/10.1002/ehf2.12760>
140. Muller I, Vogl T, Pappritz K, Miteva K, Savvatis K, Rohde D, et al. Pathogenic role of the damage-associated molecular patterns S100A8 and S100A9 in Coxsackievirus B3-induced myocarditis. *Circ Heart Fail* 2017;10. <https://doi.org/10.1161/CIRCHEARTFAILURE.117.004125>, 10
141. Low D, Subramaniam R, Lin L, Aomatsu T, Mizoguchi A, Ng A, et al. Chitinase 3-like 1 induces survival and proliferation of intestinal epithelial cells during chronic inflammation and colitis-associated cancer by regulating S100A9. *Oncotarget* 2015;6:36535–36550. <https://doi.org/10.18632/oncotarget.5440>
142. Marinkovic G, Koenis DS, de Camp L, Jablonowski R, Graber N, de Waard V, et al. S100A9 links inflammation and repair in myocardial infarction. *Circ Res* 2020;127:664–676. <https://doi.org/10.1161/CIRCRESAHA.120.315865>
143. Papp Z, Pollesello P, Borbely A. Recognizing early signs in the translational phase is essential for drug development in cardiovascular medicine. *ESC Heart Fail* 2023;10:2147–2149. <https://doi.org/10.1002/ehf2.14402>
144. Li DL, Wang ZV, Ding G, Tan W, Luo X, Criollo A, et al. Doxorubicin blocks cardiomyocyte autophagic flux by inhibiting lysosome acidification. *Circulation* 2016;133:1668–1687. <https://doi.org/10.1161/CIRCULATIONAHA.115.017443>
145. Robinson EL, Ameri P, Delrue L, Vanderheyden M, Bartunek J, Altieri P, et al. Differential expression of epigenetic modifiers in early and late cardiotoxic heart failure reveals DNA methylation as a key regulator of cardiotoxicity. *Front Cardiovasc Med* 2023;10:884174. <https://doi.org/10.3389/fcvm.2023.884174>
146. Zhou X, Liu Y, Shen Y, Chen L, Hu W, Yan Y, et al. Rescue of cardiac dysfunction during chemotherapy in acute myeloid leukaemia by blocking IL-1 α . *Eur Heart J* 2024;45:2235–2250. <https://doi.org/10.1093/eurheartj/ehae188>
147. Lee J, Mehrotra S, Zare-Eelanjegh E, Rodrigues R, Akbarinejad A, Ge D, et al. A heart-breast cancer-on-a-chip platform for disease modeling and monitoring of cardiotoxicity induced by cancer chemotherapy. *Small* 2021;17:e2004258. <https://doi.org/10.1002/sml.202004258>
148. Nickel AG, von Hardenberg A, Hohl M, Nickel A G, Löffler JR, Kohlhaas M, et al. Reversal of mitochondrial transhydrogenase causes oxidative stress in heart failure. *Cell Metab* 2015;22:472–484. <https://doi.org/10.1016/j.cmet.2015.07.008>
149. Zeiss CJ, Gatti DM, Toro-Salazar O, Davis C, Lutz CM, Spinale F, et al. Doxorubicin-induced cardiotoxicity in collaborative cross (CC) mice recapitulates individual cardiotoxicity in humans. *G3 (Bethesda)* 2019;9:2637–2646. <https://doi.org/10.1534/g3.119.400232>
150. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: The ARRIVE guidelines for reporting animal research. *PLoS Biol* 2010;8:e1000412. <https://doi.org/10.1371/journal.pbio.1000412>
151. Wilcox NS, Rotz SJ, Mullen M, Song EJ, Ky Hamilton B, Moslehi J, et al. Sex-specific cardiovascular risks of cancer and its therapies. *Circ Res* 2022;130:632–651. <https://doi.org/10.1161/CIRCRESAHA.121.319901>
152. Rosano GMC, Stolfo D, Anderson L, Abdelhamid M, Adamo M, Bauersachs J, et al. Differences in presentation, diagnosis and management of heart failure in women. A scientific statement of the Heart Failure Association of the ESC. *Eur J Heart Fail* 2024;26:1669–1686. <https://doi.org/10.1002/ehf.3284>
153. Waldhorn I, Turetsky T, Steiner D, Gil Y, Benyamini H, Gropp M, et al. Modeling sex differences in humans using isogenic induced pluripotent stem cells. *Stem Cell Reports* 2022;17:2732–2744. <https://doi.org/10.1016/j.stemcr.2022.10.017>
154. Pinheiro EA, Magdy T, Burrige PW. Human in vitro models for assessing the genomic basis of chemotherapy-induced cardiovascular toxicity. *J Cardiovasc Transl Res* 2020;13:377–389. <https://doi.org/10.1007/s12265-020-09962-x>
155. Balaji S, Antony AK, Tonchev H, Scichilone G, Morsy M, Deen H, et al. Racial disparity in anthracycline-induced cardiotoxicity in breast cancer patients. *Biomedicine* 2023;11:2286. <https://doi.org/10.3390/biomed11082286>
156. D'Arienzo A, Verrazzo A, Pagliuca M, D'Arienzo A, Napolitano F, Parola S, et al. Toxicity profile of antibody-drug conjugates in breast cancer: Practical considerations. *EClinicalMedicine* 2023;62:102113. <https://doi.org/10.1016/j.eclinm.2023.102113>
157. Ursini-Siegel J, Schade B, Cardiff RD, Muller WJ. Insights from transgenic mouse models of ERBB2-induced breast cancer. *Nat Rev Cancer* 2007;7:389–397. <https://doi.org/10.1038/nrc2127>
158. Hoopes PJ, Tavakkoli AD, Moodie KA, Maurer KJ, Meehan KR, Wallin DJ, et al. Porcine-human glioma xenograft model. Immunosuppression and model reproducibility. *Cancer Treat Res Commun* 2024;38:100789. <https://doi.org/10.1016/j.ctarc.2024.100789>
159. Farmakis D, Papingiotis G, Filippos G. Genetic predisposition to cardiovascular disease in patients with cancer: A clinical perspective. *JACC CardioOncol* 2023;5:402–405. <https://doi.org/10.1016/j.jaccao.2022.12.008>
160. Fonoudi H, Jouni M, Cejas RB, Magdy T, Blancard M, Ge N, et al. Functional validation of doxorubicin-induced cardiotoxicity-related genes. *JACC CardioOncol* 2024;6:38–50. <https://doi.org/10.1016/j.jaccao.2023.11.008>
161. Magdy T, Jiang Z, Jouni M, Fonoudi H, Lyra-Leite D, Jung G, et al. RARG variant predictive of doxorubicin-induced cardiotoxicity identifies a cardioprotective therapy. *Cell Stem Cell* 2021;28:2076–2089. <https://doi.org/10.1016/j.stem.2021.08.006>
162. Magdy T, Jouni M, Kuo HH, Weddle CJ, Lyra-Leite D, Fonoudi H, et al. Identification of drug transporter genomic variants and inhibitors that protect against doxorubicin-induced cardiotoxicity. *Circulation* 2022;145:279–294. <https://doi.org/10.1161/CIRCULATIONAHA.121.055801>
163. Van Norman GA. Overcoming the declining trends in innovation and investment in cardiovascular therapeutics: Beyond EROOM's law. *JACC Basic Transl Sci* 2017;2:613–625. <https://doi.org/10.1016/j.jacbts.2017.09.002>

164. Ringel MS, Shah NA, Baedeker M, Lim CT, Lamichhane A, Schulze U. Occlusion in the flow of new drugs for cardiovascular disease. *Clin Pharmacol Ther* 2017;**102**:246–253. <https://doi.org/10.1002/cpt.691>
165. Prescott JS, Andrews PA, Baker RW, Bogdanffy MS, Fields FO, Keller DA, et al. Evaluation of therapeutics for advanced-stage heart failure and other severely-debilitating or life-threatening diseases. *Clin Pharmacol Ther* 2017;**102**:219–227. <https://doi.org/10.1002/cpt.730>
166. Walker, N.. Drug Approval Trends: Significant Acceleration in Recent Years. July 1, 2020. <https://www.pharmasalmanc.com/articles/drug-approval-trends-significant-acceleration-in-recent-years>. Accessed October 28, 2024.
167. Romero-Sanchez A, Perdomo-Charry G, Burbano-Vallejo EL. Exploring the entrepreneurial landscape of university-industry collaboration on public university spin-off creation: A systematic literature review. *Heliyon* 2024;**10**:e27258. <https://doi.org/10.1016/j.heliyon.2024.e27258>
168. Sanberg PR, Gharib M, Harker PT, Kaler EVV, Marchase RB, Sands TD, et al. Changing the academic culture: Valuing patents and commercialization toward tenure and career advancement. *Proc Natl Acad Sci U S A* 2014;**111**:6542–6547. <https://doi.org/10.1073/pnas.1404094111>
169. Fradley MG, Nguyen NHK, Madnick D, Chen Y, DeMichele A, Makhlin I, et al. Adverse cardiovascular events associated with cyclin-dependent kinase 4/6 inhibitors in patients with metastatic breast cancer. *J Am Heart Assoc* 2023;**12**:e029361. <https://doi.org/10.1161/JAHA.123.029361>
170. Ferrara AL, Loffredo S, Tocchetti CG. Inflammation in cardio-oncology: Beyond immunotherapies. *Eur Heart J* 2024;**45**:2251–2253. <https://doi.org/10.1093/eurheartj/ehae333>
171. Matz I, Pappritz K, Springer J, Van Linthout S. Left ventricle- and skeletal muscle-derived fibroblasts exhibit a differential inflammatory and metabolic responsiveness to interleukin-6. *Front Immunol* 2022;**13**:947267. <https://doi.org/10.3389/fimmu.2022.947267>
172. Sayed N, Ameen M, Wu JC. Personalized medicine in cardio-oncology: The role of induced pluripotent stem cell. *Cardiovasc Res* 2019;**115**:949–959. <https://doi.org/10.1093/cvr/cvz024>