

Supporting information

S2 Table. Evaluation of the haematological and biochemical parameters measured in the relapsed patients and the rest of the mITT set at baseline.

Parameter	Relapsed patients	Non-relapsed patients	<i>p</i> -value relapsed vs non-relapsed
Albumin g/L (Low:30;Upper:50)	39.30 ± 5.86	39.13 ± 4.27	0.77
Creatinine mg/dL (Low:0.2;Upper:1.4)	0.4 ± 0.14	0.41 ± 0.20	0.94
Pottasium mmol/L (Low:3;Upper:5)	4.14 ± 0.32	3.97 ± 0.33	0.28
SGOT/AST U/L (Upper:<=40)	35 ± 4.06	30.91 ± 10.18	0.12
SGPT/ALT U/L (Upper:<=40)	25.8 ± 9.44	26.57 ± 12.53	0.99
Total Bilirubin mg/dL (Upper:<=1.2)	0.40 ± 0.24	0.48 ± 0.29	0.61
Basophils %	0 ± 0	0 ± 0	ND
Eosinophils %	2.60 ± 2.51	2.41 ± 1.56	0.69
Haematocrit %	36 ± 2.24	36.94 ± 3.57	0.47
Haemoglobin g/dL (Low:12;Upper:17.5)	11.56 ± 1.20	12.45 ± 1.26	0.13

Lymphocytes %	45 ± 14.40	45.58 ± 8.16	0.88
Monocytes %	7 ± 3.24	7.87 ± 2.83	0.68
Neutrophil %	45.20 ± 9.58	44.15 ± 8.92	0.79
Platelets x10³/μL (Low:150;Upper:500)	345 ± 111.88	341.40 ± 95.17	0.60
RBC x 10⁶/μL	4.74 ± 0.38	4.68 ± 0.47	0.61
WBC x 10⁶/μL (Low:4; Upper:10)	7.46 ± 1.07	7.14 ± 3.47	0.30

Data reported is presented as concentrations or percentages of each parameter. Statistical differences were evaluated by Mann-Whitney test.